

A DEGENERATION FORMULA OF GW-INVARIANTS

JUN LI

ABSTRACT. This is the sequel to the paper [Li]. In this paper, we construct the virtual moduli cycles of the degeneration of the moduli of stable morphisms constructed in [Li]. We also construct the virtual moduli cycles of the moduli of relative stable morphisms of a pair of a smooth divisor in a smooth variety. Based on these, we prove a degeneration formula of the Gromov-Witten invariants.

0. INTRODUCTION

This is the second part of the project initiated in [Li].

Like Donaldson invariants of 4-manifolds, Gromov-Witten invariants are intersection theories on the moduli spaces of stable morphisms to varieties or symplectic manifolds. However, for Gromov-Witten invariants one needs to use the virtual intersection theories to define the Gromov-Witten invariants. Namely the intersection theories defined based on the virtual moduli cycles. Such cycles were first constructed by Tian and the author [LT1, LT2] for algebraic varieties, and an alternative construction was achieved by Behrend-Fantechi [Beh, BF]. For Gromov-Witten invariants of general symplectic manifolds see [Ru1, RT, FO, LT3, Ru2, Si1] and the equivalence of these constructions see [LT4, Si2, KKP].

The goal of this project is to investigate the structure of the Gromov-Witten invariants when X degenerates to a singular scheme. Specifically, we will prove a degeneration formula of the Gromov-Witten invariants in algebraic geometry. This is the analogous of the Donaldson-Floer theory in gauge theory. Such a degeneration theory was investigated by several groups using analysis [EGH, IP1, IP2, LR, Tia].

Here is the situation we will study in this paper. Let $W \rightarrow C$ be a family of projective schemes over a smooth curve C with $0 \in C$ a distinguished point so that the fibers W_t over $t \neq 0 \in C$ are smooth varieties and the special fiber W_0 has two smooth irreducible components Y_1 and Y_2 intersecting transversally along a connected smooth divisor $D \in W_0$. We will call $Y_i^{\text{rel}} \triangleq (Y_i, D_i)$, where $D_i = D \subset Y_i$, the relative pairs after decomposing W_0 . In this paper, we will construct the Gromov-Witten invariants of W_t for all $t \in C$; We will construct the relative Gromov-Witten invariants of any relative pair $Z^{\text{rel}} \triangleq (Z, D)$ of a smooth projective variety and a smooth divisor; We will prove that the Gromov-Witten invariants of W_t are locally constant for $t \in C$; Finally we will prove a degeneration formula relating the Gromov-Witten invariants of W_0 with the relative Gromov-Witten invariants of Y_1^{rel} and Y_2^{rel} , one in cycle form and the other in numerical form.

Date: October 1, 2001.

Supported partially by NSF grants.

The first part of this paper is devoted to construct the Gromov-Witten invariants of the singular scheme W_0 . To achieve this, we need to construct a new relative moduli space of stable morphisms to W/C . This is achieved by refining the notion of stable morphisms, worked out in the first part of this project [Li]. Recall that in [Li] we constructed a stack $\mathfrak{W}$ that includes all expanded degenerations of W . We then introduced the notion of pre-stable, pre-deformable and stable morphisms to $\mathfrak{W}$. For simplicity, we denote by Γ the triple consisting of the genus, the number of marked points and the degree of the stable morphisms. The main result of [Li] is the existence of the moduli space $\mathfrak{M}(\mathfrak{W}, \Gamma)$ of stable morphisms to $\mathfrak{W}$ of topological type Γ as a Deligne-Mumford stack, separated and proper over C . Applying parallel construction to a relative pair Z^{rel} we constructed the stack of expanded relative pairs $\mathfrak{Z}^{\text{rel}}$. We then defined the notion of relative stable morphisms to $\mathfrak{Z}^{\text{rel}}$ and showed that the moduli $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ of relative stable morphisms to $\mathfrak{Z}^{\text{rel}}$ with topological type¹ Γ is also a separated and proper Deligne-Mumford stack.

In this paper, we first construct the standard obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$, $\mathfrak{M}(\mathfrak{W}_0, \Gamma) = \mathfrak{M}(\mathfrak{W}, \Gamma) \times_C 0$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$. We then show that they are all perfect, thus allow us to construct their respective virtual moduli cycles. Based on the virtual moduli cycle $[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}}$, we define the Gromov-Witten invariants of W_0

$$\Psi_{\Gamma}^{W_0} : H^*(W_0)^{\times k} \times H^*(\mathfrak{M}_{g,k}) \longrightarrow \mathbb{Q}$$

in the standard way, where $\Gamma = (g, k, A)$. Similarly, for Z^{rel} we define its relative Gromov-Witten invariants

$$\Psi_{\Gamma}^{Z^{\text{rel}}} : H^*(Z)^{\times k} \times H^*(\mathfrak{M}_{\Gamma^{\circ}}) \longrightarrow H_*(D^r)$$

to be

$$\Psi_{\Gamma}^{Z^{\text{rel}}}(\alpha, \beta) = \mathbf{q}_*(\text{ev}^*(\alpha) \cup \pi_{\Gamma}^*(\beta) [\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)]^{\text{virt}}) \in H_*(D^r).$$

Here k (resp. r) is the number of ordinary (resp. distinguished) marked points of the domain curves, ev is the evaluation morphism associated to the ordinary marked points and $\mathbf{q} : \mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma) \rightarrow D^r$ is the evaluation associated to the distinguished marked points. The symbol Γ is the topological type of the relative stable morphisms and $\pi_{\Gamma} : \mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma) \rightarrow \mathfrak{M}_{\Gamma^{\circ}}$ is the forgetful morphism, where $\mathfrak{M}_{\Gamma^{\circ}}$ is the moduli of stable nodal curves² whose topology is given by the data in Γ .

The invariants $\Phi_{\Gamma}^{W_0}$ and $\Phi_{\Gamma'}^{Z^{\text{rel}}}$ have the expected properties. For instance, $\Phi_{\Gamma}^{W_t}$ is locally constant for $t \in C$, and $\Phi_{\Gamma'}^{Z^{\text{rel}}}$ is invariant under any smooth deformation of Z^{rel} .

The second part of this paper is to derive a degeneration formula of the Gromov-Witten invariants associated to the (degeneration) family W . As explained in [Li], we expect to have a formula relating the Gromov-Witten invariants of W_0 , and hence of W_t , to the relative Gromov-Witten invariants of Y_i^{rel} . In this paper, we

¹Here the topological type of a relative stable morphism $f : X \rightarrow Z^{\text{rel}}$ is given by a weighted graph Γ . Γ consists of vertices, ordered legs and ordered roots, where each leg and root has one end attached to one vertex. Each vertex represents a connected component of the domain X , each leg (resp. root) represents a marked point (resp. distinguished marked point) on X and all distinguished marked points are mapped to D under f . The vertices and roots of Γ are assigned weights, representing the degree of f along the associated component of X and the contact order of f along the normal direction of $D \subset Z$. Here naturally we insist that f maps all distinguished points to D .

²By which we mean nodal curves having no vector fields, not necessary connected.

prove the following degeneration formula

$$\Phi_{\Gamma}^{W_t}(\alpha(t), \beta) = \sum_{\eta \in \bar{\Omega}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \sum_{j \in K_{\eta}} [\Psi_{\Gamma_1}^{Y^{\text{rel}}} (j_1^* \alpha(0), \beta_{\eta,1,j}) \bullet \Psi_{\Gamma_2}^{Y^{\text{rel}}} (j_2^* \alpha(0), \beta_{\eta,2,j})]_0.$$

We will explain the notation momentarily. We call the above the degeneration formula in numerical form. There is a parallel degeneration formula in cycle form

$$[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}} = \sum_{\eta \in \bar{\Omega}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}).$$

The degeneration formula in numerical form is an easy consequence of the degeneration formula in cycle form.

We now explain the notation in the degeneration formulas. Let Γ_1 be a topological type of a relative stable morphism to Y_1^{rel} . Then by evaluating on the distinguished marked points of the domains of the relative stable morphisms, we obtain the evaluation morphism

$$\mathbf{q}_1 : \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \longrightarrow D^r,$$

where r is the number of *distinguished marked points*. Now let Γ_1 and Γ_2 be so that they have identical number of roots. (Note that roots are associated to the distinguished points of the domains of relative stable morphisms.) Then we have a pair of evaluation morphisms $\mathbf{q}_1$ and $\mathbf{q}_2$ and thus can form the Cartesian product

$$\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2).$$

Let $D^r \rightarrow D^r \times D^r$ be the diagonal morphism. The virtual moduli cycle of the above Cartesian product is

$$\Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}).$$

The set $\bar{\Omega}$ is the set of equivalence classes of all admissible triples (Γ_1, Γ_2, I) . Here Γ_1 and Γ_2 are two weighted graphs associated to two relative stable morphisms. They are required to have identical number of roots and the weights of the i -th roots of Γ_1 and Γ_2 are identical for all i . Hence if (f_1, f_2) be an element in $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$ with X_i the domain of f_i , then we can glue the i -th distinguished marked points (associated to the i -th root) of X_1 with the i -th distinguished marked point of X_2 for all i to obtain a morphism $f : X \rightarrow W_0$. As part of the requirement on the triple (Γ_1, Γ_2, I) the curve X is connected of arithmetic genus g and f is a stable morphism topological type Γ . Lastly, I is a rule concerning the ordering of the union of the ordinary marked points of X_1 and X_2 . Note that in case η has r roots, then any permutation $\sigma \in S_r$ defines a new element η^{σ} by reordering the roots of η according to σ . For $\eta_1, \eta_2 \in \Omega$, we say $\eta_1 \sim \eta_2$ if $\eta_1 = \eta_2^{\sigma}$ for some σ . The set $\bar{\Omega}$ is the set of equivalence classes $\Omega / \sim$. The notation $\text{Eq}(\eta)$ appeared in the degeneration formula is the set of all $\sigma \in S_r$ so that $\eta = \eta^{\sigma}$.

Let $\eta \in \bar{\Omega}$, then the above construction associates to every pair (f_1, f_2) a stable morphism f in $\mathfrak{M}(\mathfrak{W}, \Gamma)$. This defines a morphism

$$\Phi_{\eta} : \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2) \longrightarrow \mathfrak{M}(\mathfrak{W}_0, \Gamma).$$

A Lemma in [Li] asserts that Φ_{η} is a local immersion and the degree of Φ_{η} , as morphism to $\text{Im}(\Phi_{\eta})$, is $|\text{Eq}(\eta)|$. Then the degeneration formula in cycle form

asserts that the union of

$$\Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}})$$

with multiplicity $\mathfrak{m}(\eta)/|\text{Eq}(\eta)|$ is the virtual moduli cycle of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. Here $\mathfrak{m}(\eta)$ is the product of the weights of the roots of η (or of the Γ_1 in η).

We now explain briefly the strategy to prove the degeneration formula. Let $(\mathbf{1}, \mathbf{t})$ be the pair of the trivial line bundle on $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and the pull back of a section $t \in \Gamma(\mathcal{O}_C)$ so that $t^{-1}(0)$ is the origin $0 \in C$. Then the virtual moduli cycle $[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}}$ is the intersection

$$[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}} \cap \mathbf{t}^{-1}(0) \in A_* \mathfrak{M}(\mathfrak{W}_0, \Gamma).$$

Using the notion of localized top Chern class, this is $c_1(\mathbf{1}, \mathbf{t})[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}$.

It turns out that to each $\eta \in \bar{\Omega}$ there associates a pair $(\mathbf{L}_\eta, \mathbf{t}_\eta)$ of a line bundle $\mathbf{L}_\eta$ on $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and a section $\mathbf{t}_\eta \in \Gamma(\mathbf{L}_\eta)$ so that $\mathbf{1} \cong \otimes_{\eta \in \bar{\Omega}} \mathbf{L}_\eta$ as line bundles and under this isomorphism $\mathbf{t} = \prod_{\eta \in \bar{\Omega}} \mathbf{t}_\eta$. This says that $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ “virtually” is a union of normal crossing divisors, each associated to an $\eta \in \bar{\Omega}$ and is defined by the vanishing of $\mathbf{t}_\eta$. This can be seen as follows: Let $f \in \mathfrak{M}(\mathfrak{W}_0, \Gamma)$ be a general point, say represented by $f: X \rightarrow W_0$. Then $f_1 = f|_{X_1}$ with $X_1 = f^{-1}(Y_1)$ defines a relative stable morphism to Y_1^{rel} . Similarly we have the induced relative stable morphism $f_2: X_2 \rightarrow Y_2^{\text{rel}}$. Let Γ_1 and Γ_2 be the topological types of f_1 and f_2 , respectively. The fact that f can be reconstructed from the pair (f_1, f_2) provides us a triple $\eta = (\Gamma_1, \Gamma_2, I)$, which belongs to $\bar{\Omega}$. The general points $f \in \mathfrak{M}(\mathfrak{W}_0, \Gamma)$ that share identical η defines a closed subset in $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. This set is homeomorphic to $\mathbf{t}_\eta^{-1}(0)$. The miracle is that such closed subset carries a natural closed subscheme structure, and is in fact defined by the vanishing of a Cartier divisor $(\mathbf{L}_\eta, \mathbf{t}_\eta)$ and the tensor product of all such $(\mathbf{L}_\eta, \mathbf{t}_\eta)$ is the $(\mathbf{1}, \mathbf{t})$ that defines the moduli stack $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. Then by applying the known identity of the localized top Chern class, we have

$$[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}} = c_1(\mathbf{1}, \mathbf{t})[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}} = \sum_{\eta \in \bar{\Omega}} c_1(\mathbf{L}_\eta, \mathbf{t}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}.$$

To prove the degeneration formula in cycle form, we need to show that

$$c_1(\mathbf{L}_\eta, \mathbf{t}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}} = \frac{\mathfrak{m}(\eta)}{|\text{Eq}(\eta)|} \Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}).$$

This is done as follows: First we will show that the vanishing locus $\mathbf{t}_\eta^{-1}(0) \subset \mathfrak{M}(\mathfrak{W}, \Gamma)$ is homeomorphic to the image stack of Φ_η . More than that, we will show that the cycle $c_1(\mathbf{L}_\eta, \mathbf{t}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}$ is a multiple of the virtual moduli cycle of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$, endowed with the obstruction theory induced by the Cartesian product. This leads to the formula above. Here the multiplicity is $\mathfrak{m}(\eta)/|\text{Eq}(\eta)|$, where both the numerator and the denominator are as mentioned before.

Finally, we explain the notations in the degeneration formula. This formula is an immediate corollary of the degeneration formula in cycle form. The only new symbols are j_{i*} , K_η and $\beta_{\eta, i, j}$. First $j_i: Y_i \rightarrow W_0$ is the inclusion and hence $j_i^* \alpha(0)$ is the pull back cohomology. Secondly, given any $\eta = (\Gamma_1, \Gamma_2, I) \in \bar{\Omega}$, we can form the moduli space of stable curves (not necessary connected but with no vector fields) whose topology are determined by Γ_i . We denote such moduli space by $\mathfrak{M}_{\Gamma_i^0}$. For any pair $(C_1, C_2) \in \mathfrak{M}_{\Gamma_1^0} \times \mathfrak{M}_{\Gamma_2^0}$ we can glue C_1 with C_2 pairwise along all pairs of

the i -th distinguished nodes. This defines a morphism $G_\eta : \mathfrak{M}_{\Gamma_1^0} \times \mathfrak{M}_{\Gamma_2^0} \longrightarrow \mathfrak{M}_{g,k}$. The terms $\beta_{\eta,i,j}$ are terms appeared in the Kunneth decomposition,

$$G_\eta^*(\beta) = \sum_{j \in K_\eta} \beta_{\eta,1,j} \boxplus \beta_{\eta,2,j},$$

assuming it exists.

As mentioned in the introduction of [Li], the construction of the moduli stacks $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$, and the derivation of the degeneration formula in this paper will be useful in studying several problems in algebraic geometry, some related to mathematical physics. Some of these will be addressed in the future research.

The degeneration formula of Gromov-Witten invariants for symplectic manifolds has been pursued by several groups prior this work. (In symplectic setting this is when a smooth symplectic manifold degenerates to a union of two smooth symplectic manifolds intersecting transversally, called the symplectic sum.) In [Tia], Tian studied the symplectic sums for semi-positive symplectic manifolds and showed how to derive the decomposition formula of the Gromov-Witten invariants in this setting. Later, A.Li-Ruan [LR] worked out a version of the relative Gromov-Witten invariants and the degeneration formula of Gromov-Witten invariants for general symplectic manifolds and symplectic sum. The degeneration formula in numerical form proved in this paper is analogous to the degeneration formula in [LR]. A parallel theory was developed by Ionel-Parker around the same time [IP1, IP2, IP3]. Their formula works for more general cases and is largely analogous to ours. It contains a correction term, which is expected to be trivial when the symplectic sum is along a (holomorphic) divisor. The SFT theory of Eliashberg-Givental-Hofer [EGH] is a very general theory part of which can be interpreted as research along this line. The degeneration formula in cycle form proved here is new.

This paper consists of five sections. In the first section, we work out the obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$, $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ and other related moduli stacks. The main result of this section is that the standard obstruction theories of these moduli stacks are perfect. The second section is devoted to construct the virtual moduli cycles of these moduli stacks. We present a modified construction of virtual moduli cycles which allow one to construct such cycles without assuming the existence of global locally free sheaves that resolve the obstruction sheaves, as assumed in [LT2] (and also assumed in [BF] but recently removed in [Kr2]). The Gromov-Witten invariants of W_0 and the relative Gromov-Witten invariants of Z^{rel} are constructed in this section. In section three, we constructed the line bundles with sections $(\mathbf{L}_\eta, \mathbf{t}_\eta)$ mentioned in the introduction and in section four we demonstrated how to derive the degeneration formula, assuming a series of key Lemmas. The last section is devoted to the proof of these key Lemmas. In Appendix, we give a expression of the obstruction space of a closed point in $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and in $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ in terms of some known cohomology groups.

1. DEFORMATION THEORY OF LOG-MORPHISMS

In this section we will first recall the notion of morphisms between schemes with log-structures (in short log-morphism). We will then show that the notion of pre-deformable morphisms introduced in [Li] is a special class of log-morphisms. In the end, we will work out the deformation theory of log-morphisms.

1.1. Pre-deformable morphisms and log-morphisms. Let $f: \mathcal{X} \rightarrow W[n]$ be a flat family of pre-deformable morphisms over S with $g: S \rightarrow \mathbf{A}^{n+1}$ the morphism underlying f (cf. [Li, §2]). In this subsection, we will give $\mathcal{X}$ and $W[n]$ canonical log structures and show that f is a morphism between schemes with log structures. We will then use the sheaf of log-differentials to describe the deformation of pre-deformable morphisms. All materials concerning schemes with log structures are drawn from the papers of Kato [Ka1, Ka2].

We first recall the notion of logarithmic structures (log structures) on schemes, following [Ka1, Ka2]. Let X be any scheme with $\mathcal{O}_X$ its structure sheaf. We view $\mathcal{O}_X$ as a sheaf of monoids under multiplication.

Definition 1.1. A pre-log structure on X is a homomorphism $\alpha: \mathcal{M} \rightarrow \mathcal{O}_X$ of sheaves of monoids, where $\mathcal{M}$ is a sheaf of commutative monoids on the étale site $X_{\text{ét}}$ of X . The pre-log structure $\alpha: \mathcal{M} \rightarrow \mathcal{O}_X$ is said to be a log structure if α induces an isomorphism $\alpha: \alpha^{-1}(\mathcal{O}_X^\times) \rightarrow \mathcal{O}_X^\times$, where $\mathcal{O}_X^\times$ is the subsheaf of invertible elements in $\mathcal{O}_X$.

Given a pre-log structure, one can construct canonically an associated log structure $\alpha^a: \mathcal{M}^a \rightarrow \mathcal{O}_X$, where $\mathcal{M}^a = (\mathcal{M} \oplus \mathcal{O}_X^\times) / \alpha^{-1}(\mathcal{O}_X^\times)$ (cf. [Ka2, §2]). A morphism $(X, \mathcal{M}) \rightarrow (Y, \mathcal{N})$ of schemes with log structures is defined to be a pair (f, h) of a morphism of schemes $f: X \rightarrow Y$ and a homomorphism $h: f^{-1}(\mathcal{N}) \rightarrow \mathcal{M}$ that satisfy the obvious commutativity condition: The composite $f^{-1}(\mathcal{N}) \rightarrow \mathcal{M} \rightarrow \mathcal{O}_X$ is identical to $f^{-1}(\mathcal{N}) \rightarrow f^{-1}(\mathcal{O}_Y) \rightarrow \mathcal{O}_X$. For convenience, given a scheme $(X, \mathcal{M})$ with local log structure, we shall abbreviate it to $X^\dagger$ with the local log structure $\mathcal{M}$ implicitly understood. Accordingly, we will abbreviate a morphism $(X, \mathcal{M}) \rightarrow (Y, \mathcal{N})$ between schemes with local log structures by $f: X^\dagger \rightarrow Y^\dagger$.

A typical example of a scheme with log structure is the log structure of a pair (X, D) of a smooth scheme X and a divisor $D \subset X$ with normal crossing singularities (cf. [Ka1, (1.5)]). Let $\pi_n: W[n] \rightarrow \mathbf{A}^{n+1}$ be the expanded family constructed in [Li, §2] associated to a family $W \rightarrow C$ and an étale $C \rightarrow \mathbf{A}^1$. Then $W[n] \times_{\mathbf{A}^1} 0 \subset W[n]$ is a divisor with normal crossing singularities having $n+2$ irreducible components. Here $\mathbf{A}^{n+1} \rightarrow \mathbf{A}^1$ is the morphism $(z_1, \dots, z_{n+1}) \mapsto z_1 \cdots z_{n+1}$. We let $W[n]^\dagger$ and $\mathbf{A}^{n+1\dagger}$ be $W[n]$ and $\mathbf{A}^{n+1}$ endowed with the log-structures induced by the pairs

$$W[n] \times_{\mathbf{A}^1} 0 \subset W[n] \quad \text{and} \quad \mathbf{A}^{n+1} \times_{\mathbf{A}^1} 0 \subset \mathbf{A}^{n+1},$$

respectively.

Let $X^\dagger$ be a log-scheme. A chart of $X^\dagger$ consists of an étale neighborhood $U \subset X$, a constant sheaf of monoids P and a homomorphism $P \rightarrow \mathcal{O}_U$ so that the associated log structure P^a of P is isomorphic to the log structure on U . We now describe the charts of $W[n]^\dagger$. Recall that the projection π_n is smooth away from $n+1$ disjoint smooth codimension 2 subvarieties $\mathbf{D}_1, \dots, \mathbf{D}_{n+1}$, indexed so that the component $\mathbf{D}_l$ surjects onto the l -th coordinate hyperplane $H_l \subset \mathbf{A}^{n+1}$. Let $y \in \mathbf{D}_l$ be any point. A chart of $W[n]$ along y consists of a pair $(\mathcal{W}, \psi)$, where $\mathcal{W}$ is a Zariski open subset of $y \in W[n]$ and ψ is a smooth morphism

$$(1.1) \quad \psi: \mathcal{W} \longrightarrow \text{Spec } \mathbb{k}[w_1, w_2] \otimes_{\mathbb{k}[t_l]} \Gamma(\mathbf{A}^{n+1}) \triangleq \Theta_l$$

so that the canonical projections $\mathcal{W} \rightarrow \mathbf{A}^{n+1}$ is the composite of ψ with the projection $\Theta_l \rightarrow \mathbf{A}^{n+1}$. Here the two projections in the fiber product are defined via $t_l \mapsto w_1 w_2$ and by viewing t_l as the l -th standard coordinate variable of $\mathbf{A}^{n+1}$.

Let $\mathbb{N}^2 \rightarrow \mathcal{O}_{\mathcal{W}}$ be the homomorphism of monoids³ defined by $e_i \mapsto \psi^*(w_i)$ and let $\mathbb{N}^{n+1} \rightarrow \mathcal{O}_{\mathbf{A}^{n+1}}$ be defined via $e_l \mapsto t_l$. We then form the product $\mathbb{N}^2 \times_{\mathbb{N}_l} \mathbb{N}^{n+1}$ where $\mathbb{N}_l \equiv \mathbb{N}$, $\mathbb{N}_l \rightarrow \mathbb{N}^2$ is defined by $e \mapsto e_1 + e_2$ while $\mathbb{N}_l \rightarrow \mathbb{N}^{n+1}$ is the inclusion as the l -th copy in $\mathbb{N}^{n+1}$. Because of the relation $t_l \mapsto w_1 w_2$, the homomorphism

$$(1.2) \quad \mathbb{N}^2 \times_{\mathbb{N}_l} \mathbb{N}^{n+1} \longrightarrow \mathcal{O}_{\mathcal{W}}$$

induced by $\mathbb{N}^2 \rightarrow \mathcal{O}_{\mathcal{W}}$ and $\mathbb{N}^{n+1} \rightarrow \mathcal{O}_{\mathbf{A}^{n+1}} \rightarrow \mathcal{O}_{\mathcal{W}}$ defines a pre-logarithmic structure on $\mathcal{W}$. This defines a chart of $W[n]^\dagger$ near $y \in \mathbf{D}_l$. Similarly, the log-structure of $\mathbf{A}^{n+1}$ is given by the homomorphism $\mathbb{N}^{n+1} \rightarrow \mathcal{O}_{\mathbf{A}^{n+1}}$ via $e_l \mapsto t_l$.

Now let $f : \mathcal{X} \rightarrow W[n]$ be a flat family of pre-deformable morphisms over S , defined in [Li, §2]. Our next step is to show that we can endow $\mathcal{X}/S$ with a log structure so that f becomes a morphism between schemes with log structures. We will use the following convention throughout this paper. We call $p : U \rightarrow V$ an étale neighborhood of the family $\mathcal{X}/S$ if U and V are étale neighborhoods of $\mathcal{X}$ and S , respectively, such that the diagram

$$(1.3) \quad \begin{array}{ccc} U & \longrightarrow & \mathcal{X} \\ p \downarrow & & \pi \downarrow \\ V & \longrightarrow & S. \end{array}$$

is commutative.

Definition 1.2. Let $\pi : \mathcal{X} \rightarrow S$ be a flat family of nodal curves and let $x \in \mathcal{X}$ be a node of the fibers of the family $\mathcal{X}/S$. A chart of the nodes of $\mathcal{X}/S$ near x consists of an affine étale neighborhood $p : \mathcal{U} \rightarrow \mathcal{V}$ of $\mathcal{X}/S$ near x , two regular functions z_1 and $z_2 \in \Gamma(\mathcal{O}_{\mathcal{U}})$ and a regular function $s \in \Gamma(\mathcal{O}_{\mathcal{V}})$ satisfying $z_1 z_2 = p^*(s)$ such that the homomorphism

$$(1.4) \quad \phi : \mathbb{k}[z_1, z_2] \otimes_{\mathbb{k}[s]} \Gamma(\mathcal{O}_{\mathcal{V}}) \longrightarrow \Gamma(\mathcal{O}_{\mathcal{U}})$$

is an étale homomorphism, that $\{z_1 = z_2 = 0\} \subset \mathcal{U}$ is connected and that the induced homomorphism

$$\hat{\phi} : (\mathbb{k}[z_1, z_2] \otimes_{\mathbb{k}[s]} \Gamma(\mathcal{O}_{\mathcal{V}}))^\wedge \longrightarrow \Gamma(\mathcal{O}_{\mathcal{U}})^\wedge$$

is an isomorphism.

A few remarks are in order. First ϕ is defined by viewing z_1, z_2 and s on one hand as formal variables and on the other hand as regular functions. This should cause no confusion since their roles are clear from the context. Also, the homomorphism $\mathbb{k}[s] \rightarrow \mathbb{k}[z_1, z_2]$ is defined by $s \mapsto z_1 z_2$. As a convention, in this section we will use $\hat{L}$ and $\hat{h}$ (or $L^\wedge$ and $h^\wedge$) to denote the I -adic completion of the ring L and the image of $h \in L$ under the homomorphism $L \rightarrow \hat{L}$, assuming $I = (z_1, z_2)$ is an ideal of L .

Lemma 1.3. Let $\pi : \mathcal{X} \rightarrow S$ be a flat family of nodal curves and let $x \in \mathcal{X}$ be a node in $\mathcal{X}_y$, where $y = \pi(x)$. Then there exists a chart of the nodes of $\mathcal{X}/S$ near x .

Proof. The proof is straightforward and will be omitted. $\square$

We next recall the notion of morphisms of pure contact. We let $(\mathcal{U}/\mathcal{V}, \phi)$ be a chart of a node of $\mathcal{X}/S$, as in Definition 1.2. For convenience we let $A = \Gamma(\mathcal{O}_{\mathcal{V}})$ and $R = \Gamma(\mathcal{O}_{\mathcal{U}})$. As before we will view z_1 and z_2 (resp. s) as regular functions of

³We will denote by $\mathbb{N}$ the additive monoid of non-negative integers, by $\mathbb{N}^k$ the direct product monoid with standard generators $e_1, \dots, e_k$.

$\mathcal{U}$ (resp. $\mathcal{V}$), thus are elements in R (resp. A). We let $\mathcal{S}$ (resp. $\mathcal{T}$) be the set of all $a \in R$ (resp. $a \in A$) so that their residue classes $\bar{a} \in R/(z_1, z_2)$ (resp. $\bar{a} \in A/(s)$) are units. The set $\mathcal{S}$ and $\mathcal{T}$ form multiplicative systems in R and A respectively. We let $R_{\mathcal{S}}$ and $A_{\mathcal{T}}$ be the localization of the respective rings. As before, we let $\mathbb{k}[t] \rightarrow \mathbb{k}[w_1, w_2]$ be defined by $t \mapsto w_1 w_2$. We let $\hat{A}$ be the (s) -adic completion of A .

Definition 1.4. Let $\mathbb{k}[t] \rightarrow A$ be a homomorphism and $\varphi: \mathbb{k}[w_1, w_2] \rightarrow R$ be a $\mathbb{k}[t]$ -homomorphism. We say φ is formally of pure contact of order m if the induced homomorphism $\hat{\varphi}: \mathbb{k}[w_1, w_2] \rightarrow (\mathbb{k}[[z_1, z_2]] \otimes_{\mathbb{k}[[s]]} A)^{\wedge}$ is of pure contact of order m , as defined in Definition 2.3 in [Li]. Namely, there are units $\hat{h}_1, \hat{h}_2 \in (\mathbb{k}[[z_1, z_2]] \times_{\mathbb{k}[[s]]} A)^{\wedge}$ satisfying $\hat{h}_1 \hat{h}_2 \in \hat{A}$ so that $\hat{\varphi}(w_i) = \hat{h}_i z_i^m$, possibly after exchanging w_1 and w_2 . We say φ is of pure contact of order m if there are units $h_1, h_2 \in R_{\mathcal{S}}$ satisfying $h_1 h_2 \in A_{\mathcal{T}}$ such that $\varphi(w_i) = z_i^m h_i$ in $R_{\mathcal{S}}$, possibly after exchanging w_1 and w_2 .

We have the following facts whose proof can be found in the Appendix.

Lemma 1.5. The notion of pure contact is independent of the choice of the charts of the nodes of $\mathcal{U}/\mathcal{V}$.

Lemma 1.6. Let the notation be as in Definition 1.4. Then φ is of pure contact if and only if it is formally of pure contact.

Let us continue to denote by f a pre-deformable morphism $f: \mathcal{X} \rightarrow W[n]$ over S . To define log-structure on $\mathcal{X}$ we need to introduce charts of f . Let $x \in \mathcal{X}$ be any closed point so that $f(x) \notin \mathbf{D}$. Here $\mathbf{D}$ is the union of all $\mathbf{D}_l$. A chart of f near x is a triple $(\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}, \mathcal{W}_{\alpha}, f_{\alpha})$ of an étale neighborhood $\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}$ of $x \in \mathcal{X}$, a Zariski neighborhood $\mathcal{W}_{\alpha} \subset W[n] - \mathbf{D}$ and $f_{\alpha} = f|_{\mathcal{U}_{\alpha}}$ so that $f_{\alpha}(\mathcal{U}) \subset \mathcal{W}_{\alpha}$. Charts of this type will be called of the first kind. Next let $x \in \mathcal{X}$ be a distinguished node of $\mathcal{X}/S$, namely $f(x) \in \mathbf{D}_{l_{\alpha}}$ for some l_{α} . We pick a chart $(\mathcal{W}_{\alpha}, \psi_{\alpha})$ of $f(x) \in W[n]$ with ψ_{α} as in (1.1). Because f is pre-deformable, by Lemma 1.6 we can find a chart of the nodes of $\mathcal{X}/S$ near x , say given by $(\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}, \phi_{\alpha})$ as in Definition 1.2, so that $\mathcal{U}_{\alpha} \rightarrow \mathcal{X} \rightarrow W[n]$ lifts to $f_{\alpha}: \mathcal{U}_{\alpha} \rightarrow \mathcal{W}_{\alpha}$, and further there are units $h_{\alpha,1}, h_{\alpha,2} \in \Gamma(\mathcal{O}_{\mathcal{U}_{\alpha}})^{\times}$, elements $g_{\alpha,1}, \dots, g_{\alpha,n+1} \in \Gamma(\mathcal{O}_{\mathcal{V}_{\alpha}})$ and an integer m_{α} so that

$$(1.5) \quad h_{\alpha,1} h_{\alpha,2} \in \Gamma(\mathcal{O}_{\mathcal{V}_{\alpha}}), \quad f_{\alpha}^*(w_{\alpha,i}) = z_{\alpha,i}^{m_{\alpha}} h_{\alpha,i} \quad \text{and} \quad f_{\alpha}^*(t_j) = g_{\alpha,j}$$

⁴ for $i = 1, 2$ and $j = 1, \dots, n+1$. Here we comment that we will not distinguish $w_{\alpha,i}$ (resp. $z_{\alpha,i}$) with $\psi_{\alpha}^*(w_{\alpha,i})$ (resp. $\phi_{\alpha}^*(z_{\alpha,i})$). In other words, we will view $w_{\alpha,i} \in \mathcal{O}_{\mathcal{W}_{\alpha}}$ (resp. $z_{\alpha,i} \in \mathcal{O}_{\mathcal{U}_{\alpha}}$). Similarly we will view $t_l \in \mathcal{O}_{W[n]}$ via the pull back $\mathcal{O}_{\mathbf{A}^{n+1}} \rightarrow \mathcal{O}_{W[n]}$. Note that since $w_{\alpha,1} w_{\alpha,2} = t_{l_{\alpha}}$ and $z_{\alpha,1} z_{\alpha,2} = s_{\alpha}$, we must have

$$(1.6) \quad g_{\alpha, l_{\alpha}} = s_{\alpha}^{m_{\alpha}} (h_{\alpha,1} h_{\alpha,2}).$$

We will call such triplet $(\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}, \mathcal{W}_{\alpha}, f_{\alpha})$ with $(\phi_{\alpha}, \psi_{\alpha})$ understood a chart of f of the second kind.

Simplification 1.7. In case $(\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}, \mathcal{W}_{\alpha}, f_{\alpha})$ is a chart of the second kind, for simplicity we assume $h_{\alpha,1} \equiv h_{\alpha,2} \equiv 1$. This is possible possibly after an étale base change of $\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}$.

⁴By this we mean it is in the image of the pull back homomorphism $\Gamma(\mathcal{O}_{\mathcal{V}_{\alpha}}) \rightarrow \Gamma(\mathcal{O}_{\mathcal{U}_{\alpha}})$.

We now cover $\mathcal{X}$ by charts of f , say $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, \mathcal{W}_\alpha, f_\alpha)$ indexed by $\alpha \in \Lambda$, of the first or the second kinds satisfying the Simplification 1.7. Let α be a chart of the second kind with ϕ_α and ψ_α understood. We then let $M_\alpha^0 = \mathbb{N}^2$ (resp. $N_\alpha^0 = \mathbb{N}$) and let $M_\alpha^0 \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ (resp. $N_\alpha^0 \rightarrow \mathcal{O}_{\mathcal{V}_\alpha}$) be the pre-log structure defined by $e_{\alpha,i} \mapsto z_{\alpha,i}$ (resp. $e_\alpha \mapsto s_\alpha$). Note that $N_\alpha^0 \rightarrow M_\alpha^0$ defined by $e_\alpha \mapsto e_{\alpha,1} + e_{\alpha,2}$ makes the projection $\mathcal{U}_\alpha \rightarrow \mathcal{V}_\alpha$ a morphism between schemes with (their respective associated) log structures. We now define the desired log structures on S . Let $\xi \in S$ be any closed point and let $\mathcal{X}_\xi$ be the fiber of $\mathcal{X}$ over ξ with $\Sigma = f^{-1}(\mathbf{D}) \cap \mathcal{X}_\xi$ be the set of distinguished nodes of $\mathcal{X}_\xi$. We then let Λ_0 be a collection of $\alpha \in \Lambda$ so that $\{\mathcal{U}_\alpha\}_{\alpha \in \Lambda_0}$ covers a neighborhood of $\mathcal{X}_\xi \subset \mathcal{X}$. We let K_l be those α so that $f^{-1}(\mathbf{D}_l) \cap \mathcal{U}_\alpha \neq \emptyset$. We let K be the union of $K_1, \dots, K_{n+1}$. By eliminating redundant α from Λ_0 we can assume that each node of $\mathcal{X}_\xi \cap f^{-1}(\mathbf{D}_l)$ is covered by at most one $\alpha \in K_l$. We then pick an étale neighborhood $\mathcal{V}$ of $\xi \in S$ so that $\mathcal{V} \rightarrow S$ factor through $\mathcal{V}_\alpha \rightarrow S$ for all $\alpha \in \Lambda_0$. For each $l \in [n+1]$ we let $\bar{N}_l = \oplus_{\alpha \in K_l} N_\alpha^0$ and let N_l be the quotient (monoid) of $\bar{N}_l$ by the relations $m_\alpha e_\alpha = m_\beta e_\beta$ for all pairs $\alpha, \beta \in K_l$. In case $K_l = \emptyset$, we agree $N_l = \bar{N}_l = \mathbb{N}$. Note that for $\alpha \in K_l$ the homomorphism $\bar{N}_l \rightarrow \mathcal{O}_{\mathcal{V}}$ defined by $e_\alpha \mapsto s_\alpha$ and the pull back $\mathcal{O}_{\mathcal{V}_\alpha} \rightarrow \mathcal{O}_{\mathcal{V}}$ descends to a homomorphism $N_l \rightarrow \mathcal{O}_{\mathcal{V}}$, because of the relations (1.5) and (1.6). When $K_l = \emptyset$, we define $N_l \rightarrow \mathcal{O}_{\mathcal{V}}$ via $e \mapsto g^*(t_l)$. We define the pre-log structures $N_{\mathcal{V}}$ and $\bar{N}_{\mathcal{V}}$ on $\mathcal{V}$ to be the ones given by the direct sums

$$(1.7) \quad N_{\mathcal{V}} \triangleq \oplus_{l=1}^{n+1} N_l \longrightarrow \mathcal{O}_{\mathcal{V}} \quad \text{and} \quad \bar{N}_{\mathcal{V}} \triangleq \oplus_{l=1}^{n+1} \bar{N}_l \longrightarrow \mathcal{O}_{\mathcal{V}}.$$

We denote by $\mathcal{N}_{\mathcal{V}}$ and $\bar{\mathcal{N}}_{\mathcal{V}}$ the associated log structures on $\mathcal{V}$.

We next define the desired pre-log structure on $\mathcal{U}_\alpha$. By replacing $\mathcal{U}_\alpha$ by $\mathcal{U}_\alpha \times_{\mathcal{V}_\alpha} \mathcal{V}$, we can assume $\mathcal{V}_\alpha = \mathcal{V}$. Accordingly we let N_α and $\bar{N}_\alpha \rightarrow \mathcal{O}_{\mathcal{V}}$ be that induced by $N_{\mathcal{V}}$ and $\bar{N}_{\mathcal{V}}$. Now let l be so that $\alpha \in K_l$. Recall that $\mathcal{U}_\alpha$ has a pre-log structure given by $M_\alpha^0 \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$. Let $N_\alpha^0 \rightarrow M_\alpha^0$ be as before (defined by $e_\alpha \mapsto e_{\alpha,1} + e_{\alpha,2}$). Then we have the obvious homomorphism

$$(1.8) \quad \bar{M}_\alpha = M_\alpha^0 \times_{N_\alpha^0} \bar{N}_l \oplus (\oplus_{l' \neq l} \bar{N}_{l'}) \longrightarrow \mathcal{O}_{\mathcal{U}_\alpha}$$

and

$$(1.9) \quad M_\alpha = M_\alpha^0 \times_{N_\alpha^0} N_l \oplus (\oplus_{l' \neq l} N_{l'}) \longrightarrow \mathcal{O}_{\mathcal{U}_\alpha}.$$

Here $M_\alpha^0 \times_{N_\alpha^0} \bar{N}_l \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ is induced by $M_\alpha^0 \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ and $\bar{N}_l \rightarrow \mathcal{O}_{\mathcal{V}_\alpha} \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ while $\bar{N}_{l'} \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ is the composite of $N_{l'} \rightarrow \mathcal{O}_{\mathcal{V}_\alpha} \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$. The arrow in (1.9) is defined similarly. They define two pre-log structures on $\mathcal{U}_\alpha$. We let $\bar{\mathcal{M}}_\alpha \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ and $\mathcal{M}_\alpha \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$ be the associated log-structures. Note that the obvious $N_\alpha \rightarrow M_\alpha$ and $\bar{M}_\alpha \rightarrow \bar{N}_\alpha$ make the projection $\mathcal{U}_\alpha \rightarrow \mathcal{V}_\alpha$ a morphism between schemes with respective log-structures.

Proposition 1.8. *The log structures $(\mathcal{U}_\alpha, \mathcal{M}_\alpha)$ and $(\mathcal{V}_\alpha, \mathcal{N}_\alpha)$ patch together to form log structures $\mathcal{M}$ on $\mathcal{X}$ and $\mathcal{N}$ on S . The same conclusion holds for $(\mathcal{U}_\alpha, \bar{\mathcal{M}}_\alpha)$ and $(\mathcal{V}_\alpha, \bar{\mathcal{N}}_\alpha)$. The collection of homomorphisms $N_\alpha \rightarrow M_\alpha$ makes $\mathcal{X}^\dagger \equiv (\mathcal{X}, \mathcal{M})$ a log scheme over $S^\dagger \equiv (S, \mathcal{N})$. Further, the morphism f is naturally a morphism between schemes with log structures $\mathcal{X}^\dagger/S^\dagger \rightarrow W[n]^\dagger/\mathbf{A}^{n+1\dagger}$.*

Proof. The fact that the so defined log structures on $\mathcal{U}_\alpha$ and $\mathcal{V}_\alpha$ patch together to form log structures on $\mathcal{X}$ and S is obvious. We now study the morphism f . We first investigate the morphism $g: S \rightarrow \mathbf{A}^{n+1}$ underlying f . Let $\xi \in S$ be any closed point with $(\mathcal{V}, N_{\mathcal{V}})$ a chart of $S^\dagger$, constructed before. Recall that the log

structure on $\mathbf{A}^{n+1\dagger}$ is given by the pre-log structure $\mathbb{N}^{n+1} \rightarrow \Gamma(\mathbf{A}^{n+1})$ via $e_l \mapsto t_l$. To show $g|_{\mathcal{V}} : \mathcal{V} \rightarrow \mathbf{A}^{n+1}$ is a morphism between schemes with log structures we need to define a homomorphism $\mathbb{N}^{n+1} \rightarrow N_{\mathcal{V}}$ that satisfies the obvious compatibility condition. By definition $N_{\mathcal{V}} = \oplus_l N_l$ and N_l is $\mathbb{N}$ in case $K_l = \emptyset$ and is the quotient of $\tilde{N}_l = \oplus_{\alpha \in K_l} N_{\alpha}^0$ otherwise. In the first case we define $\mathbb{N} \rightarrow N_l$ to be the unique isomorphism. In the later case we let $\mathbb{N} \rightarrow N_l$ be induced by $1 \mapsto m_{\alpha} e_{\alpha} \in N_l$ for some $\alpha \in K_l$. By the definition of N_l such definition is independent of $\alpha \in K_l$. The desired homomorphism $\mathbb{N}^{n+1} \rightarrow N_{\mathcal{V}}$ is the direct sum of these $n+1$ copies $\mathbb{N} \rightarrow N_l$. It is direct to check that this defines $\mathcal{V} \rightarrow \mathbf{A}^{n+1}$ a morphism between schemes with log structures. By working over a covering of S , this concludes that g is a morphism between $S^{\dagger} \rightarrow \mathbf{A}^{n+1\dagger}$.

The definition making f a morphism between $\mathcal{X}^{\dagger}$ and $W[n]^{\dagger}$ and compatible to $g : S^{\dagger} \rightarrow \mathbf{A}^{n+1\dagger}$ is similar, relying on the relations (1.5) and (1.6) and the assumption that all $h_{\alpha,i} \equiv 1$. This completes the proof of the Proposition. $\square$

We conclude this subsection by commenting the equivalence of the deformations of pre-deformable morphisms and the deformations of log-morphisms.

Definition 1.9. 1. A log-extension of $(\mathcal{V}_{\alpha}, \mathcal{N}_{\alpha})$ by I consists of an extension $\tilde{\mathcal{V}}_{\alpha}$ of $\mathcal{V}_{\alpha}$ by I as schemes, and an extension $N_{\alpha} \rightarrow \mathcal{O}_{\tilde{\mathcal{V}}_{\alpha}}$ of $N_{\alpha} \rightarrow \mathcal{O}_{\mathcal{V}_{\alpha}}$. We denote such extension by $(\tilde{\mathcal{V}}_{\alpha}, \tilde{\mathcal{N}}_{\alpha})$.

2. A flat log-extension of $(\mathcal{U}_{\alpha}/\mathcal{V}_{\alpha}, \mathcal{M}_{\alpha}/\mathcal{N}_{\alpha})$ by I consists of an extension $(\tilde{\mathcal{V}}_{\alpha}, \tilde{\mathcal{N}}_{\alpha})$ of $(\mathcal{V}_{\alpha}, \mathcal{N}_{\alpha})$ by I , a flat extension $\tilde{\mathcal{U}}_{\alpha} \rightarrow \tilde{\mathcal{V}}_{\alpha}$ of $\mathcal{U}_{\alpha} \rightarrow \mathcal{V}_{\alpha}$ and an extension $M_{\alpha} \rightarrow \mathcal{O}_{\tilde{\mathcal{U}}_{\alpha}}$ of the pre-log structure $M_{\alpha} \rightarrow \mathcal{O}_{\mathcal{U}_{\alpha}}$ of which the following holds: a. The projection $\tilde{\mathcal{U}}_{\alpha} \rightarrow \tilde{\mathcal{V}}_{\alpha}$ is a log-morphism under the given $N_{\alpha} \rightarrow M_{\alpha}$; b. Away from the distinguished nodes of $\mathcal{U}_{\alpha}$ the pre-log structure $M_{\alpha} \rightarrow \mathcal{O}_{\tilde{\mathcal{U}}_{\alpha}}$ is the pull back of $N_{\alpha} \rightarrow \mathcal{O}_{\tilde{\mathcal{V}}_{\alpha}}$; c. Near the distinguished nodes in $\mathcal{U}_{\alpha}$ the projection $\tilde{\mathcal{U}}_{\alpha} \rightarrow \tilde{\mathcal{V}}_{\alpha}$ is log-smooth.

We have the following Lemma which says that extending f as pre-deformable morphism is equivalent to extending f as log-morphisms.

Lemma 1.10. Let $f : \mathcal{X}/S \rightarrow W[n]/\mathbf{A}^{n+1}$ be as before with the canonical log structures understood. Let $\tilde{S} \supset S$ be an extension of S . Suppose $\tilde{S}^{\dagger}$ is a log-extension of S , $\tilde{\mathcal{X}}^{\dagger} \rightarrow \tilde{S}^{\dagger}$ is a flat log-extension of $\mathcal{X}/S$ and $\tilde{f} : \tilde{\mathcal{X}}^{\dagger}/\tilde{S}^{\dagger} \rightarrow W[n]^{\dagger}/\mathbf{A}^{n+1\dagger}$ is an extension of f as log-morphism. Then $\tilde{f}$ is a pre-deformable extension of f and the log structures on $\tilde{\mathcal{X}}/\tilde{S}$ induced by $\tilde{f}$ coincide with the log structure of $\tilde{\mathcal{X}}^{\dagger}/\tilde{S}^{\dagger}$.

Proof. The proof is straightforward. Since we do not need this Lemma in this paper, we shall omit the proof here. $\square$

1.2. Deformation of pre-deformable morphisms. The goal of this subsection is to work out the deformation theory of pre-deformable morphisms. Based on the equivalence Lemma, it is natural to work out the deformation theory of pre-deformable morphisms in the frame work of log-morphisms. However, the deformation theory of log-morphisms worked out in [Ka1, Ka2] deal with the question on how to extend families over $S^{\dagger}$ to $\tilde{S}^{\dagger}$. In our situation, the log structure on S for a pre-deformable family $f : \mathcal{X}/S \rightarrow W[n]/\mathbf{A}^{n+1}$ relies on the morphism f . Hence for any extension of base $S \subset \tilde{S}$ the extension of log structure to $\tilde{S}$ is part of the extension problem. Hence the deformation theory of Kato [Ka2] can not be applied

directly. In this subsection, we will work with the deformation of pre-deformable directly.

We begin with recalling some basic notion in deformation theory. Our treatment follows [Art, LT2]. Let A be an $\bar{A}$ -algebra. We let $\mathfrak{Tri}_{A/\bar{A}}$ be the category whose objects consists of are all triples (B, I, φ) where B are $\bar{A}$ -algebras, I are ideals of B such that $I^2 = 0$ and φ are $\bar{A}$ -homomorphisms $A \rightarrow B/I$. Let $\xi = (B, I, \varphi)$ and $\xi' = (B', I', \varphi')$ be two objects in $\mathfrak{Tri}_{A/\bar{A}}$. A morphism from ξ to ξ' consists of an $\bar{A}$ -homomorphism $r: B \rightarrow B'$ so that $r(I) \subset I'$ and $\varphi' = \varphi \circ r_0$ where $r_0: B/I \rightarrow B'/I'$ is the induced homomorphism. We let $\mathfrak{Mod}_A$ be the category whose objects are pairs (B, I) where B are A -algebras and I are B -modules. Morphisms from (B, I) to (B', I') are pairs $(r, \tilde{r})$ where $r: B \rightarrow B'$ are A -homomorphisms and $\tilde{r}$ are B -homomorphisms $I \rightarrow I'$. We let $\mathfrak{Mod}_A^*$ be the category whose objects are (v, B, I) where $(B, I) \in \text{Ob}(\mathfrak{Mod}_A)$ and $v \in I$. Morphisms from (B, I, v) to (B', I', v') are pairs $(r, \tilde{r})$ as in $\mathfrak{Mod}_A$ so that $\tilde{r}(v) = v'$.

We define $\mathfrak{Def}_{A/\bar{A}}: \mathfrak{Tri}_{A/\bar{A}} \rightarrow (\text{Sets})$ be the functor that associates to any $\xi = (B, I, \varphi)$ the set of all $\bar{A}$ -homomorphisms $A \rightarrow B$ extending $\varphi: A \rightarrow B/I$. (In case $\bar{A}$ is understood, we will omit the subscript $\bar{A}$.) It is known that under some mild conditions [Shl] this set admits B/I -module structures. Once this is the case, then after fixing a reference element $a \in \mathfrak{Def}_A(\xi)$ we can give $\mathfrak{Def}_A(\xi)$ a natural B/I -module structure. In particular, if $B = B/I * I$, which is the trivial extension of B/I by the module I , then $\mathfrak{Def}_A(\xi)$ contains the trivial extension $B \rightarrow A$ induced by $B = B/I * I \rightarrow B/I \rightarrow A$. Using this as the reference element, we obtain a canonical module structure on $\mathfrak{Def}_A(\xi)$. Note that there is a natural functor $\mathfrak{Mod}_A \rightarrow \mathfrak{Tri}_{A/\bar{A}}$ that sends (B, I) to $(B * I, I, \varphi)$ where $\varphi: A \rightarrow B$ is the obvious homomorphism. We let $\mathfrak{Def}_A^1: \mathfrak{Mod}_A \rightarrow \mathfrak{Mod}_A$ be the composite of $\mathfrak{Def}_A$ with $\mathfrak{Mod}_A \rightarrow \mathfrak{Tri}_{A/\bar{A}}$. We call $\mathfrak{Def}_A^1$ the functor of the first order deformations.

Now let $E^\bullet$ be any complex of A -modules. Then for any integer i the i -th cohomology of $E^\bullet$ defines a functor $\mathfrak{h}^i(E^\bullet): \mathfrak{Mod}_A \rightarrow \mathfrak{Mod}_A$ via $(B, I) \mapsto (B, h^i(E^\bullet \otimes_A I))$.

Definition 1.11. *Let $S = \text{Spec } A$ be an affine scheme over $T = \text{Spec } \bar{A}$. A perfect obstruction theory of S/T consists of a two term complex of finitely generated free A -modules $E^\bullet = [E^1 \rightarrow E^2]$ (indexed at $[1, 2]$) and an obstruction assignment ob taking value in the second cohomology of $E^\bullet$ of which the following hold:*

1. *The functor $\mathfrak{Def}_A^1$ is isomorphic to the functor $\mathfrak{h}^1(E^\bullet)$.*
2. *For any triple $(B, I, \varphi) \in \text{Ob}(\mathfrak{Tri}_{A/\bar{A}})$, the element*

$$\text{ob}(B, I, \varphi) \in \mathfrak{h}^2(E^\bullet)(I) = h^2(E^\bullet \otimes_A I)$$

is the obstruction class to extending $\varphi: A \rightarrow B/I$ to an $\bar{A}$ -homomorphism $A \rightarrow B$.

3. *The obstruction assignment*

$$(B, I, \varphi) \mapsto (B, h^2(E^\bullet \otimes_A I), \text{ob}(B, I, \varphi))$$

is a functor from $\mathfrak{Tri}_{A/\bar{A}}$ to $\mathfrak{Mod}_A^$. Namely it satisfies the base change property.*

A few remarks are in order here. First, in [LT2], we only considered the case where $T = \text{Spec } \mathbb{k}$. Here we need to study the relative case for the proof of the degeneration formula. When $T = \text{Spec } \mathbb{k}$, we will omit T from the notation. When T is non-trivial, we will call the obstruction theory so defined the relative obstruction theory. Secondly, when we restrict to the subcategory of all triples (B, I, φ) so that B are Artin local rings with residue fields $\mathbb{k}$, then the above data is the obstruction

theory to deforming $\varphi(\mathrm{Spec} \mathbb{k})$ in S . Thirdly, since $\mathrm{Hom}_A(\Omega_{A/\bar{A}} \otimes_A B/I, I)$ is canonically isomorphic to $\mathfrak{Def}_{A/\bar{A}}^1(I)$, we have $\ker\{E^{2\vee} \rightarrow E^{1\vee}\} \cong \Omega_{A/\bar{A}}$. Lastly, in the definition we can replace $E^\bullet$ by its associated complex of sheaves of $\mathcal{O}_{\mathrm{Spec} B}$ -modules and modify the wording accordingly. This is convenient if we work with obstruction of schemes or stacks. We will call $h^2(E^\bullet)$ the obstruction module and call its associated sheaf (of $\mathcal{O}_{\mathrm{Spec} B}$ -modules) obstruction sheaf.

We now investigate the deformation theory of pre-deformable morphisms. Let Γ be the data (g, k, b) representing the genus, the number of marked points and the degree of the maps. We let $\mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$ be the moduli space of stable morphisms to $W[n]$ of prescribed topological type that are also pre-deformable as morphisms to the family $W[n]/C[n]$ and are stable as morphisms to the stack $\mathfrak{W}$. As argued in section 2 in [Li], it is a Deligne-Mumford stack and it comes with a tautological morphism

$$\mathfrak{M}(W[n], \Gamma)^{\mathrm{st}} \longrightarrow \mathfrak{M}(\mathfrak{W}, \Gamma).$$

In the remainder of this section, we will cover $\mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$ by affine étale charts and construct canonical obstruction theory of each of these charts. The obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ will be the descent of the obstruction theory of $\mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$. Since this study is local, during our study we are free to shrink an open chart $S \rightarrow \mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$ if necessary.

In the remainder of this subsection we fix an affine étale chart $S \rightarrow \mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$ with $f: \mathcal{X} \rightarrow W[n]$ its universal family. We let $S = \mathrm{Spec} A$ and let $\mathcal{D} \subset \mathcal{X}$ be the divisor of the union of all marked sections of $\mathcal{X}/S$. We fix a collection of charts $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha)$ of f that covers $\mathcal{X}$ satisfying the simplification assumption. In case α is a chart of the second kind, we will reserve the symbols $\mathcal{W}_\alpha, z_{\alpha,i}, w_{\alpha,i}, \phi_\alpha, \psi_\alpha, m_\alpha$ and l_α for the data associated to the chart α . Our first task is to show that locally there are no obstruction to extending pre-deformable morphisms f_α .

We begin with the notion of flat extensions of an étale neighborhood $\mathcal{U}_\alpha/\mathcal{V}_\alpha$. Let I be an A -module. We say $\tilde{\mathcal{V}}_\alpha$ is an extension of $\mathcal{V}_\alpha$ by I if $\mathcal{V}_\alpha \subset \tilde{\mathcal{V}}_\alpha$ is a subscheme with the ideal sheaf $\tilde{\mathcal{I}}$ of $\mathcal{V}_\alpha \subset \tilde{\mathcal{V}}_\alpha$ satisfying $(\tilde{\mathcal{I}})^2 = 0$ and the module $\Gamma(\tilde{\mathcal{I}})$ is isomorphic to I as $\Gamma(\mathcal{O}_{\mathcal{V}})$ -modules. We say $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ is a small extension of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ by I if $\tilde{\mathcal{V}}_\alpha$ is an extension of $\mathcal{V}_\alpha$ by I and $\tilde{\mathcal{U}}_\alpha$ is a flat extension of $\mathcal{U}_\alpha$ over $\tilde{\mathcal{V}}_\alpha$. Now let $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ be an extension of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ by I . We first consider the case where α is a chart of the second kind. By Lemma 1.6, we can assume that the ϕ_α of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ in (1.4) is extended to $\tilde{\phi}_\alpha$ of $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$. We say an extension $\tilde{f}_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$ of $f_\alpha: \mathcal{U}_\alpha \rightarrow W[n]$ is a pre-deformable extension if $\tilde{\mathcal{U}}_\alpha \rightarrow W[n] \rightarrow \mathbf{A}^{n+1}$ factor through $\tilde{\mathcal{V}}_\alpha \rightarrow \mathbf{A}^{n+1}$ and if the family $\tilde{f}_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$ is pre-deformable along $\mathbf{D}_{l_\alpha}$.

We now define the space $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$ that will parameterize all such extensions. The group $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$ is the set of triples

$$(1.10) \quad (\varphi, \eta_1, \eta_2) \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, \mathcal{I}_\alpha) \oplus \mathcal{I}_\alpha^{\oplus 2}, \quad \mathcal{I}_\alpha = I \otimes_A \mathcal{O}_{\mathcal{U}_\alpha}$$

that obey the following condition:

$$(1.11) \quad \varphi(f^*dw_{\alpha,i}) = f^*(w_{\alpha,i}) \cdot \eta_i, \quad \varphi(f^*dt_l) \in I_\alpha \quad \text{and} \quad \eta_1 + \eta_2 \in I_\alpha$$

for $i = 1, 2$ and $l = 1, \dots, n+1$. Here $I_\alpha = I \otimes_A \mathcal{O}_{\mathcal{V}_\alpha}$. Since $f_\alpha(\mathcal{U}_\alpha) \subset \mathcal{W}_\alpha$ and $w_{\alpha,i} \in \Gamma(\mathcal{O}_{\mathcal{W}_\alpha})$, $f^*dw_{\alpha,i} \in f^*\Omega_{W[n]} \otimes_{\mathcal{O}_{\mathcal{X}}} \mathcal{O}_{\mathcal{U}_\alpha}$ and hence $\varphi(f^*dw_{\alpha,i})$ makes sense. Note also that because $t_{l_\alpha} = w_{\alpha,1}w_{\alpha,2}$, from (1.11) we have

$$(1.12) \quad s_\alpha^{m_\alpha}(\eta_1 + \eta_2) = \varphi(f^*dt_{l_\alpha}).$$

Clearly, $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$ is an A -module and is A -flat if I is A -flat.

When α is chart of the first kind, we define $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$ be the subgroup of $\varphi \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, \mathcal{I})$ such that $\varphi(dt_l) \in I_\alpha$ for all l . Note that in case α and β are two charts of f and $p_\alpha: \mathcal{U}_{\alpha\beta} \triangleq \mathcal{U}_\alpha \times_{\mathcal{X}} \mathcal{U}_\beta \rightarrow \mathcal{U}_\alpha$ is the projection. Then there is a canonical restriction A -homomorphism

$$p_\alpha^*: \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger \longrightarrow \mathrm{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{W[n]}, I)^\dagger.$$

We now state and prove the following local deformation Lemma.

Lemma 1.12. *Let $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ be an extension of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ by I . Then f_α automatically extends to a pre-deformable morphism $\tilde{f}_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$. Further, after fixing one such extension, say $\tilde{f}'_\alpha$, the space of all such extensions is canonically isomorphic to the space $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$.*

Proof. We will consider the case where α is a chart of the second kind. The other case is simpler. First, we extend ϕ_α to a parameterization $\tilde{\phi}_\alpha$ of $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ with $\tilde{z}_{\alpha,i}$ and $\tilde{s}_\alpha$ the corresponding extensions of $z_{\alpha,i}$ and s_α , respectively. We consider the composite $\psi_\alpha \circ f_\alpha: \mathcal{U}_\alpha \rightarrow \Theta_{l_\alpha}$, where Θ_{l_α} is defined in (1.1). Let $h_{\alpha,i} \equiv 1 \in \Gamma(\mathcal{O}_{\mathcal{U}_\alpha})$ and $g_{\alpha,l} \in \Gamma(\mathcal{O}_{\mathcal{V}_\alpha})$ be part of the definition of the chart of f_α , as in (1.5) and (1.6). Since both $\mathcal{U}_\alpha$ and $\mathcal{V}_\alpha$ are affine, we can extend $h_{\alpha,i}$ and $g_{\alpha,l}$ to $\tilde{h}_\alpha \in \Gamma(\mathcal{O}_{\tilde{\mathcal{U}}_\alpha})$ and $\tilde{g}_{\alpha,l} \in \Gamma(\mathcal{O}_{\tilde{\mathcal{V}}_\alpha})$, respectively, so that $\tilde{h}_{\alpha,1}\tilde{h}_{\alpha,2} \in \Gamma(\mathcal{O}_{\tilde{\mathcal{V}}_\alpha})$ and $\tilde{g}_{\alpha,l_\alpha} = \tilde{s}_\alpha^{m_\alpha}(\tilde{h}_{\alpha,1}\tilde{h}_{\alpha,2})$. We then define $F: \tilde{\mathcal{U}}_\alpha \rightarrow \Theta_{l_\alpha}$ by $w_{\alpha,i} \mapsto \tilde{z}_{\alpha,i}^{m_\alpha}\tilde{h}_{\alpha,i}$ for $i = 1, 2$ and $t_l \mapsto \tilde{g}_{\alpha,l}$ for $l = 1, \dots, n+1$. Since $\mathcal{W}_\alpha \rightarrow \Theta_{l_\alpha}$ is smooth, we can lift F to an extension $\tilde{f}_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$ of f_α . Namely, $\tilde{f}_\alpha$ is an extension of f_α such that $\psi_\alpha \circ \tilde{f}_\alpha = F$. The morphism $\tilde{f}_\alpha$ is a desired extension.

Now let $\tilde{f}'_\alpha$ be a fixed pre-deformable extension of f_α to $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$. Let $\tilde{f}_\alpha$ be any pre-deformable extension of f_α . As morphisms from $\tilde{\mathcal{U}}_\alpha$ to Θ_{l_α} , $\psi_\alpha \circ \tilde{f}_\alpha$ is defined by

$$(1.13) \quad (\psi_\alpha \circ \tilde{f}_\alpha)^*(w_{\alpha,i}) = \tilde{z}_{\alpha,i}^{m_\alpha}\tilde{h}_{\alpha,i} \quad \text{and} \quad (\psi_\alpha \circ \tilde{f}_\alpha)^*(t_l) = \tilde{g}_{\alpha,l}$$

and $\psi_\alpha \circ \tilde{f}'_\alpha$ is defined by

$$(1.14) \quad (\psi_\alpha \circ \tilde{f}'_\alpha)^*(w_{\alpha,i}) = \tilde{z}_{\alpha,i}^{m_\alpha}\tilde{h}'_{\alpha,i} \quad \text{and} \quad (\psi_\alpha \circ \tilde{f}'_\alpha)^*(t_l) = \tilde{g}'_{\alpha,l}.$$

We let $\eta_i = \tilde{h}_{\alpha,i} - \tilde{h}'_{\alpha,i} \in \mathcal{I}_\alpha$. Then $\eta_1 + \eta_2 \in I_\alpha$ since $\tilde{h}_{\alpha,1}\tilde{h}_{\alpha,2}$ and $\tilde{h}'_{\alpha,1}\tilde{h}'_{\alpha,2} \in \Gamma(\mathcal{O}_{\tilde{\mathcal{V}}_\alpha})$. Now let $\varphi \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, \mathcal{I})$ be defined by the difference⁵ $\mathbf{d}(\tilde{f}_\alpha - \tilde{f}'_\alpha)$. It can be easily checked, based on (1.13) and (1.14), that $(\varphi, \eta_1, \eta_2)$ is in $\mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$.

It remains to check that this correspondence is one-one and onto, which is straightforward. This completes the proof of the Lemma. $\square$

Remark 1.13. *In the remainder part of this paper, we will call $(\varphi, \eta_1, \eta_2)$ the log-difference of $\tilde{f}_\alpha$ and $\tilde{f}'_\alpha$, denoted by $\mathbf{d}^\dagger(\tilde{f}_\alpha - \tilde{f}'_\alpha)$. Furthermore, if $\tilde{f}_1$, $\tilde{f}_2$ and $\tilde{f}_3$*

⁵Let $\iota: A' \rightarrow A$ be a small ring extension with $I = \ker\{\iota\}$. We call $A' \rightarrow A$ a small extension if $I^2 = 0$. Let B be any ring. Let $f_1, f_2: B \rightarrow A$ be two ring homomorphisms so that $\iota \circ f_1 = \iota \circ f_2$. We define the difference of f_1 and f_2 to be the map $\mathbf{d}(f_1 - f_2): \Omega_B \rightarrow I$ defined by $b \otimes 1 - 1 \otimes b \mapsto f_1(b) - f_2(b) \in I$. It is an A -homomorphism $\Omega_B \otimes_B A \rightarrow I$. Note that once f_1 is fixed, then f_2 is uniquely determined by $\mathbf{d}(f_1 - f_2)$. Conversely, any $\varphi \in \mathrm{Hom}_A(\Omega_B \otimes_B A, I)$ defines a unique homomorphism $f_2: B \rightarrow A$ so that $\mathbf{d}(f_1 - f_2) = \varphi$.

are three pre-deformable extensions of f_α to $\tilde{\mathcal{U}}_\alpha$, then $\mathbf{d}^\dagger(\tilde{f}_3 - \tilde{f}_1) = \mathbf{d}^\dagger(\tilde{f}_3 - \tilde{f}_2) + \mathbf{d}^\dagger(\tilde{f}_2 - \tilde{f}_1)$.

Before we proceed, we introduce more convention which we will follow throughout this paper. We let $\iota_\alpha: \mathcal{U}_\alpha \rightarrow \mathcal{X}$ and $j_\alpha: \mathcal{V}_\alpha \rightarrow S$ be the tautological projections. As a convention, for each $\alpha \in \Lambda$ we let $A_\alpha = \Gamma(\mathcal{V}_\alpha)$ with $A \rightarrow A_\alpha$ the tautological homomorphism, for α and $\beta \in \Lambda$ we will denote by $\mathcal{U}_{\alpha\beta}$ the product $\mathcal{U}_\alpha \times_{\mathcal{X}} \mathcal{U}_\beta$ and by p_α and p_β the projections $\mathcal{U}_{\alpha\beta} \rightarrow \mathcal{U}_\alpha$ and to $\mathcal{U}_\beta$ respectively. The same convention applies to multi-indices in the obvious way. We let $\pi_\alpha: \mathcal{U}_\alpha \rightarrow \mathcal{V}_\alpha$ be the projection. For sheaf of $\mathcal{O}_{\mathcal{X}}$ -modules $\mathcal{A}$ (resp. A -module I ; divisor $\mathcal{D}$) we will denote by $\mathcal{A}_\alpha$ (resp. I_α , resp. $\mathcal{D}_\alpha$) the pull back sheaf $\iota_\alpha^* \mathcal{A}$ (resp. $I \otimes_A A_\alpha$; resp. $\mathcal{D} \times_{\mathcal{X}} \mathcal{U}_\alpha$). Also, for A -module I , we will use $\mathcal{I}$ (resp. $\mathcal{I}_\alpha$) to denote the sheaf of $\mathcal{O}_{\mathcal{X}}$ -modules $\mathcal{O}_{\mathcal{X}} \otimes_A I$ (resp. $\mathcal{O}_{\mathcal{U}_\alpha} \times_A I$). We denote by $d_\alpha: \mathcal{F}_\alpha^0 \rightarrow \mathcal{F}_\alpha^2$ the pull back of $d: F^0 \rightarrow F^1$ in (1.15). For sheaves of $\mathcal{O}_{\mathcal{X}}$ -modules $\mathcal{A}$ and $\mathcal{B}$ we agree that $\mathrm{Hom}_{\mathcal{U}_\alpha}(\mathcal{A}, \mathcal{B}) = \mathrm{Hom}_{\mathcal{U}_\alpha}(\mathcal{A}_\alpha, \mathcal{B}_\alpha)$ and for A -modules I and J we agree $\mathrm{Hom}_{A_\alpha}(I, J) = \mathrm{Hom}_{A_\alpha}(I_\alpha, J_\alpha)$.

We now study the deformation of $f: \mathcal{X}/S \rightarrow W[n]/\mathbf{A}^{n+1}$. By the deformation theory of nodal curves with sections, there is a complex of free A -modules

$$(1.15) \quad F^\bullet = [F^0 \xrightarrow{d} F^1]$$

so that for any A -module I ,

$$(1.16) \quad \mathrm{Ext}_{\mathcal{X}}^i(\Omega_{\mathcal{X}/S}(\mathcal{D}), \mathcal{I}) = h^i(F^\bullet \otimes_A I).$$

Now let $\tilde{S} \triangleq \mathrm{Spec} A * F^{1\vee}$ ⁶ with $S \subset \tilde{S}$ be the immersion induced by the obvious projection $A * F^{1\vee} \rightarrow A$. Let $\mathbf{1} \in F^1 \otimes_A F^{1\vee}$ be the identity element and let $[\mathbf{1}] \in \mathrm{Ext}_{\mathcal{X}}^1(\Omega_{\mathcal{X}/S}(\mathcal{D}), F^{1\vee})$ be the associated element. The element $[\mathbf{1}]$ defines a family $\tilde{\mathcal{X}}/\tilde{S}$ extending the family $\mathcal{X}/S$, using (1.16). It has the following properties:

Firstly, let I be any A -module and let $a \in F^1 \otimes_A I$ be any element. Let $T \triangleq \mathrm{Spec} A * I$ and $\mathcal{X}_T/T$ be the extension of $\mathcal{X}$ defined by the cohomology class $[a] \in \mathrm{Ext}_{\mathcal{X}}^1(\Omega_{\mathcal{X}/S}(\mathcal{D}), I)$ of $a \in F^1 \otimes_A I$. Then $\mathcal{X}_T$ is the pull back of $\tilde{\mathcal{X}}$ under the morphism $T \rightarrow \tilde{S}$ defined by $A * F^{1\vee} \rightarrow A * I$ via $(x, y) \mapsto (x, a(y))$. Namely we have isomorphism $\tilde{\mathcal{X}} \times_{\tilde{S}} T \cong \mathcal{X}_T$.

Secondly, let

$$\tilde{T} \triangleq \mathrm{Spec} A * F^{0\vee} \rightarrow \tilde{S} = \mathrm{Spec} A * F^{1\vee}$$

be the morphism defined by $A * F^{1\vee} \rightarrow A * F^{0\vee}$ via $(x, y) \mapsto (x, d^\vee(y))$. Then we have isomorphism over $\tilde{T}$

$$(1.17) \quad \tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T} \cong \mathcal{X} \times_S \tilde{T},$$

where the projection $\tilde{T} \rightarrow S$ is defined by the obvious inclusion $A \hookrightarrow A * F^{0\vee}$.

As mentioned, $\mathbf{1} \in F^1 \otimes_A F^{1\vee}$ defines an extension $\tilde{\mathcal{X}}/\tilde{S}$, which we fix from now on. We let $\tilde{\mathcal{U}}_\alpha$ (resp. $\tilde{\mathcal{V}}_\alpha$) be the étale neighborhood of $\tilde{\mathcal{X}}$ (resp. $\tilde{S}$) that is the minimal extension of $\mathcal{U}_\alpha$ (resp. $\mathcal{V}_\alpha$)⁷. Then $\{\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha\}$ forms a covering of $\tilde{\mathcal{X}}/\tilde{S}$. By Lemma 1.12, $f_\alpha: \mathcal{U}_\alpha \rightarrow \mathcal{W}_\alpha$ can be extended to a pre-deformable morphism $\tilde{\mathcal{U}}_\alpha \rightarrow W[n]$. For each α we pick one such extension $\zeta_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$ once and for all. Now let $\alpha, \beta \in \Lambda$ be a pair so that $\mathcal{U}_{\alpha\beta} \neq \emptyset$. Let $\tilde{\mathcal{U}}_{\alpha\beta} = \tilde{\mathcal{U}}_\alpha \times_{\tilde{\mathcal{X}}} \tilde{\mathcal{U}}_\beta$ and let

⁶In this section we will denote by $A * I$ the trivial ring extension of A by the A -module I .

⁷We say $\tilde{\mathcal{U}}_\alpha$ is a minimal extension of the étale neighborhood $\mathcal{U}_\alpha$ if $\tilde{\mathcal{U}}_\alpha \rightarrow \tilde{\mathcal{X}}$ is an étale neighborhood and $\tilde{\mathcal{U}}_\alpha \times_{\tilde{\mathcal{X}}} \tilde{\mathcal{X}} \cong \mathcal{U}_\alpha$.

$\tilde{p}_\alpha : \tilde{\mathcal{U}}_{\alpha\beta} \rightarrow \tilde{\mathcal{U}}_\alpha$ be the projection, following our convention. Since both $\zeta_\alpha \circ \tilde{p}_\alpha$ and $\zeta_\beta \circ \tilde{p}_\beta$ are pre-deformable extensions of $f_{\alpha\beta} : \mathcal{U}_{\alpha\beta} \rightarrow W[n]$, by Lemma 1.12 their difference defines an element

$$\zeta_{\alpha\beta} \triangleq \mathbf{d}^\dagger(\zeta_\beta \circ \tilde{p}_\beta - \zeta_\alpha \circ \tilde{p}_\alpha) \in \mathrm{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{W[n]}, F^{1\vee})^\dagger.$$

This defines a homomorphism

$$(1.18) \quad \zeta_{\alpha\beta}(\cdot) : F^1 \longrightarrow \mathrm{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{W[n]}, A)^\dagger.$$

Now to each α we construct a homomorphism

$$(1.19) \quad \zeta_\alpha(\cdot) : F^0 \longrightarrow \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, A)^\dagger.$$

Let $\mathbf{1} \in F^0 \otimes_A F^{0\vee}$ be the identity element. Then $d(\mathbf{1}) \in F^1 \otimes_A F^{0\vee}$ (d is the differential in the complex $F^\bullet$) defines a homomorphism $d(\mathbf{1}) : F^{1\vee} \rightarrow F^{0\vee}$ which induces a morphism

$$\tau : T \triangleq \mathrm{Spec} A * F^{0\vee} \rightarrow \mathrm{Spec} A * F^{1\vee} = \tilde{S}.$$

Let $\mathcal{X}_T/T$ be the pull back family of $\tilde{\mathcal{X}}/\tilde{S}$ via τ with $q : \mathcal{X}_T \rightarrow \tilde{\mathcal{X}}$ the induced the projection. By the second property of the complex $F^\bullet$, $\mathcal{X}_T$ is isomorphic to $\mathcal{X} \times_S T$, where $T \rightarrow S$ is induced by the obvious inclusion $A \rightarrow A * F^{0\vee}$. We let $q_0 : \mathcal{X}_T \rightarrow \mathcal{X}$ be the induced projection. Clearly, $U_\alpha = \tilde{\mathcal{U}}_\alpha \times_{\tilde{S}} T$ is canonically isomorphic to $\mathcal{U}_\alpha \times_S T$ under the isomorphism $\mathcal{X}_T \cong \mathcal{X} \times_S T$. Then restricting to U_α , both $\zeta_\alpha \circ q$ and $f \circ q_0$ are pre-deformable extensions of f_α . By Lemma 1.12, the difference

$$\mathbf{d}^\dagger(\zeta_\alpha \circ q - f \circ q_0) \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, F^{0\vee})^\dagger$$

It defines the desired homomorphism $\zeta_\alpha(\cdot)$ in (1.19).

Lemma 1.14. *Both $\zeta_\alpha(\cdot)$ and $\zeta_{\alpha\beta}(\cdot)$ are homomorphisms of A -modules. Further, for any $a \in F^0$ we have*

$$-p_\alpha^* \zeta_\alpha(a) + p_\beta^* \zeta_\beta(a) = \zeta_{\alpha\beta}(d(a)) \in \mathrm{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{W[n]}, A)^\dagger,$$

where $p_\alpha : \mathcal{U}_{\alpha\beta} \rightarrow \mathcal{U}_\alpha$ is the projection and $p_\alpha^* \zeta_\alpha(\cdot)$ is the pull back homomorphism.

Proof. The proof is straightforward and will be omitted. $\square$

In the next part, we will construct the complex that will be a part of the perfect obstruction theory of $S \subset \mathfrak{M}(W[n], \Gamma)^{\mathrm{st}}$. Let I be any A -module. We let $\mathbf{D}(I)^\bullet$ be the Čech complex

$$\mathbf{D}(I)^\bullet = \mathbf{C}^\bullet(\Lambda, \mathrm{Hom}(f^*\Omega_{W[n]}, I)^\dagger)$$

associating to the covering Λ , where

$$\Gamma(\mathcal{U}_{\alpha_1 \dots \alpha_k}, \mathrm{Hom}(f^*\Omega_{W[n]}, I)^\dagger) = \mathrm{Hom}_{\mathcal{U}_{\alpha_1 \dots \alpha_k}}(f^*\Omega_{W[n]}, I)^\dagger$$

with $\partial : \mathbf{D}(I)^\bullet \rightarrow \mathbf{D}(I)^{\bullet+1}$ the coboundary differential in the Čech complex. We let

$$\delta_0 : F^0 \otimes_A I \longrightarrow \mathbf{C}^0(\Lambda, \mathrm{Hom}(f^*\Omega_{W[n]}, I)^\dagger)$$

be defined by $\delta(a)_\alpha = \zeta_\alpha(a)$ and let

$$\delta_1 : F^1 \otimes_A I \longrightarrow \mathbf{C}^1(\Lambda, \mathrm{Hom}(f^*\Omega_{W[n]}, I)^\dagger)$$

be defined by $\delta_1(b)_{\alpha\beta} = \zeta_{\alpha\beta}(b)$. By Lemma 1.14, δ_i are homomorphisms of A -modules. We now show that this defines a homomorphism of complexes

$$\delta : F^\bullet \otimes_A I \longrightarrow \mathbf{D}(I)^\bullet.$$

To prove this, we need to check that $\delta_1 \circ d = \partial \circ \delta_0$ on $F^0 \otimes_A I$ and $\partial \circ \delta_1 = 0$ on $F^1 \otimes_A I$. We will check $\partial \circ \delta = 0$ on $F^1 \otimes_A I$ and leave the other to the readers. Let $b \in F^1 \otimes_A I$ be any element. By definition,

$$(\partial \circ \delta_1)(b)_{\alpha\beta\gamma} = \delta_1(b)_{\alpha\beta} - \delta_1(b)_{\alpha\gamma} + \delta_1(b)_{\beta\gamma} = \zeta_{\alpha\beta}(b) - \zeta_{\alpha\gamma}(b) + \zeta_{\beta\gamma}(b)$$

as elements in $\text{Hom}_{\mathcal{U}_{\alpha\beta\gamma}}(f^*\Omega_{W[n]}, I)^\dagger$, where the summation is taken after pulling back each term in the summation to this module in the obvious way. But this vanishes because of the relation $\zeta_{\beta\gamma} - \zeta_{\alpha\gamma} + \zeta_{\alpha\beta} = 0$, following Remark 1.13. This shows that $\partial \circ \delta_1 = 0$ on $F^1 \otimes_A I$.

In the end, we define the complex $\mathbf{E}(I)^\bullet$ by

$$(1.20) \quad \mathbf{E}(I)^k = \bigoplus_{i+j=k} (F^i \otimes_A I \oplus \mathbf{D}(I)^{j-1})$$

with the differential $d_{\mathbf{E}}: \mathbf{E}(I)^k \rightarrow \mathbf{E}(I)^{k+1}$ defined so that its restriction to $F^0 \otimes_A I$ and $F^1 \otimes_A I$ are $d \oplus -\delta_0$ and δ_1 , respectively, and its restriction to $\mathbf{D}(I)^i$ is ∂ . When $I = A$, we will abbreviate $\mathbf{D}(A)^\bullet$ and $\mathbf{E}(A)^\bullet$ to $\mathbf{D}^\bullet$ and $\mathbf{E}^\bullet$ respectively.

Lemma 1.15. *We assume Λ is a sufficiently fine cover of f . Then for any A -module I , $\mathbf{E}(I)^\bullet$ is canonically isomorphic to $\mathbf{E}^\bullet \otimes_A I$ as complexes. Further, the complex $\mathbf{E}^\bullet$ is a complex of flat A -modules.*

Proof. The proof is straightforward and will be omitted. $\square$

We remark that the complex $\mathbf{E}^\bullet$ just constructed depends on the choice of the atlas Λ . To emphasize this dependence we shall denote it by $\mathbf{E}_\Lambda^\bullet$. Let Λ' be any atlas of f that is a refinement of Λ with associated complex $\mathbf{E}_{\Lambda'}^\bullet$. Following [Mil, III.2], we can define a homomorphism of complexes $\mathbf{E}_\Lambda^\bullet \rightarrow \mathbf{E}_{\Lambda'}^\bullet$ which induces a map of cohomologies

$$\rho(\Lambda, \Lambda') : h^i(\mathbf{E}_\Lambda^\bullet \otimes_A I) \longrightarrow h^i(\mathbf{E}_{\Lambda'}^\bullet \otimes_A I)$$

and then form the direct limit $\lim_{\rightarrow} h^i(\mathbf{E}_\Lambda \otimes_A I)$ taken over all charts of f . Note that this limit is a functor from $\mathfrak{Mod}_A$ to $\mathfrak{Mod}_A$, denoted by $\mathfrak{h}^i(\mathbf{E}^\bullet)$. We now assume Λ is fine enough so that $\mathfrak{h}^i(\mathbf{E}^\bullet)(I) = h^i(\mathbf{E}_\Lambda^\bullet \otimes_A I)$ for all A -module I . We fix such a Λ once and for all, and abbreviate the resulting complex $\mathbf{E}_\Lambda^\bullet$ to $\mathbf{E}^\bullet$.

We now prove the main result of this section.

Proposition 1.16. *Let $\mathfrak{Def}_A^1$ be the functor of the first order deformations of morphisms to $S = \text{Spec } A$, which is naturally a functor from A -modules to A -modules. Then $\mathfrak{Def}_A^1$ is naturally isomorphic to the functor $\mathfrak{h}^1(\mathbf{E}^\bullet)$.*

Proof. Let $\xi = (B, I)$ be any object in $\mathfrak{Mod}_A$. We first show that there is a canonical isomorphism $\mathfrak{Def}_A^1(\xi) \cong h^1(\mathbf{E}^\bullet \otimes_A I)$.

Let $T = \text{Spec } B$ and $\tilde{T} = \text{Spec } B * I$, the trivial extension by I . Since B is an A -algebra, there is a tautological morphism $T \rightarrow S$. Let $x \in \mathfrak{Def}_S^1(\xi)$ ⁸ be any element, associated to an extension $\tilde{T} \rightarrow S$ of $T \rightarrow S$, and let $f_T : \mathcal{X}_T \rightarrow W[n]$ (resp. $f_{\tilde{T}} : \mathcal{X}_{\tilde{T}} \rightarrow W[n]$) be the pull back family of f via $T \rightarrow S$ (resp. $\tilde{T} \rightarrow S$). We let $\mathcal{D}_T \subset \mathcal{X}_T$ and $\mathcal{D}_{\tilde{T}} \subset \mathcal{X}_{\tilde{T}}$ be the associated divisors of marked points and let $\mathcal{I} = \mathcal{O}_{\mathcal{X}_T} \otimes_B I$. First of all, since $\mathcal{X}_{\tilde{T}}$ is a flat extension of $\mathcal{X}_T$ to $\tilde{T}$, it associates to a unique element in

$$(1.21) \quad \text{Ext}_{\mathcal{X}_T}^1(\Omega_{\mathcal{X}_T/T}(\mathcal{D}_T), \mathcal{I}) \cong h^1(F^\bullet \otimes_A I).$$

⁸We use the convention $\mathfrak{Def}_S = \mathfrak{Def}_A$ when $S = \text{Spec } A$.

Hence it is represented by $[a] \in h^1(F^\bullet \otimes_A I)$ for an $a \in F^1 \otimes_A I$. Now let $\tilde{S}$ be the trivial extension of S by $F^{1\vee}$ and let $\tilde{\mathcal{X}}/\tilde{S}$ be the family defined before (1.17). Then a defines a morphism $\varphi_a: \tilde{T} \rightarrow \tilde{S}$ via the homomorphism $A \oplus F^{1\vee} \rightarrow B \oplus I$ defined by $(x, y) \mapsto (\bar{x}, a(y))$, where $\bar{x}$ is the image of x in B . By the first property of (1.16), a defines an isomorphism $\gamma_a: \mathcal{X}_{\tilde{T}} \cong \tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T}$, where the projection $\tilde{T} \rightarrow \tilde{S}$ is via φ_a . We let $\tilde{U}_\alpha = \tilde{\mathcal{U}}_\alpha \times_{\tilde{S}} \tilde{T}$ and $\tilde{V}_\alpha = \tilde{\mathcal{V}}_\alpha \times_{\tilde{S}} \tilde{T}$. Then each $\tilde{U}_\alpha/\tilde{V}_\alpha$ extends to a chart $(\tilde{U}_\alpha/\tilde{V}_\alpha, f_{\tilde{T},\alpha})$ of $\tilde{T}$, where $f_{\tilde{T},\alpha} \triangleq f_{\tilde{T}}|_{\tilde{U}}$. We let $\tilde{\rho}_\alpha: \tilde{U}_\alpha \rightarrow \tilde{\mathcal{U}}_\alpha$ be the projection. Then over each $\tilde{U}_\alpha$ we have two pre-deformable extensions of f_T : One is $f_{\tilde{T},\alpha}$ and the other is the composite $\zeta_\alpha \circ \tilde{\rho}_\alpha$. Let

$$b_\alpha = \mathbf{d}^\dagger(f_{\tilde{T},\alpha} - \zeta_\alpha \circ \tilde{\rho}_\alpha) \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f_T^* \Omega_{W[n]}, I)^\dagger$$

and let $b = \{b_\alpha\} \in \mathbf{D}^0 \otimes_A I$. We claim that $(a, b) \in \mathbf{E}^1 \otimes_A I$ is in the kernel of $d_{\mathbf{E}}$. For this, we only need to check that

$$d_{\mathbf{E}}(a, b)_{\alpha\beta} = \delta(a)_{\alpha\beta} + \partial(b)_{\alpha\beta} = \zeta_{\alpha\beta}(a) + (b_\beta - b_\alpha) = \zeta_{\alpha\beta}(a) - \mathbf{d}^\dagger(\zeta_\beta \circ \tilde{\rho}_\beta - \zeta_\alpha \circ \tilde{\rho}_\alpha)$$

vanishes for all pairs (α, β) . But this follows immediately from the definition of $\zeta_{\alpha\beta}(\cdot)$. This shows that (a, b) defines a cohomology class $[(a, b)] \in h^1(\mathbf{E}^\bullet \otimes_A I)$.

Next, we show that $[(a, b)]$ is independent of the choices of a and the isomorphisms $\gamma_a: \mathcal{X}_{\tilde{T}} \cong \tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T}$. Let $a' \in F^1 \times_A I$ be another element so that $[a'] = [a]$ in (1.21). Then $a - a' = d_I(c)$ for a $c \in F^0 \otimes_A I$, where $d_I: F^0 \otimes_A I \rightarrow F^1 \otimes_A I$ is induced by d in (1.15). Now let

$$\varphi_{a'}: \tilde{T} \rightarrow \tilde{S}, \quad \gamma_{a'}: \mathcal{X}_{\tilde{T}} \cong \tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T}, \quad \tilde{\rho}'_\alpha: \tilde{U}_\alpha \rightarrow \tilde{\mathcal{U}}_\alpha, \quad b'_\alpha = \mathbf{d}^\dagger(f_{\tilde{T},\alpha} - \zeta_\alpha \circ \tilde{\rho}'_\alpha)$$

be objects defined similarly with a replaced by a' . Let $b' = \{b'_\alpha\}$. We claim that

$$d_{\mathbf{E}}(c) = (a, b) - (a', b') \in \mathbf{E}^1 \otimes_A I.$$

Once this is established then $[(a, b)] = [(a', b')] \in h^1(\mathbf{E}^\bullet \otimes_A I)$, which shows that $[(a, b)]$ only depends on the class $x \in \mathfrak{Def}_S^1(v)$. This way we obtain a map

$$\mathbf{T}(\xi): \mathfrak{Def}_S^1(v) \longrightarrow h^1(\mathbf{E}^\bullet \otimes_A I).$$

Now we prove the claim. Since

$$d_{\mathbf{E}}(c) = (\partial(c), -\delta(c)) = (d_I(c), -\delta(c))$$

and $d_I(c) = a - a'$, it suffices to show that $\delta(c)_\alpha = \zeta(c)_\alpha$ is identical to

$$-(b - b')_\alpha = -(b_\alpha - b'_\alpha) = -\mathbf{d}^\dagger(f_{\tilde{T},\alpha} - \zeta_\alpha \circ \tilde{\rho}_\alpha) + \mathbf{d}^\dagger(f_{\tilde{T},\alpha} - \zeta_\alpha \circ \tilde{\rho}'_\alpha) = \mathbf{d}^\dagger(\zeta_\alpha \circ \tilde{\rho}_\alpha - \zeta_\alpha \circ \tilde{\rho}'_\alpha).$$

The proof that $\zeta_\alpha(c) = \mathbf{d}^\dagger(\zeta_\alpha \circ \tilde{\rho}_\alpha - \zeta_\alpha \circ \tilde{\rho}'_\alpha)$ is routine and will be omitted.

We now show that the map $\mathbf{T}(\xi)$ is one-one and onto. We give an outline of the proof since it is standard. We first show that it is one-one. Let $x \in \mathfrak{Def}_S^1(\xi)$ be any element so that $\mathbf{T}(\xi)(x) = 0 \in h^1(\mathbf{E}^\bullet \otimes_A I)$. We let (a, b) be the pair constructed associated to the family $f_{\tilde{T}}: \mathcal{X}_{\tilde{T}} \rightarrow W[n]$ following the previous discussion. Since $[(a, b)] = 0$, there is a $c \in F^0 \otimes_A I$ so that $(a, b) = d_{\mathbf{E}}(c)$. This implies at first that $a = d_I(c)$. Hence $\mathcal{X}_{\tilde{T}} \cong \mathcal{X}_T \times_T \tilde{T}$ under the obvious projection $\tilde{T} \rightarrow T$ via $B \rightarrow B \oplus I$. Let $\rho: \mathcal{X}_{\tilde{T}} \rightarrow \mathcal{X}_T$ be the projection. Because $\mathbf{T}(\xi)(x)$ is well-defined, $\mathbf{T}(\xi)(x)$ is also represented by $(0, b')$, where $b' = \{b'_\alpha\}$ and

$$b'_\alpha = \mathbf{d}^\dagger(f_{\tilde{T},\alpha} - f_{T,\alpha} \circ \rho_\alpha) \in \mathrm{Hom}_{U_\alpha}(f_T^* \Omega_{W[n]}, I)^\dagger.$$

Again, since $\mathbf{T}(\xi)(x) = 0$, there must be a $c \in F^0 \otimes_A I$ so that $d_I(c) = 0$ and $\delta(c) = b'$. Hence c lifts to an element in $\mathrm{Ext}_{\mathcal{X}_T}^0(\Omega_{\mathcal{X}_T/T}(\mathcal{D}_T), \mathcal{I})$, which defines a

new isomorphism $\gamma: \mathcal{X}_{\tilde{T}} \cong \mathcal{X}_T \times_T \tilde{T}$. Further, $\delta(c) = b'$ implies that under this new isomorphism $f_{\tilde{T}}$ is the constant extension of f_T . Namely, $f_{\tilde{T}}$ is the pull back of f_T via the projection $\tilde{T} \rightarrow T$. This proves that $x = 0$ in $\mathfrak{Def}_S^1(\xi)$ and hence $\mathbf{T}(\xi)$ is one-one.

The onto part is similar. Since this argument is standard in deformation theory, we will omit it here. In the end, we need to check that $\mathbf{T}(\xi)$ is a homomorphism of modules and that $\mathbf{T}$ is an isomorphism of functors. This is straightforward and will be omitted. $\square$

Corollary 1.17. *Let $\xi = (B, I, \varphi_0)$ be any object in $\mathfrak{Tri}_S$. Suppose $\mathfrak{Def}_S(\xi) \neq \emptyset$, then $\mathfrak{Def}_S(\xi)$ is isomorphic to the set $h^1(\mathbf{E}^\bullet \otimes_A I)$.*

Proof. Since S is an étale chart of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$, which is a Deligne-Mumford stack, the standard fact in deformation theory shows that once an extension $\varphi: \text{Spec } B \rightarrow S$ of φ_0 is fixed, then the space of all such extensions is canonically isomorphic to $\mathfrak{Def}_S^1(\xi_0) \cong h^1(\mathbf{E}^\bullet \otimes_A I)$, where $\xi_0 = (B/I * I, I, \varphi_0)$. This proves the corollary. $\square$

Proposition 1.18. *There is a natural obstruction theory to deformation of the family of pre-deformable $f: \mathcal{X} \rightarrow W[n]$ over S that takes values in $\mathfrak{h}^2(\mathbf{E}^\bullet)$.*

Proof. The construction of the obstruction class is standard, as shown in [LT1] for smooth targets. Let $\xi = (B, I, \varphi)$ be an object in $\mathfrak{Tri}_S$. Let $T = \text{Spec } B/I$ and let $f_T: \mathcal{X}_T \rightarrow W[n]$ with $\mathcal{D}_T \subset \mathcal{X}_T$ be the divisor of marked points be the pull back of f via $T \rightarrow S$. Then to find an extension $\tilde{r}: \tilde{T} \rightarrow S$ of $T \rightarrow S$ is equivalent to extend the family f_T to a family of pre-deformable morphisms over $\tilde{T}$. We now construct the obstruction class to this extension problem.

Since deformation of pointed curves is unobstructed, we can extend $\mathcal{X}_T$ to a family over $\tilde{T}$. By our choice of $F^\bullet$, such extension can be realized by an extension $\tilde{T} \rightarrow \tilde{S}$ of $T \rightarrow S$. We let $\mathcal{X}_{\tilde{T}}$ be the pull back family $\tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T}$. Let $\tilde{U}_\alpha = \tilde{U}_\alpha \times_{\tilde{S}} \tilde{T}$, $\tilde{V}_\alpha = \tilde{V}_\alpha \times_{\tilde{S}} \tilde{T}$, $U_\alpha = \tilde{U}_\alpha \times_{\mathcal{X}_{\tilde{T}}} \mathcal{X}_T$ and let $V_\alpha = \tilde{V}_\alpha \times_{\mathcal{X}_{\tilde{T}}} \mathcal{X}_T$. Also we extend U_α/V_α to a chart $(U_\alpha/V_\alpha, f_{T,\alpha}, \mathcal{W}_\alpha)$ of f_T . This way $\tilde{U}_\alpha/\tilde{V}_\alpha$ is a minimal extension of U_α/V_α to $\mathcal{X}_{\tilde{T}}/\tilde{T}$. Next we let $\tilde{\rho}_\alpha: \tilde{U}_\alpha \rightarrow \tilde{U}_\alpha$ be the projection. For any α we pick a family of pre-deformable extension $\tilde{h}_\alpha: \tilde{U}_\alpha \rightarrow W[n]$ of $f_{T,\alpha}$. We let $\tilde{p}_\alpha: \tilde{U}_{\alpha\beta} \rightarrow \tilde{U}_\alpha$ be the projection. We then let

$$(1.22) \quad b_{\alpha\beta} = -\mathbf{d}^\dagger(\tilde{h}_\beta \circ \tilde{p}_\beta - \tilde{h}_\alpha \circ \tilde{p}_\alpha) \in \text{Hom}_{U_{\alpha\beta}}(f_T^* \Omega_{W[n]}, I)^\dagger.$$

We let $b = \{b_{\alpha\beta}\}$, which belongs to $\mathbf{D}^1 \otimes_A I \subset \mathbf{E}^2 \otimes_A I$. It follows from the Remark 1.13 that b is a cocycle, and thus defines a cohomology class $[b]$ in $h^2(\mathbf{E}^\bullet \otimes_A I)$. The technical part of the proof is to check that the cohomology class $[b]$ is independent of the choices of $\tilde{T} \rightarrow \tilde{S}$ and $\tilde{h}_\alpha$. The argument for this is straightforward though tedious, and will be omitted.

We now show that it is an obstruction class to extending f_T to families of pre-deformable morphisms over $\tilde{T}$. First, if such extensions do exist, then we can choose $\mathcal{X}_{\tilde{T}}$ and $\tilde{h}_\alpha$ be data coming from one of such extensions. Then the corresponding $b' = 0$ as cocycle and thus $[b] = [b'] = 0$. This shows that $[b] = 0$ whenever extensions of f_T exist. Now assume $[b] \neq 0$. We first look at the case where the cycle $b \in \mathbf{D}^1 \otimes_A I$ is a coboundary in $\mathbf{D}^\bullet \otimes_A I$. Namely, there is a $c \in \mathbf{D}^0 \otimes_A I$ so that $b = \partial(c)$. Let $c = \{c_\alpha\}$ with $c_\alpha \in \text{Hom}_{U_\alpha}(f_T^* \Omega_{W[n]}, I)^\dagger$. Then by Lemma 1.12, we

can find pre-deformable extension $f_{\tilde{T},\alpha}:\tilde{U}_\alpha \rightarrow W[n]$ of $f_{T,\alpha}$ so that $\mathbf{d}^\dagger(f_{\tilde{T},\alpha} - \tilde{h}_\alpha) = c_\alpha$. Then over $\tilde{U}_{\alpha\beta}$, the difference of the pull backs $f_{\tilde{T},\beta} \circ \tilde{p}_\beta$ and $f_{\tilde{T},\alpha} \circ \tilde{p}_\alpha$ is

$$\begin{aligned} \mathbf{d}^\dagger(f_{\tilde{T},\beta} \circ \tilde{p}_\beta - f_{\tilde{T},\alpha} \circ \tilde{p}_\alpha) &= \mathbf{d}^\dagger(f_{\tilde{T},\beta} \circ \tilde{p}_\beta - \tilde{h}_\beta \circ \tilde{p}_\beta) - \mathbf{d}^\dagger(f_{\tilde{T},\alpha} \circ \tilde{p}_\alpha - \tilde{h}_\alpha \circ \tilde{p}_\alpha) + \\ &\quad + \mathbf{d}^\dagger(\tilde{h}_\beta \circ \tilde{p}_\beta - \tilde{h}_\alpha \circ \tilde{p}_\alpha) = (c_\beta - c_\alpha) - b_{\alpha\beta} = 0. \end{aligned}$$

Hence $\{f_{\tilde{T},\alpha}\}$ patch together to form a desired extension $\mathcal{X}_{\tilde{T}} \rightarrow W[n]$ of f_T .

In general, assume $[b] = 0$, then there is an $a \in F^0 \otimes_A I$ and $c \in \mathbf{D}^0 \otimes_A I$ so that

$$b = d_{\mathbf{E}}(a, c) = \delta(a) + \partial(c).$$

Now let $\tilde{r}': \tilde{T} \rightarrow \tilde{S}$ be a new extension of $T \rightarrow S$ so that $\mathbf{d}(\tilde{r}' - \tilde{r}) = a$, where $\tilde{r}: \tilde{T} \rightarrow \tilde{S}$ is the morphism used to construct the cycle b . One checks that if one uses the new extension $\tilde{\mathcal{X}} \times_{\tilde{S}} \tilde{T}$, where $\tilde{T} \rightarrow \tilde{S}$ is via $\tilde{r}'$, to construct a similar cycle $b' \in \mathbf{D}^1 \otimes_A I$, then b' is a coboundary in $\mathbf{D}^\bullet \otimes_A I$. This reduces the situation to the previous case studied, and hence confirms that a pre-deformable extension of f_T over $\tilde{T}$ can be found. This shows that $[b]$ is an obstruction class to extending f_T to families of pre-deformable morphisms over $\tilde{T}$, or equivalently the obstruction class to extending $T \rightarrow S$ to $\tilde{T} \rightarrow S$.

We define

$$\mathfrak{ob}(B, I, \varphi) = [b] \in h^2(\mathbf{E}^\bullet \otimes_A I).$$

Again it is direct to check that this assignment defines a functor from $\mathfrak{Tri}_S$ to $\mathfrak{Mod}_S^*$. This completes the proof of the Proposition. $\square$

We now summarize the result of this section in the following theorem. We need a vanishing Lemma whose proof will be provided in Proposition 5.1.

Lemma 1.19. *For sufficiently fine Λ , we have $h^i(\mathbf{D}^\bullet \otimes_A I) = 0$ for any A -module I and $i \geq 2$.*

Theorem 1.20. *Let S be an affine chart of the moduli stack $\mathfrak{M}(W[n], \Gamma)^{st}$. Then the obstruction theory just defined is a perfect obstruction theory of S .*

Proof. Let $f: \mathcal{X} \rightarrow W[n]$ be the universal family over S . It suffices to show that there is a complex of finitely generated free A -modules $E^\bullet = [E^1 \rightarrow E^2]$ so that it is quasi-isomorphic to $\mathbf{E}^\bullet$, where $\mathbf{E}^\bullet$ is the complex associated to a sufficiently fine atlas Λ of f . Since $h^i(\mathbf{D}^\bullet \otimes_A I) = 0$ for $i \geq 2$ and any I , $h^i(\mathbf{E}^\bullet \otimes_A I) = 0$ for $i \geq 3$ and any I . Hence there is a bounded subcomplex $\tilde{E}^\bullet$ of flat A -modules so that it is quasi-isomorphic to $\mathbf{E}^\bullet$. Then we can apply the standard technique [Har, III.12] to find a bounded subcomplex $E^\bullet$ of finitely generated free A -modules that is quasi-isomorphic to $\mathbf{E}^\bullet$. Finally, since $h^i(\mathbf{E} \otimes_A I) \neq 0$ only for $i = 1$ and 2 , we can choose $E^\bullet$ to be of the form $[E^1 \rightarrow E^2]$. This proves the Theorem. $\square$

In the Appendix, we will express the cohomology $h^\bullet(\mathbf{E}^\bullet)$ in terms of some known cohomologies.

1.3. Obstruction to deforming relative stable morphisms. We will follow the notation developed in [Li, section 4] concerning relative stable morphisms. Let (Z, D) be a polarized relative pair and let $Z[n]^{\text{rel}}$ be the expanded relative pair constructed there. Recall that $Z[n]^{\text{rel}}$ consists of a proper variety $Z[n]$ over $\mathbf{A}^n$ and a smooth divisor $D[n] \subset Z[n]$ that is isomorphic to $D \times \mathbf{A}^n$ under the projection $Z[n] \rightarrow Z \times \mathbf{A}^n$. The pair $(Z[n], D[n])$ over $\mathbf{A}^n$ also admits an equivariant $G[n]$

action ($= G_m^{\times n}$) whose action on $\mathbf{A}^n$ is the standard one⁹. The fibers of $Z[n]/\mathbf{A}^n$ has at most normal crossing singularities and the singular locus of all the fibers of $Z[n]/\mathbf{A}^n$ is a disjoint union of smooth varieties $\mathbf{B}_1, \dots, \mathbf{B}_n$, indexed so that B_l surjects to the l -th coordinate hyperplane $H_l \subset \mathbf{A}^n$.

In [Li] we used admissible graph to describe the topological type of relative stable morphisms to $(Z[n], D[n])$. Recall that a weighted graph Γ (introduced in [Li]) consists of a collection of vertices V_Γ , an ordered collection of weighted roots R_Γ and an ordered collection of legs L_Γ (the later two are half edges with one ends attached to vertices and the other ends left free) plus two weight functions $g, b: V_\Gamma \rightarrow \mathbb{Z}_{\geq 0}$ and a multiplicity assignment $\mu: R_\Gamma \rightarrow \mathbb{Z}^+$. We require Γ to be relatively connected in the sense that either Γ is connected or each of its vertex has at least one root attached to it. Recall that a relative morphism to $Z[n]^{\text{rel}}$ of type Γ consists of a pointed complete nodal curve X and a morphism $f: X \rightarrow Z[n]$ that has the following property: First the marked points of X are labeled by the roots and the legs of Γ . For consistency, we will reserve the notation $p_i \in X$ for $i = 1, \dots, k$ and $q_j \in X$ for $j = 1, \dots, r$ to denote the marked points associated to the ordered legs and roots of Γ . we will use $\mu_1, \dots, \mu_r$ to denote the weights of the respective roots of Γ ; Secondly, the connected components of X are labeled by $a \in V_\Gamma$ and the arithmetic genus of the component X_a is $g(a)$. Further, in case a root or a leg is attached to a vertex a then its associated marked point must lie in the connected component X_a ; Thirdly, restricting to each connected component X_a the morphism $f|_{X_a}$ with all the marked points in X_a is an ordinary stable morphism to $Z[n]$ of degree $b(a)$ (using the polarization on Z chosen implicitly); Lastly, as divisor $f^{-1}(D[n]) = \mu_1 q_1 + \dots + \mu_r q_r$. Here we implicitly assume the former is a proper divisor.

We recall the notion of relative stable morphisms to $\mathfrak{Z}^{\text{rel}}$ defined in [Li]. Let $f: X \rightarrow Z[n]$ be a relative morphism of type Γ , as described. Recall fibers of $Z[n]/\mathbf{A}^n$ have normal crossing singularities along $\mathbf{B}_1, \dots, \mathbf{B}_n$. We say f is pre-deformable if it is pre-deformable along all $\mathbf{B}_i$, as defined in [Li, Section 2]. We say f is stable as a relative morphism to $\mathfrak{Z}^{\text{rel}}$ if f is pre-deformable and $\text{Aut}_3(f)$ is finite. Here $\text{Aut}_3(f)$ is the group of all pairs (a, b) where $a: X \rightarrow X$ are automorphisms and $b \in G[n]$ so that $f \circ a = f^b$. (Here we view b as an automorphism $b: Z[n] \rightarrow Z[n]$ using the $G[n]$ action on $Z[n]$ and f^b is the composite of f with b .) As was proved in [Li], the moduli of all relative morphisms to $Z[n]^{\text{rel}}$ of type Γ that are stable as morphisms to $\mathfrak{Z}^{\text{rel}}$ form a Deligne-Mumford stack. We will denote this stack by $\mathfrak{M}(Z[n]^{\text{rel}}, \Gamma)^{\text{st}}$. The goal of this subsection is to describe the obstruction theory of this moduli stack.

Similar to the case $f: \mathcal{X} \rightarrow W[n]$, for any family of relative stable morphisms $f: \mathcal{X} \rightarrow Z[n]$ over S there is a canonical log structure on $\mathcal{X}/S$ and on $Z[n]/\mathbf{A}^n$ that makes f a morphism between log schemes. First, the log structure on $\mathbf{A}^n$ (resp. $Z[n]$) is given by the divisor $\cup_{l=1}^n H_l \subset \mathbf{A}^n$ (resp. $Z[n] \times_{\mathbf{A}^1} 0 \cup D[n] \subset Z[n]$). As to the log structure on $\mathcal{X}/S$, we first note that if we let $W/\mathbf{A}^1$ be $Z[1]/\mathbf{A}^1$, then $Z[n]/\mathbf{A}^n = W[n-1]/\mathbf{A}^n$ and f is a family of pre-deformable morphisms to $W[n-1]$. We let $(\mathcal{X}, \mathcal{M}')$ and $(S, \mathcal{N})$ be the associated log structures of $f: \mathcal{X} \rightarrow W[n-1]$. On the other hand, let r be the number of distinguished marked points of f . Then

⁹In short, $Z[1]$ is the blowing up of $Z \times \mathbf{A}^1$ along $D \times 0$ and $D[1]$ is the proper transform of $D \times \mathbf{A}^1$. $Z[2]$ is the blowing up of $Z[1] \times \mathbf{A}^1$ along $D[1] \times \mathbf{A}^1$, etc. The $G[n]$ action is the unique lifting of its standard action on $Z \times \mathbf{A}^n$.

locally the log structure of $f^{-1}(D[n]) \subset \mathcal{X}$ is given by a pre-log structure $\mathbb{N}^r \rightarrow \mathcal{O}_U$ for open $U \subset \mathcal{X}$. The the desired log structure of $\mathcal{X}$ over U is given by the associated log structure of the pre-log structure $\mathcal{M}'|_U \oplus \mathbb{N}^r \rightarrow \mathcal{O}_U$. It is obvious that this gives $\mathcal{X}$ a log structure $\mathcal{M}$, making it a log-scheme over $(S, \mathcal{N})$ and making f a log-morphism between $\mathcal{X}^\dagger/S^\dagger \rightarrow Z[n]^\dagger/\mathbf{A}^{n\dagger}$.

Let $S = \text{Spec } A$ be an affine chart of $\mathfrak{M}(Z[n]^{\text{rel}}, \Gamma)^{st}$ with $f : \mathcal{X} \rightarrow Z[n]$ the universal family and $q_i, p_j : S \rightarrow \mathcal{X}$ its marked sections. We cover f by charts of the first or the second kind. Let these charts be $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha, \mathcal{Z}_\alpha)$ indexed by Λ as defined before (1.5) with $W[n]$ (resp. $\mathbf{D}_l$; resp. $n+1$) replaced by $Z[n]$ (resp. $\mathbf{B}_l$; resp. n). We let $\mathcal{D} \subset \mathcal{X}$ be the divisor of the locus of all marked sections of $\mathcal{X}/S$. We now fix a complex of finite rank free A -modules $F^\bullet = [F^0 \xrightarrow{d} F^1]$ so that

$$h^\bullet(F^\bullet \otimes_A I) = \text{Ext}_{\mathcal{X}}^\bullet(\Omega_{\mathcal{X}/S}(\mathcal{D}), \mathcal{I}), \quad \mathcal{I} = I \otimes_{\mathcal{O}_S} \mathcal{O}_{\mathcal{X}}$$

for all A -module I . We form the group $\text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{Z[n]}, I)^\dagger$ as follows: In case $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ is a chart away from $f^{-1}(D[n])$, this group is $\text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n-1]}, I)^\dagger$ with $W[n-1]/\mathbf{A}^n = Z[n]/\mathbf{A}^n$ understood. Now let $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ be a chart of some points in $f^{-1}(D[n])$ that is away from $f^{-1}(\mathbf{B})$. Then by shrinking $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ and $\mathcal{Z}_\alpha$ if necessary, we can assume that there is a section $z_\alpha \in \Gamma(\mathcal{O}_{\mathcal{Z}_\alpha})$ so that $z_\alpha = 0$ is the divisor $\mathcal{Z}_\alpha \cap D[n]$. We define $\text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{Z[n]}, I)^\dagger$ be the subgroup of

$$(\varphi, \eta) \in \text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{Z[n]}, \mathcal{I}_\alpha) \oplus \mathcal{I}_\alpha$$

so that

$$\varphi(f^*dz_\alpha) = f_\alpha^*(z_\alpha) \cdot \eta \quad \text{and} \quad \varphi(dt_l) \in I_\alpha, \quad \forall l.$$

Using the A -modules $\text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{Z[n]}, I)^\dagger$, we can form the complex $(\mathbf{D}(I)^\bullet, \partial)$, just as we did for $W[n]$ before (1.20). As before, the element $\mathbf{1} \in F^1 \otimes_A F^{1\vee}$ defines a flat extension of $\mathcal{X}/S$ to $\tilde{\mathcal{X}}/\tilde{S}$, where $\tilde{S} \triangleq \text{Spec } A * F^{1\vee}$, with extended sections $\tilde{q}_i, \tilde{p}_j : \tilde{S} \rightarrow \tilde{\mathcal{X}}$. We let $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ be the minimal extension of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ as an étale neighborhood of $\tilde{\mathcal{X}}/\tilde{S}$. We then pick $\zeta_\alpha : \tilde{\mathcal{U}}_\alpha \rightarrow Z[n]$ that is an extension of $f_\alpha : \mathcal{U}_\alpha \rightarrow Z[n]$ so that ζ_α is pre-deformable and

$$(1.23) \quad \zeta_\alpha^{-1}(D[n]) = \sum_{j=1}^r \mu_j \tilde{q}_j(\tilde{\mathcal{V}}_\alpha).$$

To construct the corresponding complex $\mathbf{E}(I)^\bullet$ we need two homomorphisms

$$(1.24) \quad \zeta_\alpha(\cdot) : F^0 \longrightarrow \text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{Z[n]}, A)^\dagger$$

and

$$(1.25) \quad \zeta_{\alpha\beta}(\cdot) : F^1 \longrightarrow \text{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{Z[n]}, A)^\dagger.$$

First, as before we argue that the difference of ζ_α and ζ_β over $\tilde{\mathcal{U}}_{\alpha\beta}$ canonically defines an element $\zeta_{\alpha\beta} \in \text{Hom}_{\mathcal{U}_{\alpha\beta}}(f^*\Omega_{Z[n]}, F^{1\vee})^\dagger$, which naturally defines a homomorphism as required in (1.25). Here the log differential, namely $f^*(dz_\alpha)/f^*(z_\alpha) \mapsto \eta$, appear near $D[n]$ because of the constraint (1.23). The construction of (1.24) is similar. Namely, locally over $\text{Spec } A * F^{0\vee}$ there are two extensions of f . One is given by the pull back of ζ_α and the other is given by the pull back of f . Their difference then gives rise to the homomorphism (1.24).

Once all such data are constructed, we then go ahead to form the homomorphism of complexes $F^\bullet \otimes_A I \rightarrow \mathbf{D}(I)^\bullet$, form a new complex $\mathbf{E}(I)^\bullet$ and check that there is a

complex of finite rank free A -modules $E^\bullet = [E^1 \rightarrow E^2]$ so that it is quasi-isomorphic to $\mathbf{E}^\bullet$, parallel to the argument in the previous subsection.

We now state the main result of this section.

Lemma 1.21. *Let the notation be as before and let Λ be a sufficiently fine covering of f by charts of the first or the second kinds. Then the complex $\mathbf{E}^\bullet$ is a complex of flat A -modules. Further, for any A -module I we have $\mathbf{E}(I)^\bullet = \mathbf{E}^\bullet \otimes_A I$ and $h^i(\mathbf{E}^\bullet \otimes I) = 0$ for $i \neq 1, 2$. In particular, there is a complex $E^\bullet = [E^1 \rightarrow E^2]$ of finitely generated free A -modules so that it is quasi-isomorphic to $\mathbf{E}^\bullet$.*

Theorem 1.22. *Let the notation be as in the previous Lemma and let $S = \text{Spec } A$ be the affine chart of $\mathfrak{M}(Z[n]^{\text{rel}}, \Gamma)^{\text{st}}$ as before. Then there is a perfect obstruction theory of S taking value in the complex $E^\bullet$. In particular, the functor of the first order deformations $\mathfrak{Def}_S^1$ is isomorphic to $\mathfrak{h}^1(E^\bullet)$ and there is an obstruction assignment $\mathfrak{ob}$ taking values in $\mathfrak{h}^2(E^\bullet)$ that satisfies the required base change property.*

Proof. The proof of the Lemma and the Theorem are parallel to that of Lemmas 1.15, 1.19 and the Theorem 1.20. The only new ingredient is about preserving the divisor $f^{-1}(D[n]) = \sum \mu_j q_j(S)$. Since $q_j(S) \subset \mathcal{X}$ is a divisor smooth over S and $D[n] \subset Z[n]$ is a smooth divisor, that the deformation of morphisms preserving this relation is given by the sheaf of log-differentials is well-known, for example see [Kol]. Since the proof is routine and parallel to what we did before, we shall omit it. This completes the proof of the Theorem. $\square$

2. GROMOV-WITTEN INVARIANTS

In this section, we will define the virtual moduli cycle of $\mathfrak{M}(\mathfrak{W}, \Gamma)$, $\mathfrak{M}(\mathfrak{W}_t, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$, thus defining the Gromov-Witten invariants of the family W , of the singular variety W_0 and the relative Gromov-Witten invariants of the pair (Z, D) . In the next section, we will prove the decomposition (degeneration) formula relating the Gromov-Witten invariants of W_t to the relative Gromov-Witten invariants of the pairs (Y_1, D_1) and (Y_2, D_2) .

2.1. Perfect obstruction theories of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$. Recall that the construction of the virtual cycles of moduli stacks is based on the choice of their perfect obstruction theories. In this section, we will show that the perfect obstruction theories constructed in the previous section naturally induce perfect obstruction theories of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$.

Let $\mathbf{M}$ be a proper Deligne-Mumford stack with an atlas Λ consisting of finitely many affine étale morphisms $\iota_\alpha: S_\alpha = \text{Spec } A_\alpha \rightarrow \mathbf{M}$. We first recall the definition of a perfect obstruction theory of $\mathbf{M}$.

Definition 2.1. *A perfect obstruction theory of $\mathbf{M}$ (over the atlas Λ) consists of a choice of perfect obstruction theory $(E_\alpha^\bullet, \mathfrak{ob}_\alpha)$ of S_α for each $\alpha \in \Lambda$ so that they satisfy the following compatibility condition: First, let $\mathcal{O}b_\alpha$ be the obstruction sheaf (i.e. $= \text{Coker}\{E_\alpha^1 \rightarrow E_\alpha^2\}$), then the collection $\{\mathcal{O}b_\alpha\}_{\alpha \in \Lambda}$ descends to a (global) sheaf of $\mathcal{O}_{\mathbf{M}}$ -modules $\mathcal{O}b_{\mathbf{M}}$. Secondly, the obstruction assignments $\mathfrak{ob}_\alpha$ and $\mathfrak{ob}_\beta$ are identical when pulled back to $S_{\alpha\beta}$, using the given isomorphisms.*

Note that $\{\mathcal{O}b_\alpha\}$ descends means that over $S_{\alpha\beta}$ the pull back of $\mathcal{O}b_\alpha$ and of $\mathcal{O}b_\beta$ are isomorphic and that such isomorphisms satisfy the cocycle condition on $S_{\alpha\beta\gamma}$.

Our immediate goal is to show that the obstruction theories of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ naturally induces a perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$.

Theorem 2.2. *There is a natural perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ induced by the perfect obstruction theories of $\mathfrak{M}(W[n], \Gamma)^{st}$ constructed in the previous section.*

Proof. Let S be an affine chart of $\mathfrak{M}(\mathfrak{W}, \Gamma)$. Without loss of generality, we can assume S is one of the chart constructed in the proof of Theorem 3.10 in [Li]. Namely, there is a chart $\bar{S} \subset \mathfrak{M}(W[n], \Gamma)^{st}$ for some n so that S is a closed subscheme of $\bar{S}$ and $S \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ is induced by $S \rightarrow \bar{S}$ and $\mathfrak{M}(W[n], \Gamma)^{st} \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$. We let $S = \text{Spec } A$ and $\bar{S} = \text{Spec } \bar{A}$. We then let $\bar{E}^\bullet = [\bar{E}^1 \rightarrow \bar{E}^2]$ be the complex of $\bar{A}$ -modules provided by Theorem 1.20 for the chart $\bar{S}$.

We begin with the functor of the first order deformations in S . As argued in the proof of [Li, Theorem 3.10], there is a neighborhood U of $S \times \{e\} \subset S \times G[n]$ so that the morphism $S \times G[n] \rightarrow \mathfrak{M}(W[n], \Gamma)^{st}$ induced by the $G[n]$ -action lifts to an étale $\phi: U \rightarrow \bar{S}$. Since $U \rightarrow \bar{S}$ is étale, each vector $v \in T_e G[n]$ defines a first order deformation of the inclusion $S \rightarrow \bar{S}$, and hence an element $\bar{v} \in h^1(\bar{E}^\bullet \otimes_{\bar{A}} A)$. This induces a homomorphism

$$(2.1) \quad T_e G[n] \otimes_{\mathbb{K}} A \longrightarrow h^1(\bar{E}^\bullet \otimes_{\bar{A}} A) \longrightarrow \bar{E}^1 \otimes_{\bar{A}} A.$$

Since elements in $\mathfrak{M}(W[n], \Gamma)^{st}$ associate to stable morphisms to $\mathfrak{W}$, at each closed point $p \in S$ the homomorphism $T_e G[n] \rightarrow T_p \bar{S}/T_p S$ induced by the group action is injective. Hence the cokernel of (2.1) is also free. Now let E^1 be the cokernel of (2.1) and let $E^2 = \bar{E}^2 \otimes_{\bar{A}} A$. Let $L \subset \bar{E}^1 \otimes_{\bar{A}} A$ be the image module of (2.1). Since L lies in the image of the second arrow (above), the image of L in E^2 is trivial. Hence $\bar{E}^1 \otimes_{\bar{A}} A \rightarrow \bar{E}^2 \otimes_{\bar{A}} A$ lifts to $E^1 \rightarrow E^2$.

We next show that the natural obstruction theory of S takes values in the cohomology theory of $E^\bullet$. First, since (2.1) has free cokernel, $E^\bullet$ is a two-term complex of finitely generated free A -modules. Secondly, that the functor of the first order deformations in S is given by the functor $h^1(E^\bullet)$ is obvious since the morphism

$$(2.2) \quad S \times G[n] \supset U \xrightarrow{\phi} \bar{S}$$

induced by the group action is étale near $S \times \{e\}$. Finally, let $(B, I, \varphi) \in \text{Ob}(\mathfrak{Tri}_S)$, then it is also an object in $\mathfrak{Tri}_{\bar{S}}$ and thus has an obstruction class

$$\text{ob}(B, I, \varphi) \in h^2(\bar{E}^\bullet \otimes_{\bar{A}} I) = h^2(E^\bullet \otimes_A I)$$

to extending $\varphi: \text{Spec } B/I \rightarrow S$ to $\text{Spec } B \rightarrow \bar{S}$. Because $G[n]$ is smooth and $\phi: U \rightarrow \bar{S}$ (in (2.2)) is étale, $\text{ob}((B, I, \varphi))$ is also an obstruction class to extending φ to $\text{Spec } B \rightarrow S$.

Now let S_α be charts of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ with $E_\alpha^\bullet$ their complexes that are part of their obstruction theories. Then it is direct to check that the collection $\{h^2(E_\alpha^\bullet)\}$ form a sheaf over $\mathfrak{M}(\mathfrak{W}, \Gamma)$, and the obstruction assignments ob_α are compatible. This completes the proof of the Theorem. $\square$

We now state the theorem concerning the obstruction theory of $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$.

Theorem 2.3. *The perfect obstruction theory of $\mathfrak{M}(Z[n]^{\text{rel}}, \Gamma)^{st}$ constructed in the previous section naturally induces a perfect obstruction theory of $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$.*

Proof. We will omit the proof here since it is exactly the same as the proof of the previous Theorem. $\square$

The next issue is about the obstruction theory of the substack $\mathfrak{M}(\mathfrak{W}_t, \Gamma)$ defined by the fiber product

$$\mathfrak{M}(\mathfrak{W}_t, \Gamma) = \mathfrak{M}(\mathfrak{W}, \Gamma) \times_C t,$$

where $t \in C$ is a closed point. Clearly, when $t \neq 0$ the stack $\mathfrak{M}(\mathfrak{W}_t, \Gamma)$ is naturally isomorphic to the module stack of stable morphisms to W_t of topological type Γ , which itself admits a natural obstruction theory as worked out in [LT1]. The obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ deserves more attention since it was not known before.

We now study the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. Let $W_0[n] = W[n] \times_C 0$. As in the case of $\mathfrak{M}(\mathfrak{W}, \Gamma)$, we only need to work out the obstruction theory of

$$\mathfrak{M}(W_0[n], \Gamma)^{\text{st}} \triangleq \mathfrak{M}(W[n], \Gamma)^{\text{st}} \times_C 0.$$

Let $S = \text{Spec } A$ be an affine chart of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$. Then $S_0 = S \times_C 0$ is an affine chart of $\mathfrak{M}(W_0[n], \Gamma)^{\text{st}}$. We let A_0 be the quotient ring of A so that $S_0 = \text{Spec } A_0$. As before we let $f: \mathcal{X} \rightarrow W[n]$ be the universal family over S and let $f_0: \mathcal{X}_0 \rightarrow W_0[n]$ be the restriction of f to $S_0 \subset S$. We fix a sufficiently fine covering $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha, \mathcal{W}_\alpha)$ of f indexed by Λ and let $\mathbf{E}^\bullet$ be the associated complex of A -modules constructed in section 1.1. We let $\underline{\mathcal{V}}_0$ be the coverings $\{\mathcal{V}_{0,\alpha}\}$ of S_0 with $\mathcal{V}_{0,\alpha} = \mathcal{V}_\alpha \times_C 0$. Similarly we let $\mathcal{U}_{0,\alpha} = \mathcal{U}_\alpha \times_C 0$ and let $f_{0,\alpha}: \mathcal{U}_{0,\alpha} \rightarrow W_0[n]$ be the restriction of f to $\mathcal{U}_{0,\alpha}$. For any A_0 -module I , we define

$$(2.3) \quad \Gamma_{\mathcal{V}_{0,\alpha_1 \dots \alpha_m}}(I) \triangleq I \otimes_{A_0} \Gamma(\mathcal{O}_{\mathcal{V}_{0,\alpha_1 \dots \alpha_m}}).$$

Using these, we can form a Čech complex $\mathbf{C}^\bullet(\underline{\mathcal{V}}_0, A_0)$ with the standard coboundary operation. Let $\mathbf{E}_0^\bullet = \mathbf{E}^\bullet \otimes_A A_0$. We next construct a homomorphism

$$\bar{\delta}_i: \mathbf{E}_0^i \rightarrow \mathbf{C}^{i-1}(\underline{\mathcal{V}}_0, A_0) \otimes_{\mathbb{k}} T_0 C.$$

Here we understand $\mathbf{C}^{-1}(\underline{\mathcal{V}}_0, A_0) = 0$. Let $\xi \in \mathbf{E}_0^1$ be any element. We write $\xi = (a, b)$ with $a \in F^1 \otimes_A A_0$ and $b \in \mathbf{C}^0(\Lambda, \text{Hom}(f^* \Omega_{W[n]}, A_0)^\dagger)$, as in section 1. Then by the construction in section 1.1, to each $\alpha \in \Lambda$ the element a defines an extension of $\mathcal{U}_{0,\alpha}/\mathcal{V}_{0,\alpha}$ to $\tilde{\mathcal{U}}_{0,\alpha}/\tilde{\mathcal{V}}_{0,\alpha}$ by $A_{0,\alpha} (= \Gamma_{\mathcal{V}_{0,\alpha}}(A_0))$. The extension ζ_α chosen before (1.19) induces an extension $\tilde{\zeta}_{0,\alpha}: \tilde{\mathcal{U}}_{0,\alpha} \rightarrow W[n]$ of $f_{0,\alpha}: \mathcal{U}_{0,\alpha} \rightarrow W_0[n]$. We let $\pi_n: W[n] \rightarrow C$ be the tautological projection. We now consider $\pi_n \circ \tilde{\zeta}_{0,\alpha}: \tilde{\mathcal{U}}_{0,\alpha} \rightarrow C$. Since $\pi_n \circ \tilde{\zeta}_{0,\alpha}|_{\mathcal{U}_{0,\alpha}}$ factor through $0 \in C$,

$$\mathbf{d}(\pi_n \circ \tilde{\zeta}_{0,\alpha} - 0) \in \Gamma_{\mathcal{U}_{0,\alpha}}(\mathcal{A}_{0,\alpha}) \otimes_{\mathbb{k}} T_0 C, \quad \mathcal{A}_{0,\alpha} = A_0 \otimes_{A_0} \mathcal{O}_{\mathcal{U}_{0,\alpha}}.$$

Because $\tilde{\zeta}_{0,\alpha}$ is a pre-deformable extension, the above element lies in $A_{0,\alpha} \otimes_{\mathbb{k}} T_0 C$. We define $\bar{\delta}(a)_\alpha$ to be this element. As to $\bar{\delta}(b)_\alpha$, since $b_\alpha \in \Gamma(\mathcal{U}_\alpha, \text{Hom}(f^* \Omega_{W[n]}, A_0)^\dagger)$, b_α induces a homomorphism $(\pi_n \circ f)^* T_0^\vee C \rightarrow \mathcal{A}_{0,\alpha}$. Again this is an element in $A_{0,\alpha} \otimes_{\mathbb{k}} T_0 C$. We define $\bar{\delta}_1(b)_\alpha$ to be this element. Clearly, this construction carries over to the case of multi-indices. This defines a map (a homomorphism) of complexes

$$\bar{\delta}: \mathbf{E}_0^\bullet \longrightarrow \mathbf{C}^{\bullet-1}(\underline{\mathcal{V}}_0, A_0) \otimes_{\mathbb{k}} T_0 C.$$

It is direct to check that this is a homomorphism of complexes. We let $\mathbf{F}^\bullet$ be the associated complex defined by $\mathbf{F}^i = \mathbf{E}_0^i \oplus \mathbf{C}^{i-2}(\underline{\mathcal{V}}_0, A_0) \otimes_{\mathbb{k}} T_0 C$ whose differential is the obvious induced one. Then we have a short exact sequence of complexes

$$0 \implies \mathbf{C}^{\bullet-1}(\underline{\mathcal{V}}_0, A_0) \otimes_{\mathbb{k}} T_0 C \implies \mathbf{F}^\bullet \implies \mathbf{E}^\bullet \otimes_A A_0 \implies 0$$

which induces a long exact sequence of cohomologies (for any A_0 -module I)

$$\begin{aligned} 0 \longrightarrow h^1(\mathbf{F}^\bullet \otimes_{A_0} I) &\longrightarrow h^1(\mathbf{E}_0^\bullet \otimes_{A_0} I) \longrightarrow I \otimes_{\mathbb{k}} T_0 C \longrightarrow \\ &\longrightarrow h^2(\mathbf{F}^\bullet \otimes_{A_0} I) \longrightarrow h^2(\mathbf{E}_0^\bullet \otimes_{A_0} I) \longrightarrow 0. \end{aligned}$$

Here we used the fact that since $\mathcal{V}_0$ is an étale covering of S_0 , $h^j(\mathbf{C}^\bullet(\mathcal{V}_0, I)) = I$ when $j = 0$ and vanishes when $j \geq 1$.

Since terms in $\mathbf{C}^\bullet(\mathcal{V}_0, A_0)$ are flat A_0 -modules, we can pick a complex of finitely generated free A_0 -modules $F^\bullet = [F^1 \rightarrow F^2]$ so that $F^\bullet$ is quasi-isomorphic to $\mathbf{F}^\bullet$.

Proposition 2.4. *The chart S_0 admits a natural perfect obstruction theory taking values in the cohomology of the complex $F^\bullet$.*

Proof. We need to check that the functor of the first order deformations $\mathfrak{Def}_{S_0}^1$ is isomorphic to the functor $\mathfrak{h}^1(F^\bullet)$ and that there is an obstruction assignment taking values in $h^2(F^\bullet)$ that satisfies the required base change property.

The fact that the functor $\mathfrak{Def}_{S_0}^1$ is isomorphic to $\mathfrak{h}^1(F^\bullet) \equiv \mathfrak{h}^1(\mathbf{F}^\bullet)$ follows directly from the definition and will be omitted. Now we construct the obstruction assignment. Let (B, I, φ) be any object in $\mathfrak{Tri}_{S_0}$. Let $T = \text{Spec } B/I$ and $f_T: \mathcal{X}_T \rightarrow W_0[n]$ be the pull back family under $\varphi: T \rightarrow S_0$. Let $\mathcal{U}_{T,\alpha}/\mathcal{V}_{T,\alpha}$ be the pull back of $\mathcal{U}_{0,\alpha}/\mathcal{V}_{0,\alpha}$ and let $f_{T,\alpha}: \mathcal{U}_{T,\alpha} \rightarrow W_0[n]$ be the restriction of f_T to $\mathcal{U}_{T,\alpha}$. Recall that in constructing the obstruction class to extending $T \rightarrow S_0$ to $\tilde{T} = \text{Spec } B \rightarrow S$, we first extend $\mathcal{X}_T/T$ to $\mathcal{X}_{\tilde{T}}/\tilde{T}$ and extend $f_{T,\alpha}: \mathcal{U}_{T,\alpha} \rightarrow W_0[n]$ to $f_{\tilde{T},\alpha}: \mathcal{U}_{\tilde{T},\alpha} \rightarrow W[n]$ where $\mathcal{U}_{\tilde{T},\alpha}/\mathcal{V}_{\tilde{T},\alpha}$ is the minimal extension of $\mathcal{U}_{T,\alpha}/\mathcal{V}_{T,\alpha}$ in $\mathcal{X}_{\tilde{T}}/\tilde{T}$. We then use the difference of $f_{\tilde{T},\alpha}$ and $f_{\tilde{T},\beta}$ to build a cocycle $a \in \mathbf{E} \otimes_{A_0} I$. Let $\pi_n: W[n] \rightarrow C$ be the tautological projection as before. Since $\pi_n \circ f_{T,\alpha}$ factor through $0 \in C$, $\pi_n \circ f_{\tilde{T},\alpha} \in \mathcal{I}_\alpha$. Further because $f_{\tilde{T},\alpha}$ is a pre-deformable extension, it lies in $\mathcal{I}_\alpha$. Hence the collection $\{\pi_n \circ f_{\tilde{T},\alpha}\}$ defines a cochain $c \in \mathbf{C}^0(\mathcal{V}_0, I)$. It is routine to check that the pair $(a, c) \in \mathbf{F}^2 \otimes_{A_0} I$ is closed, and hence defines a cohomology class $[(a, c)] \in h^2(\mathbf{F}^\bullet \otimes_{A_0} I)$. Further, it is routine to check that this class is independent of the choice of the extensions $f_{\tilde{T},\alpha}$, and that it is an obstruction class to extending $\varphi: T \rightarrow S_0$ to $\tilde{T} \rightarrow S_0$. We define $\mathfrak{ob}_0$ be the assignment that assigns $(B, I, \varphi) \in \text{Ob}(\mathfrak{Tri}_{S_0})$ to this class in $h^2(\mathbf{F}^\bullet \otimes_{A_0} I) = h^2(F^\bullet \otimes_{A_0} I)$. $\square$

Theorem 2.5. *The obstruction theories of the charts S_0 so defined induce a perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$.*

Proof. The only thing needs to be checked is that the sheaves $h^2(\mathbf{F}^\bullet)$ over the charts S_0 of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ just constructed descends to a global sheaf over $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ and that the obstruction assignments over these charts are mutually compatible. This is routine and will be omitted. $\square$

We comment that so far all the results concerning $\mathfrak{W}_0$, including its construction, are based on the existence of the smoothing W of W_0 . It is not difficult to see that we can construct $W_0[n]$ from W_0 directly, without reference to W . Therefore, we can define $\mathfrak{W}_0$ and the moduli stack $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ directly without assuming the existence of W . The construction of the perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ without using W is a little tricky, but should be doable. Since we will not use this in this paper, we will content with assuming the existence of a smoothing W of W_0 .

2.2. Gromov-Witten invariants. The goal of this subsection is to construct the virtual moduli cycles of the moduli stacks $\mathfrak{M}(\mathfrak{W}, \Gamma)$, $\mathfrak{M}(\mathfrak{W}_t, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ and to define their respective Gromov-Witten invariants.

Currently, there are two constructions of virtual moduli cycles in algebraic geometry. One is the original construction by Tian and the author. They assumed

that the moduli space admits a perfect obstruction theory. They then constructed a global cone that function as a virtual normal cone. Such cone was constructed using the (algebraic) Kuranishi maps of the obstruction theory [LT1, LT2]. In their construction they made a technical assumption that there is a global vector bundle on the moduli space that makes the obstruction sheaf its quotient. With this vector bundle, the cone becomes a subcone of this vector bundle, and the virtual moduli cycle is the intersection of this cone with the zero section of this vector bundle, using Gysin map. The alternative construction of Behrend and Fantechi [Beh, BF] works along a parallel line. They constructed a similar cone as an Artin stack, assuming the moduli space admits a perfect obstruction theory. They then obtain a cone cycle in a vector bundle by assuming the existence of a global vector bundle, as in the original construction of Tian and the author. These two constructions yield identical cycles [KKP]. Recently, by working out the intersection theory on Artin stacks, Kresh [Kr2] showed that one can construct the virtual moduli cycle without relying on the existence of a global vector bundle as mentioned, thus removing this technical condition. This makes the construction of virtual moduli cycles more versatile. After seeing Kresh's work, we realized that by applying a simple trick we can remove the technical condition of the existence of such vector bundles in our construction of the virtual moduli cycles. In the following, we will present this modified construction.

We begin with the general situations. Let $\mathbf{M}$ be a proper and separated DM-stack. It follows from the definition that $\mathbf{M}$ can be covered by a quasi-projective scheme. We let $\mathcal{O}b$ be a sheaf of $\mathcal{O}_{\mathbf{M}}$ -modules. We assume that there is a finite collection of schemes S_α and smooth morphisms $\rho_\alpha : S_\alpha \rightarrow \mathbf{M}$, indexed by a set Λ , so that the collection of images $\rho_\alpha(S_\alpha) \subset \mathbf{M}$ form an open covering of $\mathbf{M}$. We next assume that to each $\alpha \in \Lambda$ there is a locally free sheaf of $\mathcal{O}_{S_\alpha}$ -modules $\mathcal{E}_\alpha$, a surjective homomorphism of sheaves $\mathcal{E}_\alpha \rightarrow \rho_\alpha^* \mathcal{O}b$ and a cone cycle $C_\alpha \in Z_* \text{Vect}(\mathcal{E}_\alpha)$ that satisfy the following *cycle consistency criteria*. Here we denote by $\text{Vect}(\mathcal{E}_\alpha)$ the vector bundle over S_α so that its sheaf of sections is $\mathcal{E}_\alpha$. In this paper, by abuse of notation we will view a vector bundle as its total space. We first fix a few notations before we state the criteria. Let $p \in \mathbf{M}$ be any closed point. We pick an étale morphism $\varphi : (X, \bar{p}) \rightarrow (\mathbf{M}, p)$ and let $\hat{X}_p$ be the formal completion of X along $\bar{p}$. We let G_p be the automorphism group of $p \in \mathbf{M}$. Note that G_p acts naturally on $\hat{X}_p$ and up to G_p the scheme $\hat{X}_p$ is canonical. We next let $V_{\bar{p}} = \varphi^* \mathcal{O}b \otimes_{\mathcal{O}_X} \mathbb{K}_{\bar{p}}$ and let $V_{\hat{X}_p}$ be the vector bundle $V_{\bar{p}} \times \hat{X}_p$ over $\hat{X}_p$. Again G_p acts on V_p and $V_{\hat{X}_p}$, and up to G_p they are canonical.

Cycle consistency criteria. *We say the collection $\mathcal{C} = \{(S_\alpha, \mathcal{E}_\alpha, C_\alpha)\}_\Lambda$ satisfies the cycle consistency criteria at $p \in \mathbf{M}$ if there is a cycle $C_p \in Z_* V_{\hat{X}_p}$ invariant under G_p of which the following hold. Let $\alpha \in \Lambda$ be any index, let $S_{\alpha,p} = S_\alpha \times_{\mathbf{M}} p$ and let $\hat{S}_\alpha = S_\alpha \times_{\mathbf{M}} \hat{X}_p$. We let pr_i be the i -th projection of the product $\hat{S}_\alpha = S_\alpha \times_{\mathbf{M}} \hat{X}_p$. Then there is a surjective homomorphism of vector bundles*

$$\Phi_1 : \text{Vect}(\mathcal{E}_\alpha) \times_{S_\alpha} \hat{S}_\alpha \longrightarrow V_{\hat{X}_p} \times_{\hat{X}_p} \hat{S}_\alpha$$

extending the canonical (composite) homomorphism

$$\text{Vect}(\mathcal{E}_\alpha) \times_{S_\alpha} S_{\alpha,p} \longrightarrow \text{Vect}(\text{pr}_1^* \rho_\alpha^* \mathcal{O}b|_{S_{\alpha,p}}) \equiv \text{Vect}(\text{pr}_2^* \varphi^* \mathcal{O}b|_{S_{\alpha,p}}) \equiv V_{\bar{p}} \times S_{\alpha,p}^{10}$$

¹⁰ For sheaves $\mathcal{F}$ of $\mathcal{O}_Z$ -modules and closed subscheme $X \subset Z$ we use $\mathcal{F}|_X$ to denote $\mathcal{F} \otimes_{\mathcal{O}_Z} \mathcal{O}_X$.

so that $\Phi_1^* C_{\xi} = \Phi_2^* C_{\alpha}$. Here the first arrow above is induced by $\mathcal{E}_{\alpha} \rightarrow p_{\alpha}^* \mathcal{O}b$, Φ_2 is the tautological flat morphism $\mathrm{Vect}(\mathcal{E}_{\alpha}) \times_{S_{\alpha}} \hat{S}_{\alpha} \rightarrow \mathrm{Vect}(\mathcal{E}_{\alpha})$ and Φ_i^* are the flat pull back homomorphism of cycles.

We will call the collection $\hat{\mathcal{C}} \triangleq \{C_p \subset V_{\hat{X}_p}\}_{p \in \mathbf{M}}$ satisfying the above criteria the infinitesimal models of the collection $\mathcal{C}$. Accordingly we will call $\mathcal{C}$ a local model of $\hat{\mathcal{C}}$. In the following, we say the collection $\mathcal{C}$ is consistent if there is a $\hat{\mathcal{C}}$ as above that satisfy the above criteria. Conversely, given $\hat{\mathcal{C}}$, we say it can be algebraized if there is a $\mathcal{C}$ so that they satisfy the above criteria. Note that once the infinitesimal models exist, then the property of the local model $\mathcal{C}$ is completely determined by the infinitesimal models. This is the key to many of the results concerning virtual moduli cycles.

Given the collection $\mathcal{C} = \{(S_{\alpha}, \mathcal{E}_{\alpha}, C_{\alpha})\}_{\alpha}$ over $(\mathbf{M}, \mathcal{O}b)$ we now construct a canonical cycle $[\mathcal{C}] \in A_* \mathbf{M}$ as follows. For each α we let Ξ_{α} be the set of irreducible components of C_{α} . For $a \in \Xi_{\alpha}$ we denote by N_a the irreducible variety (component) in C_{α} associated to a and let m_a be the multiplicity of N_a in C_{α} . Then we have

$$(2.4) \quad C_{\alpha} = \sum_{a \in \Xi_{\alpha}} m_a N_a \in Z_* \mathrm{Vect}(\mathcal{E}_{\alpha}).$$

For any $a \in \Xi_{\alpha}$ we define the base stack of a to be the minimal closed integral substack $\mathbf{Y}_a \subset \mathbf{M}$ so that the natural $N_a \rightarrow \mathbf{M}$ factor through $\mathbf{Y}_a \subset \mathbf{M}$. We let $j_a : \mathbf{Y}_a^0 \rightarrow \mathbf{Y}_a$ be the (maximal) dense open substack so that the pull back sheaf $j_a^* \mathcal{O}b$ is locally free. Then $\mathbf{F}_a^0 \triangleq \mathrm{Vect}(j_a^* \mathcal{O}b)$ is a vector bundle stack over $\mathbf{Y}_a^0$. Further, the natural morphism

$$(2.5) \quad \eta_a : \mathrm{Vect}(\mathcal{E}_{\alpha})|_{\rho_{\alpha}^{-1}(\mathbf{Y}_a^0)} \longrightarrow \mathbf{F}_a^0 \equiv \mathrm{Vect}(j_a^* \mathcal{O}b)$$

induced by $\mathcal{E}_{\alpha} \rightarrow \rho_{\alpha}^* \mathcal{O}b$ is flat. We let $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ be the image stack of $N_a|_{\rho_{\alpha}^{-1}(\mathbf{Y}_a^0)}$ under η_a with the reduced stack structure. Clearly, by cycle consistency criteria the flat pull back $\eta_a^* \mathbf{N}_a^0$ contains $N_a|_{\rho_{\alpha}^{-1}(\mathbf{Y}_a^0)}$ as one of its irreducible components. In the following, we will call $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ the intrinsic representative of a . Note that since we choose $\mathbf{Y}_a^0$ to be the maximal possible open substack of $\mathbf{Y}_a$ so that $j_a^* \mathcal{O}b$ is locally free, the open $\mathbf{Y}_a^0 \subset \mathbf{Y}_a$ and the representative $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ only depend on a .

Now let $b \in \Xi_{\beta}$ be any element with $\mathbf{Y}_b$ and $\mathbf{N}_b^0 \subset \mathbf{F}_b^0$ its base stack and intrinsic representative. We say $a \sim b$ if $\mathbf{Y}_a = \mathbf{Y}_b$ and $\mathbf{N}_a^0 = \mathbf{N}_b^0$ in $\mathbf{F}_a^0 \equiv \mathbf{F}_b^0$. This defines an equivalence relation $\sim$ on $\cup_{\alpha \in \Lambda} \Xi_{\alpha}$. We define $\Xi = (\cup_{\alpha \in \Lambda} \Xi_{\alpha}) / \sim$. Again by the cycle consistency criteria whenever $a \sim b$ then $m_a = m_b$. Hence each $a \in \Xi$ has an associated multiplicity m_a , a base substack $\mathbf{Y}_a$ and an intrinsic representative $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ over an open substack $\mathbf{Y}_a^0 \subset \mathbf{Y}_a$.

Assuming there is a global locally free sheaf $\mathcal{E}$ on $\mathbf{M}$ making $\mathcal{O}b$ its quotient sheaf, then over each $\mathbf{Y}_a^0$ we have a flat projection $\mathrm{Vect}(\mathcal{E})|_{\mathbf{Y}_a^0} \rightarrow \mathbf{F}_a^0$. We let $\mathbf{N}_a \subset \mathrm{Vect}(\mathcal{E})$ be the closure of the pull back of $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ under this projection. The associated cycle $[\mathcal{C}]$ is then defined to be

$$(2.6) \quad [\mathcal{C}] = \sum_{a \in \Xi} m_a 0_{\mathrm{Vect}(\mathcal{E})}^! [\mathbf{N}_a] \in A_* \mathbf{M},$$

where $0_{\mathrm{Vect}(\mathcal{E})}^!$ is the Gysin map of the 0-section of $\mathrm{Vect}(\mathcal{E})$. This is essentially the original construction of Tian and the author.

We now back to the general situation (without assuming the existence of such $\mathcal{E}$). We need to define a map $\xi: \Xi \rightarrow A_*\mathbf{M}$ so that $\xi(a)$ is the cycle $0_{\text{Vect}(\mathcal{E})}^1[\mathbf{N}_a]$ should a global $\mathcal{E}$ exist. Let $a \in \Xi$ be any element. Since $\mathbf{M}$ can be covered by a quasi-projective scheme, there is a normal projective variety Y_a and a generically finite surjective morphism $\varphi_a: Y_a \rightarrow \mathbf{Y}_a$. By abuse of notation, we also view φ_a as the composite of $Y_a \rightarrow \mathbf{Y}_a$ with $\mathbf{Y}_a \rightarrow \mathbf{M}$. Since Y_a is projective, there is a locally free sheaf of $\mathcal{O}_{Y_a}$ -modules $\mathcal{F}_a$ so that $\varphi_a^*\mathcal{O}b$ is a quotient sheaf of $\mathcal{F}_a$. We denote by F_a the vector bundle $\text{Vect}(\mathcal{F}_a)$ over Y_a . Let $Y_a^0 \subset Y_a$ be a dense open subset so that $Y_a^0 \rightarrow \mathbf{Y}_a^0$ is étale. Then the morphism $F_a|_{Y_a^0} \rightarrow \mathbf{F}_a^0$ induced by $\mathcal{F}_a \rightarrow \varphi_a^*\mathcal{O}b$ is a flat morphism. We let N_a be the closure in F_a of the flat pull-back of $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$. Note that N_a only depends on Y_a and $\mathcal{F}_a \rightarrow \varphi_a^*\mathcal{O}b$. The cycle $N_a \subset F_a$ will be called a representative of $a \in \Xi$. With $N_a \subset F_a$ chosen, we define

$$(2.7) \quad \xi(a) = \deg(\varphi_a)^{-1} \varphi_{a*} 0_{F_a}^1[N_a],$$

where 0_{F_a} is the zero section of F_a , $0_{F_a}^1$ is the Gysin homomorphism $Z_*F_a \rightarrow A_*Y_a$ (of the zero section of 0_{F_a}) and φ_{a*} is the push-forward homomorphism of cycles. The degree $\deg(\varphi_a)$ is the degree of the morphism $\varphi_a: Y_a \rightarrow \mathbf{Y}_a$ defined in [Vis]. Finally, we define

$$(2.8) \quad [\mathcal{C}] = \sum_{a \in \Xi} m_a \xi(a) \in A_*\mathbf{M}.$$

Note that this construction coincides with that in (2.6) in case a global locally free sheaf $\mathcal{E}$ exists.

We will call this construction the basic construction and call $[\mathcal{C}]$ the associated cycle of the collection $\mathcal{C}$.

Lemma 2.6. *Let the notation be as before. Then $\xi(a)$ is independent of the choice of Y_a and $\mathcal{F}_a$.*

Proof. Let $\varphi_{a,1}: Y_{a,1} \rightarrow \mathbf{Y}_a$ and $\varphi_{a,2}: Y_{a,2} \rightarrow \mathbf{Y}_a$ be two normal varieties and generically finite dominant morphisms and let $N_{a,1} \subset F_{a,1}$ and $N_{a,2} \subset F_{a,2}$ be the respective choices of the representatives of a over $Y_{a,1}$ and $Y_{a,2}$. To prove the Lemma it suffices to show that

$$\deg(\varphi_{a,1})^{-1} \varphi_{a,1*} 0_{F_{a,1}}^1[N_{a,1}] = \deg(\varphi_{a,2})^{-1} \varphi_{a,2*} 0_{F_{a,2}}^1[N_{a,2}].$$

We let Y_a be the normalization of an irreducible component of $Y_{a,1} \times_{\mathbf{Y}_a} Y_{a,2}$ that is dominant and generically finite over $\mathbf{Y}_a$. We let $p_i: Y_a \rightarrow Y_{a,i}$ be the projection induced by the i -th projection of $Y_{a,1} \times_{\mathbf{Y}_a} Y_{a,2}$. We pick a locally free sheaf $\mathcal{F}_a$ on Y_a and surjective homomorphism $\mathcal{F}_a \rightarrow p_i^* \mathcal{F}_{a,i}$ so that the diagram

$$\begin{array}{ccc} \mathcal{F}_a & \longrightarrow & p_1^* \mathcal{F}_{a,1} \\ \downarrow & & \downarrow \\ p_2^* \mathcal{F}_{a,2} & \longrightarrow & p_1^* \varphi_{a,1}^* \mathcal{O}b \equiv p_2^* \varphi_{a,2}^* \mathcal{O}b \end{array}$$

is commutative. Now let $F_a = \text{Vect}(\mathcal{F}_a)$, let $U \subset Y_a$ be a dense open subset so that the projections $U \rightarrow \mathbf{Y}_a$, $U \rightarrow Y_{a,1}$ and $U \rightarrow Y_{a,2}$ are flat. Then the flat pull-back of $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$ via the induced $F_a|_U \rightarrow \mathbf{F}_a^0$ is identical to the flat pull back of $N_{a,i} \subset F_{a,i}$ under the flat morphism $F_a|_U \rightarrow F_{a,i}$. Further, it is direct to check that $\deg(\varphi_a) = \deg(\varphi_{a,i}) \deg(p_i)$. Hence if we let $\tilde{N}_{a,i} \subset F_{a,i} \times_{Y_{a,i}} Y_a \triangleq p_i^* F_{a,i}$ be

the closure of the flat pull back of $N_{a,i}|_{U_i}$ (under $F_{a,i} \times_{Y_{a,i}} U \rightarrow F_{a,i}$), then for $i = 1$ and 2 ,

$$\varphi_{a,i*} 0_{F_{a,i}}^! [N_{a,i}] = \deg(p_i)^{-1} \varphi_{a*} 0_{p_i^* F_{a,i}}^! [\bar{N}_{a,i}] = \deg(p_i)^{-1} \varphi_{a*} 0_{F_a}^! [N_a].$$

This, combined with the identity about the degrees, proves the Lemma. $\square$

We now show how to apply this construction to construct the virtual moduli cycle of a Deligne-Mumford stack $\mathbf{M}$ endowed with a perfect obstruction theory. Let $\{S_\alpha\}_{\alpha \in \Lambda}$ be an atlas of $\mathbf{M}$ and let $\{(\mathcal{E}_\alpha^\bullet, \mathbf{ob}_\alpha)\}_\Lambda$ be the data associated to the perfect obstruction theory of $\mathbf{M}$ as in Definition 2.1. Here $\mathcal{E}_\alpha^\bullet = [\mathcal{E}_\alpha^1 \rightarrow \mathcal{E}_\alpha^2]$ is a complex of finite rank locally free sheaves of $\mathcal{O}_{S_\alpha}$ -modules. We let $\mathcal{O}_\mathbf{M}$ be the sheaf of $\mathcal{O}_\mathbf{M}$ -modules that is the descent of $\text{Coker}\{\mathcal{E}_\alpha^1 \rightarrow \mathcal{E}_\alpha^2\}$. Following [LT2, Section 3], to each α we can construct a canonical cone cycle $C_\alpha \subset \text{Vect}(\mathcal{E}_\alpha^2)$, using the relative Kuranishi-maps constructed from the perfect obstruction theory of S_α . We recall the main results in [LT2, Section 3]. Let $p \in \mathbf{M}$ be any closed point and let $\bar{p} \in S_\alpha$ be a lift of p for some $\alpha \in \Lambda$. We let $T_p = h^1(\mathcal{E}_\alpha^\bullet \otimes_{\mathcal{O}_{S_\alpha}} \mathbb{k}_{\bar{p}})$ and $O_p = h^2(\mathcal{E}_\alpha^\bullet \otimes_{\mathcal{O}_{S_\alpha}} \mathbb{k}_{\bar{p}})$. Then T_p is the Zariski tangent space of S_α at $\bar{p}$ and the obstruction theory $\mathbf{ob}_\alpha$ induces an obstruction theory to deformation of $\bar{p}$ in S_α taking values in O_p . Note that G_p naturally acts on T_p and O_p , and both T_p and O_p are canonical (depending only on p) up to the symmetry G_p . Let $X_p = \text{Spec } \mathbb{k}[[T_p^\vee]]$ and let $f_{\bar{p}} \in \mathbb{k}[[T_p^\vee]] \otimes_{\mathbb{k}} O_p$ be a Kuranishi map of the obstruction theory to deforming $\bar{p}$ in S_α . We let $\hat{X}_p = \text{Spec } \mathbb{k}[[T_p^\vee]]/(f_{\bar{p}})$. Then the formal completion of S_α along $\bar{p}$, denoted by $\hat{S}_\alpha$, is canonically isomorphic to $\hat{X}_p$. As before, we let $V_{\hat{X}_p}$ be the total scheme of the vector bundle $O_p \times \hat{X}_p$ over $\hat{X}_p$. Following the recipe in constructing the virtual moduli cycle, we consider the normal cone C_p to $\hat{X}_p$ in X_p . The cone C_p is naturally a subcone of $V_{\hat{X}_p}$. Though the Kuranishi map $f_{\bar{p}}$ is not unique and also depends on the choice of the lifting $\bar{p} \in S_\alpha$, the cone C_p is unique in the following sense.

Lemma 2.7. *Let the notation be as before. Let $0 \in \hat{X}_p$ be the unique closed point. Then the subscheme $C_p \times_{\hat{X}_p} 0 \subset V_{\hat{X}_p} \times_{\hat{X}_p} 0$ is G_p -invariant and only depends on $p \in \mathbf{M}$. In particular, the supports and the multiplicities of the irreducible components of C_p are independent of the choice of $\bar{p}$ and $f_{\bar{p}}$.*

The collection $\{C_p \subset V_{\hat{X}_p}\}_{p \in \mathbf{M}}$ form an infinitesimal models. By constructing relative Kuranishi family, Tian and the author established the following existence result [LT2, Section 3].

Lemma 2.8. *Let $\mathcal{F}$ be a locally free sheaf of $\mathcal{O}_{S_\alpha}$ -modules and $\eta: \mathcal{F} \rightarrow \mathcal{O}_{b_\alpha}$ be a surjective sheaf homomorphism. Then there is a unique cycle $C_{\mathcal{F}} \in Z_* \text{Vect}(\mathcal{F})$ that satisfies the cycle consistency criteria for the pair $(\mathbf{M}, \mathcal{O}_\mathbf{M})$ and the infinitesimal models $\{C_p \subset V_{\hat{X}_p}\}_{p \in \mathbf{M}}$.*

Applying Lemma 2.8 to $\rho_\alpha: S_\alpha \rightarrow \mathbf{M}$ and the quotient homomorphism $\mathcal{E}_\alpha^2 \rightarrow \rho_\alpha^* \mathcal{O}_\mathbf{M}$ we obtain a cycle $C_\alpha \in Z_* \text{Vect}(\mathcal{E}_\alpha^2)$. Since the cycles C_p are unique, Lemma 2.7 and 2.8 combined shows that the collection $\{(S_\alpha, \mathcal{E}_\alpha^2, C_\alpha)\}_\Lambda$ satisfies the cycle consistency criteria for $(\mathbf{M}, \mathcal{O}_\mathbf{M})$. We define the virtual moduli cycle of $\mathbf{M}$ (with the given perfect obstruction theory) to be the associated cycle of this collection, and denoted by $[\mathbf{M}]^{\text{virt}}$.

In the end, by applying this construction to the stacks $\mathfrak{M}(\mathfrak{W}, \Gamma)$, $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ we obtain moduli cycles $[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}$, $[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}}$ and $[\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)]^{\text{virt}}$. As in [LT2], for $t \in C$ we define the GW-invariant of W_t to be the homomorphism

$$\Psi_\Gamma^{W_t} : H^*(W_t)^{\times k} \times H^*(\mathfrak{M}_{g,k}) \longrightarrow H_0(\text{pt}) \cong \mathbb{Q}$$

defined by

$$\Psi_\Gamma^{W_t}(\alpha, \beta) = \mathbf{q}_{*0}(\text{ev}^*(\alpha) \cup \pi_{g,k}^*(\beta) [\mathfrak{M}(\mathfrak{W}_t, \Gamma)]^{\text{virt}}),$$

where $\pi_{g,k}$ and ev are the forgetful and evaluation morphisms, g and k are the genus and the number of marked points of the topological type Γ , $\mathbf{q} : \mathfrak{M}(\mathfrak{W}_t, \Gamma) \rightarrow \{t\} \subset C$ is the projection and $\mathbf{q}_{*0}$ is the push-forward $A_*\mathfrak{M}(\mathfrak{W}_t, \Gamma) \rightarrow H_0(pt)$ at degree 0. (Here we use H_* to denote the ordinary homology theory in case the ground field is $\mathbb{C}$. Otherwise one can use Chow rings to define the GW-invariants.)

The Gromov-Witten invariants of $\mathfrak{W}$ is the homomorphism

$$\Psi_\Gamma^{W/C} : H_C^0(R^*\pi_*\mathbb{Q}_W)^{\times k} \times H^*(\mathfrak{M}_{g,k}) \longrightarrow H_2^{\text{BM}}(C) \cong \mathbb{Q}$$

defined via a similar formula with $\mathbf{q}_{*0}$ replaced by $\mathbf{q}_{*1} : A_*\mathfrak{M}(\mathfrak{W}, \Gamma) \rightarrow H_2^{\text{BM}}(C)$. Here $\mathbb{Q}_W$ is the sheaf of locally constant sections on W taking values in $\mathbb{Q}$, $\pi : W \rightarrow C$ is the tautological projection and H_2^{BM} is the Borel-Moore homology of the open complex curve C .

Now we define the relative Gromov-Witten invariants of $Z^{\text{rel}} = (Z, D)$. Let Γ be an admissible weighted graph defined in [Li]. As mentioned before, it determines the topological type of relative morphisms to $\mathfrak{Z}^{\text{rel}}$. We let Γ^o be the sub-data (in Γ) relating to the domain curves. (Namely, the connected components, the genus and both kinds of marked points of the domain curves.) Let $\mathfrak{M}_{\Gamma^o}$ be the moduli space of stable curves with topological types Γ^o . Here a curve C of topological type Γ^o is stable if all connected components of C are stable pointed curves. Clearly, $\mathfrak{M}_{\Gamma^o}$ is a Deligne-Mumford stack. As in the ordinary case, there is a forgetful morphism $\pi_{\Gamma^o} : \mathfrak{M}(\mathfrak{Y}^{\text{rel}}, \Gamma) \longrightarrow \mathfrak{M}_{\Gamma^o}$. We define the relative GW-invariants to be the homomorphism

$$\Psi_\Gamma^{Z^{\text{rel}}} : H^*(Z)^{\times k} \times H^*(\mathfrak{M}_{\Gamma^o}) \longrightarrow H_*(D^r)$$

(recall k and r are the numbers of legs and roots of Γ) defined by

$$\Psi_\Gamma^{Z^{\text{rel}}}(\alpha, \beta) = \mathbf{q}_*(\text{ev}^*(\alpha) \cup \pi_{\Gamma^o}^*(\beta) [\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)]^{\text{virt}}) \in H_*(D^r).$$

Here

$$(2.9) \quad \mathbf{q} : \mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma) \rightarrow D^r \quad \text{and} \quad \text{ev} : \mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma) \longrightarrow Z^k$$

are the morphisms defined by evaluating on the distinguished and the ordinary marked sections respectively.

3. DEGENERATIONS OF GROMOV-WITTEN INVARIANTS

In this and the next section, we will prove the degeneration formula of the Gromov-Witten invariants of the family W/C stated in the introduction of this paper. We will state the first version of the degeneration formula in the first subsection. We will state the reduction Lemmas in subsection 6.2. The proof of these Lemmas will be postponed to the next section.

3.1. The first version of the degeneration formula. The first step to prove the degeneration formula is to express $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$ as a union of Cartier-divisors in $\mathfrak{M}(\mathfrak{W}, \Gamma)$. Here we fix a $\Gamma = (g, k, b)$ once and for all. We first define the notion of Cartier-divisor of an algebraic stack $\mathbf{M}$.

Definition 3.1. *Let $\mathbf{M}$ be an algebraic stack. A Cartier-divisor (in short C-divisor) on $\mathbf{M}$ is a pair $(\mathbf{L}, \mathbf{s})$ where $\mathbf{L}$ is a line bundle on $\mathbf{M}$ and $\mathbf{s}$ is a section of $\mathbf{L}$. An isomorphism between $(\mathbf{L}, \mathbf{s})$ and $(\mathbf{L}', \mathbf{s}')$ consists of an isomorphism $\mathbf{L} \cong \mathbf{L}'$ so that $\mathbf{s} \equiv \mathbf{s}'$ under this isomorphism.*

We comment that a C-divisor (L, s) over a scheme is a pseudo-divisor (defined in [Ful]) via (L, Z, s) where $Z = X - s^{-1}(0)$. Note that when (L, s) and (L', s') are two C-divisors, then $(L, s) \otimes (L', s') \triangleq (L \otimes L', ss')$ is also a C-divisor.

We now let Ω be the set of all admissible triples defined in [Li, Section 4] and reviewed in the introduction. Recall that $\eta = (\Gamma_1, \Gamma_2, I) \in \Omega$ is an admissible triple if Γ_1 and Γ_2 are two weighted graphs of identical numbers of roots and I is an order preserving inclusion $I: [k_1] \rightarrow [k]$ where k_i is the number of legs of Γ_i and $k = k_1 + k_2$. Let C_1 and C_2 be two curves of topological types Γ_1 and Γ_2 , respectively. Then we can identify the i -th distinguished marked point $q_{1,i} \in C_1$ with the i -th distinguished marked point $q_{2,i} \in C_2$ for all i to obtain a new curve $C \triangleq C_1 \sqcup C_2$. It has k marked points, ordered according to I . As part of the definition of Ω , we require that the multiplicities of the i -th roots of Γ_1 and Γ_2 are identical, that C is connected of genus g and $b = \sum_{x \in V(\Gamma_1) \cup V(\Gamma_2)} b(x)$ ($b(x)$ accounts for the degree of the stable morphism along the connected component labeled by x). We will call C the gluing of C_1 and C_2 along distinguished marked points. This gluing construction can be applied to a pair of families of curves. Hence for each $\eta = (\Gamma_1, \Gamma_2, I) \in \Omega$, we have a closed local immersion of stacks

$$(3.1) \quad \Phi_\eta : \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2) \longrightarrow \mathfrak{M}(\mathfrak{W}, \Gamma),$$

defined in [Li, (4.8)]. Here $\mathfrak{M}(\mathfrak{Y}_i^{\text{rel}}, \Gamma_i) \rightarrow D^r$ is the evaluation morphism $\mathbf{q}$ in (2.9) and the morphism (3.1) is defined by sending any pair $((f_1, \mathcal{X}_1), (f_2, \mathcal{X}_2))$ to the family $(f_1 \sqcup f_2, \mathcal{X}_1 \sqcup \mathcal{X}_2)$. Now let $\eta = (\Gamma_1, \Gamma_2, I) \in \Omega$. Following [Li], we define $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ be the image stack of (3.1). As was shown in [Li, Section 4],

$$(3.2) \quad \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \rightarrow \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$$

is finite, étale of pure degree $|\text{Eq}(\eta)|$. Here by abuse of notion we also use Φ_η to denote this induced morphism.

In this subsection, to each $\eta \in \Omega$ we will define a C-divisor $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ on $\mathfrak{M}(\mathfrak{W}, \Gamma)$ so that the vanishing locus (as topological space) of $\mathbf{s}_\eta$ is $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$. We begin with the study of line bundles on $\mathbf{A}^{n+1}$. We continue to use the convention introduced in [Li, Section 1] concerning subsets of $\mathbf{A}^{n+1}$. For $l \in [n+1]$ we still denote by $\mathbf{H}_l \subset \mathbf{A}^{n+1}$ the l -th coordinate hyperplane of $\mathbf{A}^{n+1}$. We define L_l be the line bundle on $\mathbf{A}^{n+1}$ so that its sheaf of sections is $\mathcal{O}_{\mathbf{A}^{n+1}}(\mathbf{H}_l)$ and define s_l be the section of L_l that is the constant section $1 \in \mathcal{O}_{\mathbf{A}^{n+1}}$ under the canonical inclusion $\mathcal{O}_{\mathbf{A}^{n+1}} \subset \mathcal{O}_{\mathbf{A}^{n+1}}(\mathbf{H}_l) \equiv \mathcal{O}_{\mathbf{A}^{n+1}}(L_l)$. Recall that $\mathbf{A}^{n+1}$ is a $G[n]$ -subscheme ($G[n] \triangleq GL(1)^{\times n}$) as defined in [Li, Section 1]. Since $\mathbf{H}_l \subset \mathbf{A}^{n+1}$ is invariant under the $G[n]$ -action, there is a unique $G[n]$ -linearization on L_l so that the section s_l is $G[n]$ -invariant. We fix such a linearization. Now let $J: [m+1] \rightarrow [n+1]$ be an order preserving embedding. Following the convention in [Li, Section 1], J

defines a standard embedding¹¹ $\gamma_J: \mathbf{A}^{m+1} \rightarrow \mathbf{A}^{n+1}$ and hence defines a pull back C-divisor $\gamma_J^*(L_l, s_l)$ on $\mathbf{A}^{m+1}$. There are two possibilities: One is when $l \neq \text{Im}(J)$. Then $\text{Im}(\gamma_J) \cap \mathbf{H}_l = \emptyset$ and hence there is a canonical isomorphism¹² $\gamma_J^*(L_l, s_l) \cong (\mathbf{1}_{\mathbf{A}^{m+1}}, 1)$. The other case is when $J(l') = l$ for some $l' \in [m+1]$, in which case we have $\gamma_J^*(L_l, s_l) \cong (L_{l'}, s_{l'})$.

Now let $J: [n_1 + 1] \rightarrow [n_2 + 1]$ be an order preserving embedding. Let S be any scheme, $\tau: S \rightarrow \mathbf{A}^{n_1+1}$ and $\rho: S \rightarrow G[n_2]$ be two morphisms with $\gamma_J: \mathbf{A}^{n_1+1} \rightarrow \mathbf{A}^{n_2+1}$ the standard embedding. As in [Li], we define $(\gamma_J \circ \tau)^\rho: S \rightarrow \mathbf{A}^{n_2+1}$ be the morphisms induced by $\gamma_J \circ \tau: S \rightarrow \mathbf{A}^{n_2+1}$ and the $G[n_2]$ -action on $\mathbf{A}^{n_2+1}$ via ρ .

Lemma 3.2. *Let J , τ and ρ be as before. In case $l_2 = J(l_1)$ then we have a natural isomorphism $((\gamma_J \circ \tau)^\rho)^*(L_{l_2}, s_{l_2}) = \tau^*(L_{l_1}, s_{l_1})$. In case $l_2 \notin \text{Im}(J)$ the same identity holds with (L_{l_1}, s_{l_1}) replaced by $(\mathbf{1}, 1)$.*

Proof. We have the canonical isomorphism $(\gamma_J \circ \tau)^*(L_{l_2}, s_{l_2}) \cong \tau^*(L_{l_1}, s_{l_1})$. The required isomorphism is then induced by the canonical isomorphism

$$((\gamma_J \circ \tau)^\rho)^*(L_{l_2}, s_{l_2}) \cong (\gamma_J \circ \tau)^*(L_{l_2}, s_{l_2})$$

induced by the $G[n_2]$ -linearization on (L_{l_2}, s_{l_2}) . This proves the Lemma. $\square$

We now construct the required C-divisor $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ on $\mathfrak{M}(\mathfrak{W}, \Gamma)$. Let $S \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ be any chart with $f: \mathcal{X} \rightarrow \mathcal{W}$ its universal family. Without loss of generality, we can assume that $\mathcal{W} = W[n] \times_{C[n]} S$ via a $\tau: S \rightarrow C[n]$. We let

$$(3.3) \quad S_\eta = S \times_{\mathfrak{M}(\mathfrak{W}, \Gamma)} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta).$$

In case $S_\eta = \emptyset$, we define $(\mathbf{L}_\eta, \mathbf{s}_\eta)|_S$ be $(\mathbf{1}_S, 1)$. When $S_\eta \neq \emptyset$, we consider the tautological projection $\rho_\eta: S_\eta \rightarrow \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ and the composite $\mathbf{p}_i \circ \rho_\eta: S_\eta \rightarrow \mathfrak{M}(\mathfrak{Y}_i^{\text{rel}}, \Gamma_i)$. Here

$$\mathbf{p}_i: \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) \longrightarrow \mathfrak{M}(\mathfrak{Y}_i^{\text{rel}}, \Gamma_i)$$

is the i -th projection, which exists if we replace S by an étale cover of S . By shrinking S if necessary, we can assume that the pull back of the universal family of $\mathfrak{M}(\mathfrak{Y}_i^{\text{rel}}, \Gamma_i)$ to S_η via the morphism $\mathbf{p}_i \circ \rho_\eta$ is given by a family $f_i: \mathcal{X}_i \rightarrow \mathcal{Y}_i$, where $\mathcal{Y}_i$ is an effective relative pair in $\mathfrak{Y}_i^{\text{rel}}(S_\eta)$ associated to a morphism $\tau_i: S_\eta \rightarrow \mathbf{A}^{n_i}$. Following the discussion leading to the proof of Proposition [Li, Prop. 4.12], the tautological family over S_η of the morphism $\Phi_\eta|_{S_\eta}: S_\eta \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ is then represented by the family

$$f_1 \sqcup f_2: \mathcal{X}_1 \sqcup \mathcal{X}_2 \longrightarrow \mathcal{Y}_1^\circ \sqcup \mathcal{Y}_2$$

with $\mathcal{Y}_1^\circ \sqcup \mathcal{Y}_2 \in \mathfrak{W}(S_\eta)$ given by the morphism $\tau_\eta: S_\eta \rightarrow C[n]$, where $n = n_1 + n_2$, defined in [Li, (4.4)]¹³. By definition $f_1 \sqcup f_2$ is isomorphic to the restriction of f to the family over S_η , denoted by $f|_{S_\eta}$. Namely there are isomorphisms shown below

¹¹For instance in case $J: [2] \rightarrow [4]$ is defined by $J(1) = 1$ and $J(2) = 3$, then $\gamma_J: \mathbf{A}^2 \rightarrow \mathbf{A}^4$ is defined by $\gamma_J(t_1, t_2) = (t_1, 1, t_2, 1)$.

¹²We use bold $\mathbf{1}_X$ with subscription X to denote the trivial line bundle on X .

¹³ $\mathcal{Y}_1^\circ \sqcup \mathcal{Y}_2$ is the result of gluing the distinguished divisors of $\mathcal{Y}_1$ and $\mathcal{Y}_2$ in the obvious way.

that make the following diagram commutative

$$(3.4) \quad \begin{array}{ccc} \mathcal{X}_1 \sqcup \mathcal{X}_2 & \xrightarrow{f_1 \sqcup f_2} & \mathcal{Y}_1^o \sqcup \mathcal{Y}_2 \\ \cong \downarrow & & \cong \downarrow \\ \mathcal{X} \times_S S_\eta & \xrightarrow{f|_{S_\eta}} & \mathcal{W} \times_S S_\eta. \end{array}$$

Now let $\mathbf{D}_1, \dots, \mathbf{D}_{n+1}$ be the $n+1$ components of the singular locus of the fibers of $W[n]$ over $C[n]$. For any closed $z \in S_\eta$ there is an integer $l_z \in [n+1]$ so that the images of the distinguished divisors $\mathcal{D}_{1,z} \subset \mathcal{Y}_{1,z}$ and of $\mathcal{D}_{2,z} \subset \mathcal{Y}_{2,z}$ under the obvious morphism

$$\mathcal{D}_{1,z} \cong \mathcal{D}_{2,z} \subset \mathcal{Y}_1^o \sqcup \mathcal{Y}_2 \xrightarrow{\cong} \mathcal{W} \times_S S_\eta \longrightarrow W[n]$$

lie in $\mathbf{D}_{l_z}$. Clearly, l_z is locally constant on S_η . Hence by shrinking S if necessary we can assume that it is constant on S_η , say is $l_\eta \in \mathbb{Z}$. In the following we will call $f|_{S_\eta} = f_1 \sqcup f_2$ the η -decomposition of f and call the divisor $\mathbf{D}_{l_\eta} \subset W[n]$ the locus where the η -decomposition of $f|_{S_\eta}$ takes place.

Definition 3.3. Let $S \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ be a chart with $f: \mathcal{X} \rightarrow \mathcal{W}$ the universal family, where $\mathcal{W} = W[n] \times_{C[n]} S$. We say S is η -admissible if there is an integer $l \in [n+1]$ so that the tautological $S_\eta \rightarrow C[n]$, where S_η is defined in (3.3), factor through $C[n] \times_{\mathbf{A}^{n+1}} H_l \subset C[n]$ and that the divisor $\mathbf{D}_l \subset W[n]$ (or the locus $\mathcal{X} \times_{W[n]} \mathbf{D}_l \subset \mathcal{X}$) is where the η -decomposition of $f|_{S_\eta}$ takes place.

Clearly for each $\eta \in \Omega$ we can find an atlas Λ of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ so that all its charts are η -admissible. Let Λ be such an atlas and let S_α be any chart in this atlas. We let $f_\alpha: \mathcal{X}_\alpha \rightarrow W[n_\alpha] \times_{C[n_\alpha]} S_\alpha$ be the universal family with $\tau_\alpha: S_\alpha \rightarrow C[n_\alpha]$ the tautological morphism. We let l_α be the integer so that $\mathbf{D}_{l_\alpha} \subset W[n_\alpha]$ is where the η -decomposition of $f_\alpha|_{S_{\alpha,\eta}}$ takes place. We then define the C-divisor $(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})$ on S_α to be the pull back of the C-divisor $(L_{l_\alpha}, s_{l_\alpha})$ on $\mathbf{A}^{n_\alpha+1}$ via $S \xrightarrow{\tau_\alpha} C[n_\alpha] \rightarrow \mathbf{A}^{n_\alpha+1}$. Because the chart S_α is η -admissible, the vanishing locus of $\mathbf{s}_{\eta,\alpha}$ is exactly $S_\alpha \cap \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$.

Lemma 3.4. The collection $(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})_{\alpha \in \Lambda}$ forms a C-divisor on $\mathfrak{M}(\mathfrak{W}, \Gamma)$.

Proof. Let S_α and S_β be two charts in Λ . We consider $S_{\alpha\beta} = S_\alpha \times_{\mathfrak{M}(\mathfrak{W}, \Gamma)} S_\beta$ with its projections ρ_α, ρ_β . Let f_α and f_β be the universal families over S_α and S_β and let $\rho_\alpha^*(f_\alpha)$ and $\rho_\beta^*(f_\beta)$ be the pull back families. We let the isomorphism of the families $\rho_\alpha^*(f_\alpha)$ and $\rho_\beta^*(f_\beta)$ be given by the diagram

$$(3.5) \quad \begin{array}{ccc} \mathcal{X}_\alpha \times_{S_\alpha} S_{\alpha\beta} & \xrightarrow{f_\alpha|_{S_{\alpha\beta}}} & \mathcal{W}_\alpha \times_{S_\alpha} S_{\alpha\beta} \\ \varphi_1 \downarrow \cong & & \varphi_2 \downarrow \cong \\ \mathcal{X}_\beta \times_{S_\beta} S_{\alpha\beta} & \xrightarrow{f_\beta|_{S_{\alpha\beta}}} & \mathcal{W}_\beta \times_{S_\beta} S_{\alpha\beta}. \end{array}$$

We distinguish two cases: The first is when $S_{\alpha\beta}$ is disjoint from $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$. Then the pull backs of $(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})$ and $(\mathbf{L}_{\eta,\beta}, \mathbf{s}_{\eta,\beta})$ to $S_{\alpha\beta}$ are canonically isomorphic to the trivial C-divisor $(\mathbf{1}_{S_{\alpha\beta}}, 1)$, hence they are naturally isomorphic to each other. The other case is when $S_{\alpha\beta} \cap \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) \neq \emptyset$. Since S_α and S_β are η -admissible, they have the associated morphisms $\tau_\alpha: S_\alpha \rightarrow C[n_\alpha]$ and $\tau_\beta: S_\beta \rightarrow C[n_\beta]$ and the associated integers l_α and l_β respectively. Because of the

isomorphisms in (3.5), the subscheme $\mathbf{D}_{l_\alpha} \times_{C[n_\alpha]} S_{\alpha\beta} \subset \mathcal{W}_\alpha \times_{S_\alpha} S_{\alpha\beta}$ is isomorphic to $\mathbf{D}_{l_\beta} \times_{C[n_\beta]} S_{\alpha\beta} \subset \mathcal{W}_\beta \times_{S_\beta} S_{\alpha\beta}$ under φ_2 . Now let T be any open subset of $S_{\alpha\beta}$ so that the restriction of φ_2 to $\mathcal{W}_\alpha \times_{S_\alpha} T \cong \mathcal{W}_\beta \times_{S_\beta} T$ is induced by a sequence of effective arrows¹⁴. Then by Lemma 3.2, the restriction to T of the pull back $\rho_\alpha^*(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})$ is canonically isomorphic to the restriction to T of $\rho_\beta^*(\mathbf{L}_{\eta,\beta}, \mathbf{s}_{\eta,\beta})$. By [Li, Lemma 1.8], we can cover $S_{\alpha\beta}$ by such T 's. Further, applying Lemma 3.2 again we see immediately that the isomorphisms $\rho_\alpha^*(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})|_T \cong \rho_\beta^*(\mathbf{L}_{\eta,\beta}, \mathbf{s}_{\eta,\beta})|_T$ patch together to form an isomorphism

$$\varphi_{\beta\alpha} : \rho_\alpha^*(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha}) \xrightarrow{\cong} \rho_\beta^*(\mathbf{L}_{\eta,\beta}, \mathbf{s}_{\eta,\beta}).$$

Since the isomorphism $\varphi_{\alpha\beta}$ canonically depends on the isomorphism (3.5), over $S_{\alpha\beta\gamma}$ we have $\varphi_{\alpha\beta} \circ \varphi_{\beta\gamma} = \varphi_{\alpha\gamma}$. Thus the collection $(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha})_{\alpha \in \Lambda}$ coupled with the isomorphisms $\varphi_{\alpha\beta}$ defines a C-divisor on $\mathfrak{M}(\mathfrak{W}, \Gamma)$. We denote the resulting C-divisor by $(\mathbf{L}_\eta, \mathbf{s}_\eta)$. $\square$

We now indicate the relation between the C-divisor $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ and the canonical log structure on $\mathfrak{M}(\mathfrak{W}, \Gamma)$. Recall that the log structure defined in section 1.1 defines a canonical log structure on $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$, which is $G[n]$ -equivariant and thus descends to a canonical log structure $\mathcal{N}$ on $\mathfrak{M}(\mathfrak{W}, \Gamma)$. The line bundle $\mathbf{L}_\eta$ with the section $\mathbf{s}_\eta$ defines also a log structure $\mathcal{L}_\eta$ on $\mathfrak{M}(\mathfrak{W}, \Gamma)$. It follows from the construction of N_α (see (1.7)) and the $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ that the identity map is a log morphism $(\mathfrak{M}(\mathfrak{W}, \Gamma), \mathcal{L}_\eta) \rightarrow (\mathfrak{M}(\mathfrak{W}, \Gamma), \mathcal{N})$. Using the chart $\mathcal{V}$ in (1.7), this is given by the homomorphism of pre-log structure $\mathbb{N} \rightarrow N_l \subset N_\mathcal{V}$ defined by $1 \mapsto m_\alpha e_\alpha$ for $\alpha \in K_l$.

Associating to each closed point $t \in C$, considered as an effective divisor in C , we have a C-divisor (L_t, r_t) such that $\mathcal{O}_C(L_t) = \mathcal{O}_C(t)$ and that r_t is the section induced by the constant section $1 \in \Gamma(C, \mathcal{O}_C)$ together with the natural homomorphism $\mathcal{O}_C(L_t) \rightarrow \mathcal{O}_C(t)$. We let $(\mathbf{L}_t, \mathbf{r}_t)$ be the C-divisor on $\mathfrak{M}(\mathfrak{W}, \Gamma)$ that is the pull back of (L_t, r_t) via the tautological projection $\mathfrak{M}(\mathfrak{W}, \Gamma) \rightarrow C$. When $t = 0 \in C$ we denote the corresponding C-divisor by $(\mathbf{L}_0, \mathbf{r}_0)$.

Let $\eta = (\Gamma_1, \Gamma_2, I) \in \Omega$ be any triple having r roots. Then for any permutation $\sigma \in S_r$ we have a new triple $\eta^\sigma = (\Gamma_1^\sigma, \Gamma_2^\sigma, I)$ that is derived by permuting the order of the r -ordered roots of Γ_1 and Γ_2 according to σ . We say two elements $\eta_1, \eta_2 \in \Omega$ are similar if there is a permutation σ so that $\eta_2 = \eta_1^\sigma$. We let $\bar{\Omega}$ be the set of similar classes of Ω . Note that when η_1 is similar to η_2 , then the C-divisors $(\mathbf{L}_{\eta_1}, \mathbf{s}_{\eta_1}) \equiv (\mathbf{L}_{\eta_2}, \mathbf{s}_{\eta_2})$.

Proposition 3.5. *The tensor product of the C-divisors $\{(\mathbf{L}_\eta, \mathbf{s}_\eta) \mid \eta \in \bar{\Omega}\}$ is isomorphic to $(\mathbf{L}_0, \mathbf{r}_0)$.*

Proof. Let Λ be an atlas of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ so that all its charts are η -admissible for all $\eta \in \bar{\Omega}$. Since $\bar{\Omega}$ is a finite set, such atlas does exist. Now let S_α be any chart in Λ . We let $\bar{\Omega}_\alpha$ be those triples $\eta \in \bar{\Omega}$ so that $\text{Im}(\Phi_\eta) \cap S_\alpha \neq \emptyset$. Recall that $(\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha}) \cong (1_{S_\alpha}, 1)$ canonically when $\eta \notin \bar{\Omega}_\alpha$. Therefore

$$\otimes_{\eta \in \bar{\Omega}} (\mathbf{L}_\eta, \mathbf{s}_\eta)|_{S_\alpha} \equiv \otimes_{\eta \in \bar{\Omega}_\alpha} (\mathbf{L}_{\eta,\alpha}, \mathbf{s}_{\eta,\alpha}).$$

Now let $f_\alpha : \mathcal{X}_\alpha \rightarrow \mathcal{W}_\alpha$ be the universal family over S_α and let $\tau_\alpha : S_\alpha \rightarrow C[n_\alpha]$ be so that $\mathcal{W}_\alpha = \tau_\alpha^* W[n_\alpha]$. To each $\eta \in \bar{\Omega}_\alpha$ we let l_η be the integer defined in Definition 3.3. The assignment $\eta \mapsto l_\eta$ defines a function $\bar{\Omega}_\alpha \rightarrow [n_\alpha + 1]$. Because

¹⁴For the definition of effective arrows please see [Li, Section 1].

of [Li, Lemma 4.13], this assignment is one-to-one. Now let $K_\alpha \subset [n_\alpha + 1]$ be the image set of this assignment and let $\bar{\tau}_\alpha: S_\alpha \rightarrow \mathbf{A}^{n_\alpha+1}$ be the composition of τ_α with the projection $C[n_\alpha] \rightarrow \mathbf{A}^{n_\alpha+1}$. Clearly, if $l \in [n_\alpha + 1] - K_\alpha$, then $\bar{\tau}_\alpha(S_\alpha) \cap \mathbf{H}_l = \emptyset$ and hence $\bar{\tau}_\alpha^*(L_l, s_l) \equiv (\mathbf{1}_{S_\alpha}, 1)^{15}$. Otherwise, $l = l_\eta$ for a unique $\eta \in \bar{\Omega}_\alpha$ and then $\bar{\tau}_\alpha^*(L_l, s_l) \equiv (\mathbf{L}_{\eta, \alpha}, \mathbf{s}_{\eta, \alpha})$. Therefore, we have canonical isomorphisms

$$\otimes_{\eta \in \bar{\Omega}_\alpha} (\mathbf{L}_{\eta, \alpha}, \mathbf{s}_{\eta, \alpha})|_{S_\alpha} \cong \otimes_{l \in K_\alpha} \bar{\tau}_\alpha^*(L_l, s_l) \cong \bar{\tau}_\alpha^*(\otimes_{l \in K_\alpha} (L_l, s_l)) \cong (\mathbf{L}_0, \mathbf{r}_0)|_{S_\alpha}.$$

Because the above isomorphisms are canonical, they are compatible over $S_{\alpha\beta}$ and hence define an isomorphism of C-divisors as required by the Proposition. $\square$

We now derive the first version of the degeneration formula. We first recall the notion of localized top Chern class of a vector bundle with a section. Let E be a rank m vector bundle over a scheme X and s a section of E . The localized top Chern class of (E, s) is the homomorphism

$$c_m(E, s): A_*X \rightarrow A_{*-m}s^{-1}(0)$$

defined in [Ful] as follows: Let Z be any variety in X . We take the normal cone $N_{s^{-1}(0) \cap Z/Z}$ to $s^{-1}(0) \cap Z$ in Z and then define $c_m(E, s)([Z]) = 0_E^!(N_{s^{-1}(0) \cap Z/Z})$, where $0_E^!$ is the Gysin map of the zero section 0_E of $E|_{s^{-1}(0)}$. This defines a homomorphism of the group of cycles. This construction can be extended to the case where X is an algebraic stack [Vis] with A_*X understood to be the cycle group with rational coefficients.

We consider the moduli stack $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and its Gromov-Witten invariants

$$\Psi_\Gamma^{W/C}: H_C^0(R^*\pi_*\mathbb{Q}_W)^{\times k} \times H^*(\mathfrak{M}_{g,k}) \longrightarrow H_2^{\text{BM}}(C) \cong \mathbb{Q}.$$

Now let $\xi \in C$ be any closed point and let $H_C^0(R^*\pi_*\mathbb{Q}_W) \rightarrow H^*(W_\xi)$, denoted by $\alpha \mapsto \alpha(\xi)$, be induced by $W_\xi \rightarrow W$. We let $H_2^{\text{BM}}(C) \rightarrow \mathbb{Q}$ be the Gysin homomorphism defined by intersecting with the divisor $\xi \in C$. Then we have a commutative diagram

$$(3.6) \quad \begin{array}{ccc} H_C^0(R^*\pi_*\mathbb{Q}_W)^{\times k} \times H^*(\mathfrak{M}_{g,k}) & \xrightarrow{\Psi_\Gamma^{W/C}} & H_2^{\text{BM}}(C) \cong \mathbb{Q} \\ \downarrow & & \downarrow \\ H^*(W_\xi)^{\times k} \times H^*(\mathfrak{M}_{g,k}) & \xrightarrow{\Psi_\Gamma^{W_\xi}} & \mathbb{Q}. \end{array}$$

This was proved in [LT1] for $\xi \neq 0$ except that there we used the existence of a global vector bundle in defining the GW-invariants. This will be proved later in this paper again. We let $\Gamma = (g, k, b)$ be the triple as before and let $\alpha \in H^0(R^*\pi_*\mathbb{Q}_W)^{\times k}$ and let $\beta \in H^*(\mathfrak{M}_{g,k})$ be any classes. As before, we let $ev_\xi: \mathfrak{M}(W_\xi, \Gamma) \rightarrow W_\xi^{\times k}$ be the evaluation morphism by the ordinary marked points of the stable morphisms, and let $\pi_{g,k}: \mathfrak{M}(\mathfrak{W}, \Gamma) \rightarrow \mathfrak{M}_{g,k}$ be the forgetful map.

Theorem 3.6. *For any closed $\xi \in C$, we have*

$$\Psi_\Gamma^{W_\xi}(\alpha(\xi), \beta) = \mathbf{q}_{*0} \left(\sum_{\eta \in \bar{\Omega}} (ev_0^*(\alpha(0)) \cup \pi_{g,k}^*(\beta)) (c_1(\mathbf{L}_\eta, \mathbf{s}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}) \right).$$

¹⁵There is an exceptional case I should mention here. It is when there are $s \in S_\alpha$ so that $\bar{\tau}_\alpha(s) \in H_l$ while $f_\alpha^{-1}(\mathbf{D}_l) \cap \mathcal{X}_s = \emptyset$. Note that since $\mathcal{X}_s$ is connected, this is possible only when $l = 1$ or $n_\alpha + 1$. In either case, we agree that $\mathcal{X}_s$ decomposes into $C_1 \sqcup C_2$ with either $C_1 = \emptyset$ or $C_2 = \emptyset$, and the corresponding $\eta = (\Gamma_1, \emptyset, \emptyset)$ or $\eta = (\emptyset, \Gamma_2, \emptyset)$, which we agree is in $\bar{\Omega}$. In this case we let l_η be 1 or $n_\alpha + 1$. With this agreement, this statement is true without exception.

Proof. First, by the commutativity of the diagram (3.6), we have $\Psi_\Gamma^{W_\xi}(\alpha(\xi), \beta) = \Psi_\Gamma^{W_0}(\alpha(0), \beta)$. In the later part of this paper, we will show that

$$\Psi_\Gamma^{W_0}(\alpha(0), \beta) = \mathbf{q}_{*0} \left((ev_0^*(\alpha(0)) \cup \pi_{g,k}^*(\beta)) (c_1(\mathbf{L}_0, \mathbf{r}_0)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}) \right).$$

However, since $(\mathbf{L}_0, \mathbf{r}_0)$ is the tensor product of all $(\mathbf{L}_\eta, \mathbf{s}_\eta)$, the Chern class operations

$$c_1(\mathbf{L}_0, \mathbf{r}_0) = \sum_{\eta \in \bar{\Omega}} c_1(\mathbf{L}_\eta, \mathbf{s}_\eta) : A_* \mathfrak{M}(\mathfrak{W}, \Gamma) \rightarrow A_* \mathfrak{M}(\mathfrak{W}_0, \Gamma).$$

The theorem then follows immediately. $\square$

3.2. Statement of the degeneration formula. In this subsection, we will first construct the virtual moduli cycles of several moduli stacks related to the substack $\mathfrak{W}_0 \subset \mathfrak{W}$. After that, we will state the final version of the degeneration formula of the Gromov-Witten invariants of this paper. We will leave the proof of the key Lemmas to the next section.

Let $\eta \in \bar{\Omega}$ be any admissible triple. Associated to η we have the substack $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) \subset \mathfrak{M}(\mathfrak{W}, \Gamma)$ that is the image stack of Φ_η in (3.1). Let $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ be the $\mathbf{C}$ -divisor on $\mathfrak{M}(\mathfrak{W}, \Gamma)$ defined in the previous subsection. We define the substack

$$\mathfrak{M}(\mathfrak{W}_0, \eta) = \mathbf{s}_\eta^{-1}(0) \subset \mathfrak{M}(\mathfrak{W}, \Gamma).$$

Note that we have an increasing chain of closed substacks

$$\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) \subset \mathfrak{M}(\mathfrak{W}_0, \eta) \subset \mathfrak{M}(\mathfrak{W}_0, \Gamma) \subset \mathfrak{M}(\mathfrak{W}, \Gamma),$$

where the first inclusion induces a homeomorphism on topological spaces. In the previous section, we have constructed the perfect-obstruction theories of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$, and thus have constructed their virtual moduli cycles. In the first part of this subsection, we will show that the natural obstruction theories of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ and of $\mathfrak{M}(\mathfrak{W}_0, \eta)$ are also perfect. Thus they have natural virtual moduli cycles.

We first investigate the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)$. The discussion is parallel to the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. We now present the details. Recall that for each n we have the moduli $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ of stable morphisms to $W[n]$ of topological type Γ that are also stable as morphisms to the stack $\mathfrak{W}$. We let $\mathfrak{M}(W[n], \Gamma)^{\text{st}} \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ be the tautological morphism and let

$$\mathfrak{M}(W_0[n], \eta)^{\text{st}} = \mathfrak{M}(W[n], \Gamma)^{\text{st}} \times_{\mathfrak{M}(\mathfrak{W}, \Gamma)} \mathfrak{M}(\mathfrak{W}_0, \eta).$$

As in the case of $\mathfrak{M}(\mathfrak{W}, \Gamma)$, to study the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)$ it suffices to work out the obstruction theory of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}$, which we will do now. Of course for étale charts S of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ we can define the notion of η -admissible as in Definition 3.3. Now let S be an η -admissible chart of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ and let $S_\eta = S \times_{\mathfrak{M}(W[n], \Gamma)^{\text{st}}} \mathfrak{M}(W_0[n], \eta)^{\text{st}}$. As before, we let $f: \mathcal{X} \rightarrow W[n]$ be the universal family over S . We let l be the integer so that $f^{-1}(\mathbf{D}_l)$ is where the η -decomposition of $f|_{S_\eta}$ takes place. We fix a covering of f by charts $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha)$ of the first and the second kinds, indexed by Λ . Following our convention, we denote by $\underline{\mathcal{U}}$ the étale covering $\{\mathcal{U}_\alpha\}_\Lambda$ of $\mathcal{X}$. We let $f_{\eta, \alpha}: \mathcal{U}_{\eta, \alpha} \rightarrow W[n]$ be the restriction of f_α to $\mathcal{U}_{\eta, \alpha} \triangleq \mathcal{U}_\alpha \times_S S_\eta$. We let $\mathcal{V}_{\eta, \alpha} = \mathcal{V}_\alpha \times_S S_\eta$ and let $\underline{\mathcal{V}}_\eta$ be the covering $\{\mathcal{V}_{\eta, \alpha}\}_\Lambda$ of S_η . We let $A = \Gamma(\mathcal{O}_S)$ and let $\mathbf{E}^\bullet \equiv \mathbf{E}(A)^\bullet$ be the complex constructed in section

2 associated to the covering Λ . Now let L_η be the line bundle on S_η that is the pull back of $\mathbf{L}_\eta$ via $S_\eta \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$. We form the ordinary Čech complex

$$(3.7) \quad \mathbf{C}_\eta^\bullet \triangleq \mathbf{C}^\bullet(\underline{\mathcal{V}}_\eta, L_\eta)$$

of the invertible sheaf (line bundle) L_η over S_η associated to the covering $\underline{\mathcal{V}}_\eta$. Let $A_\eta = \Gamma(\mathcal{O}_{S_\eta})$. We now define a homomorphism of the complex:

$$(3.8) \quad \delta : \mathbf{E}^\bullet \otimes_A A_\eta \Longrightarrow \mathbf{C}_\eta^{\bullet-1}$$

as follows: Let $(a, b) \in \mathbf{E}^1 \otimes_A A_\eta$ be any element. As argued in the proof of Proposition 1.16, a defines, to each $\alpha \in \Lambda$, a flat extension $\tilde{\mathcal{U}}_{\eta, \alpha} / \tilde{\mathcal{V}}_{\eta, \alpha}$ of $\mathcal{U}_{\eta, \alpha} / \mathcal{V}_{\eta, \alpha}$ by the module $\Gamma(\mathcal{O}_{\mathcal{V}_{\eta, \alpha}})$ and pre-deformable extensions $\zeta_{\eta, \alpha} : \tilde{\mathcal{U}}_{\eta, \alpha} \rightarrow W[n]$ of $f_{\eta, \alpha} : \mathcal{U}_{\eta, \alpha} \rightarrow W[n]$. Let $\rho_{\eta, \alpha} : \mathcal{V}_{\eta, \alpha} \rightarrow \mathbf{A}^{n+1}$ be the tautological morphism (induced by $S \rightarrow \mathbf{A}^{n+1}$) and let $\tilde{\rho}_{\eta, \alpha} : \tilde{\mathcal{V}}_{\eta, \alpha} \rightarrow \mathbf{A}^{n+1}$ be the composite of $\zeta_{\eta, \alpha}$ and the projection $W[n]_\alpha \rightarrow \mathbf{A}^{n+1}$. Since $t_l \circ \rho_{\eta, \alpha} = 0$,

$$\xi_\alpha(a) \triangleq \mathbf{d}(t_l \circ \tilde{\rho}_{\eta, \alpha} - 0) \in \Gamma(\mathcal{V}_{\eta, \alpha}, \rho_{\eta, \alpha}^* N_{\mathbf{H}_l / \mathbf{A}^{n+1}}) \equiv \Gamma(\mathcal{V}_{\eta, \alpha}, L_\eta),$$

where $N_{\mathbf{H}_l / \mathbf{A}^{n+1}}$ is the normal bundle to $\mathbf{H}_l$ in $\mathbf{A}^{n+1}$. Here we have used the fact that $\rho_{\eta, \alpha}^* N_{\mathbf{H}_l / \mathbf{A}^{n+1}}$ is canonically isomorphic to L_l . On the other hand, for $b = \{b_\alpha\}$ with $b_\alpha \in \text{Hom}_{\mathcal{U}_\alpha}(f^* \Omega_{W[n]}, A_\eta)^\dagger$, $b_\alpha(f^*(dt_l)) \in I$, where $\pi_n : W[n] \rightarrow \mathbf{A}^{n+1}$ is the projection. We define $\xi(b_\alpha) = b_\alpha(f^*(dt_l))$. This defines a homomorphism

$$(3.9) \quad \delta_k : \mathbf{E}^k \otimes_A A_\eta \longrightarrow \mathbf{C}_\eta^{k-1} \triangleq \mathbf{C}^{k-1}(\underline{\mathcal{V}}_\eta, L_\eta)$$

via $\delta_1((a, b))_\alpha = \xi_\alpha(a) + \xi(b_\alpha)$ for $k = 1$. Note that for $k > 1$ elements in $\mathbf{E}^k$ are of the form $b = \{b_{\alpha_0 \dots \alpha_k}\}$ and we can define $\delta_k(b_{\alpha_0 \dots \alpha_k})$ similarly. This defines δ_k for $k > 1$. It is direct to check that the so defined map is a homomorphism of complexes. With this homomorphism of complexes, we can form a complex

$$(3.10) \quad \mathbf{E}_\eta^k \triangleq \mathbf{C}_\eta^{k-2} \oplus \mathbf{E}^k \otimes_A A_\eta$$

with the induced differential. It is clear that the statement of Lemma 1.15 holds true to $\mathbf{E}_\eta^\bullet$. Namely, for sufficiently fine admissible covering Λ of f , $\mathbf{E}_\eta^\bullet$ is a complex of flat A_η -modules. Further, to each A_η -module I we have the following exact sequence of complexes

$$(3.11) \quad 0 \Longrightarrow \mathbf{C}_\eta^{\bullet-2} \otimes_{A_\eta} I \Longrightarrow \mathbf{E}_\eta^\bullet \otimes_{A_\eta} I \Longrightarrow \mathbf{E}^\bullet \otimes_{A_\eta} I \Longrightarrow 0$$

which induces a long exact sequence of cohomologies

$$(3.12) \quad \begin{aligned} 0 \longrightarrow h^1(\mathbf{E}_\eta^\bullet \otimes_{A_\eta} I) &\longrightarrow h^1(\mathbf{E}^\bullet \otimes_A I) \longrightarrow I \otimes_{A_\eta} \mathcal{O}_{S_\eta}(L_\eta) \longrightarrow \\ &\longrightarrow h^2(\mathbf{E}_\eta^\bullet \otimes_{A_\eta} I) \longrightarrow h^2(\mathbf{E}^\bullet \otimes_A I) \longrightarrow 0. \end{aligned}$$

In particular, $h^k(\mathbf{E}_\eta^\bullet \otimes_{A_\eta} I) = 0$ except $k = 1, 2$.

Proposition 3.7. *First, the functor of the first order deformations of S_η is naturally isomorphic to the functor $\mathfrak{h}^1(\mathbf{E}_\eta^\bullet)$. Secondly, there is a natural obstruction theory to deformation of the families of pre-deformable morphisms $f|_{S_\eta} : \mathcal{X}|_{S_\eta} \rightarrow W[n] \times_{\mathbf{A}^{n+1}} \mathbf{H}_l$ taking values in $\mathfrak{h}^2(\mathbf{E}_\eta^\bullet)$. Finally, such obstruction theory is perfect.*

Proof. We will omit the proof here because it is parallel to the treatment of the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \Gamma)$. $\square$

Proposition 3.8. *The perfect obstruction theory constructed in Proposition 3.7 induces a perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)$, which in turn defines a natural virtual moduli cycle $[\mathfrak{M}(\mathfrak{W}_0, \eta)]^{\text{virt}}$.*

Proof. The proof is parallel to the construction of the perfect obstruction theory of $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and of the virtual moduli cycle $[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}$. We shall omit the details here. $\square$

We next work out the obstruction theory of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$. We let $S = \text{Spec } A \rightarrow \mathfrak{M}(W[n], \Gamma)^{\text{st}}$ be an η -admissible chart and let $f : \mathcal{X} \rightarrow W[n]$ be the universal family over S . We then pick a sufficiently fine covering $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha)$ indexed by Λ . Following section 1.2, we pick the complex $[F^0 \xrightarrow{d} F^1]$ as in (1.5) and pick extensions ζ_α as defined before (1.19). Based on these data we can form the complex $\mathbf{E}^\bullet$ as in (1.20) so that its cohomology is part of the obstruction theory of S . We let $S_0 = \text{Spec } A_0 = S \times_{\mathfrak{M}(\mathfrak{W}, \Gamma)} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ and let $f_0 : \mathcal{X}_0 \rightarrow W[n]$ be the restriction of f to fibers over S_0 . We let $l \in [n+1]$ be the integer associated to η defined in Definition 3.3. By definition, the family f_0 can be decomposed into two families of relative stable morphisms of types Γ_1 and Γ_2 respectively along a multi-section $\Sigma \subset \mathcal{X}_{0, \text{node}}$ over S_0 . We let $\mathcal{U}_{0, \alpha} = \mathcal{U}_\alpha \times_S S_0$, $\mathcal{V}_{0, \alpha} = \mathcal{V}_\alpha \times_S S_0$ and let $f_{0, \alpha} = f_\alpha|_{\mathcal{U}_{0, \alpha}}$. Note that Σ is étale over S_0 . Now let $\hat{\mathcal{X}}_0$ be the formal completion of $\mathcal{X}_0$ along $\Sigma \subset \mathcal{X}_0$. Then since $\Sigma \rightarrow S$ is finite and étale, and since $\Sigma \subset \mathcal{X}_0$ is a multiple section of the nodal points of the fibers of $\mathcal{X}_0/S_0$, the extension sheaf $\mathcal{E}xt_{\mathcal{X}_0}^1(\Omega_{\hat{\mathcal{X}}_0/S_0}, \mathcal{O}_{\hat{\mathcal{X}}_0})$ is an invertible sheaf of $\mathcal{O}_\Sigma$ -modules. We denote this sheaf by $\mathcal{M}_\Sigma$. Then we have a natural homomorphism

$$\text{Ext}_{\mathcal{X}_0}^1(\Omega_{\mathcal{X}_0/S_0}, \mathcal{O}_{\mathcal{X}_0}) \longrightarrow \rho_* (\mathcal{E}xt_{\mathcal{X}_0}^1(\Omega_{\mathcal{X}_0/S_0}, \mathcal{O}_{\mathcal{X}_0})) \equiv \rho_* \mathcal{M}_\Sigma,$$

where $\rho : \mathcal{X}_0 \rightarrow S_0$ is the projection, which defines a canonical homomorphism

$$(3.13) \quad F^1 \otimes_A A_0 \longrightarrow \text{Ext}_{\mathcal{X}_0}^1(\Omega_{\mathcal{X}_0/S_0}, \mathcal{O}_{\mathcal{X}_0}) \longrightarrow \bar{\rho}_* \mathcal{M}_\Sigma.$$

Here $\bar{\rho} : \Sigma \rightarrow S_0$ is the tautological projection induced by ρ . Clearly, the composite of (3.13) is surjective. We let $F_{\eta, 0}^0 = F^0 \otimes_A A_0$ and let $F_{\eta, 0}^1$ be the kernel of (3.13). Clearly, $F_{\eta, 0}^1$ is a free A_0 -module and $F^0 \otimes_A A_0 \rightarrow F^1 \otimes_A A_0$ factor through $F_{\eta, 0}^0 \rightarrow F_{\eta, 0}^1$.

We now construct the complex that will give the obstruction theory of S_0 (which is a chart of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$). We let $\mathbf{E}^\bullet$ be the complex associated to the family f and the covering Λ mentioned before. We let $\mathbf{E}_{\eta, 0}^k \triangleq \mathbf{E}^k \otimes_A A_0$ for $k \neq 1$ and let $\mathbf{E}_{\eta, 0}^1$ be the kernel of the composite $\mathbf{E}^1 \otimes_A A_0 \xrightarrow{\text{pr}} F^1 \otimes_A A_0 \rightarrow \bar{\rho}_* \mathcal{M}_\Sigma$. Clearly, the differentials in $\mathbf{E}^\bullet$ induce differentials in $\mathbf{E}_{\eta, 0}^\bullet$.

Proposition 3.9. *The Lemma 1.15 holds true for the complex $\mathbf{E}_{\eta, 0}^\bullet$. The Proposition 3.7 holds true for the charts S_0 with $\mathbf{E}_\eta^\bullet$ replaced by $\mathbf{E}_{\eta, 0}^\bullet$. The Proposition 3.8 holds true for the moduli stack $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$.*

Proof. We shall sketch the construction of the obstruction classes. The remainder part of the proof is similar to that of the Propositions 3.7 and 3.8 and will be omitted. Here we will follow closely the convention introduced in the proof of Proposition 1.18. Let $\xi = (B, I, \varphi)$ be an object in $\mathfrak{Tris}_0$. Let $f_T : \mathcal{X}_T \rightarrow W[n]$ be the pull back of f via $T \rightarrow S_0$, where $T = \text{Spec } B/I$, and let $\Sigma_T = \mathcal{X}_T \times_{\mathcal{X}_0} \Sigma$. By the definition of the subscheme $S_0 \subset S$, the formal completion of $\mathcal{X}_T$ along Σ_T is isomorphic to $\text{Spec } \mathbb{k}[[z_1, z_2]]/(z_1 z_2) \times \Sigma_T$, at least after shrinking S if necessary. Now let $\tilde{T} = \text{Spec } B$. Then by the deformation theory of nodal curves, we can

find a flat extension $\mathcal{X}_{\tilde{T}}/\tilde{T}$ of $\mathcal{X}_T/T$ so that the formal completion of $\mathcal{X}_{\tilde{T}}$ along Σ_T is isomorphic to $\mathrm{Spec} \mathbb{k}[[z_1, z_2]]/(z_1 z_2) \times \Sigma_{\tilde{T}}$, where $\Sigma_{\tilde{T}}$ is étale over $\tilde{T}$ so that $\Sigma_{\tilde{T}} \times_{\tilde{T}} T \equiv \Sigma_T$. In other words, the multiple section $\Sigma_T \subset \mathcal{X}$ extends to a multiple section $\Sigma_{\tilde{T}} \subset \mathcal{X}_{\tilde{T}}$ and the extended family $\mathcal{X}_{\tilde{T}}$ can be decomposed along $\Sigma_{\tilde{T}}$. Once we have chosen such an extension, to each $\alpha \in \Lambda$ we can pick a pre-deformable extension $\tilde{h}_\alpha : \tilde{\mathcal{U}}_{0,\alpha} \rightarrow W[n]$ of $f_{T,\alpha}$ so that the composite $\tilde{\mathcal{U}}_{0,\alpha} \rightarrow W[n] \rightarrow \mathbf{A}^{n+1}$ factor through the l -th coordinate hyperplane $\mathbf{H}_l \subset \mathbf{A}^{n+1}$. We let $b_{\alpha,\beta}$ be as defined in (1.22). Then $b = \{b_{\alpha,\beta}\} \in \mathbf{E}^2 \otimes_A I \equiv \mathbf{E}_{\eta,0}^2 \otimes_{A_0} I$. We define

$$\mathbf{ob}(\xi) = [b] \in h^2(\mathbf{E}_{\eta,0}^\bullet \otimes_{A_0} I).$$

It is direct to check that this defines an obstruction class. It is routine to check that such choices of obstruction classes satisfies the base change property. $\square$

Applying the basic construction of virtual moduli cycles formulated in the previous section, we obtain cycles $[\mathfrak{M}(\mathfrak{W}_0, \eta)]^{\mathrm{virt}}$ and $[\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)]^{\mathrm{virt}}$.

We are now ready to state the degeneration formula of the Gromov-Witten invariants of the family W . Let $\eta = (\Gamma_1, \Gamma_2, I)$ be an admissible triple in $\bar{\Omega}$. Let μ_i be the weight of the i -th root of Γ_1 and Γ_2 , which are the same. We define the multiplicity of η to be $\mathbf{m}(\eta) = \prod_{i=1}^r \mu_i$.

Lemma 3.10. *We have the identity*

$$c_1(\mathbf{L}_0, \mathbf{s}_0)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\mathrm{virt}} = [\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\mathrm{virt}} \in A_* \mathfrak{M}(\mathfrak{W}_0, \Gamma).$$

Lemma 3.11. *We have the identity*

$$[\mathfrak{M}(\mathfrak{W}_0, \eta)]^{\mathrm{virt}} = c_1(\mathbf{L}_\eta, \mathbf{s}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\mathrm{virt}} \in A_* \mathfrak{M}(\mathfrak{W}_0, \eta).$$

Lemma 3.12. *Under the natural isomorphism*

$$A_* \mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta) \cong A_* \mathfrak{M}(\mathfrak{W}_0, \eta)$$

induced by the homeomorphism $\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta) \simeq \mathfrak{M}(\mathfrak{W}_0, \eta)$,

$$\mathbf{m}(\eta)[\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)]^{\mathrm{virt}} = [\mathfrak{M}(\mathfrak{W}_0, \eta)]^{\mathrm{virt}} \in A_* \mathfrak{M}(\mathfrak{W}_0, \eta).$$

Using the identity of C-divisors $(\mathbf{L}_0, \mathbf{s}_0) = \otimes_{\eta \in \bar{\Omega}} (\mathbf{L}_\eta, \mathbf{s}_\eta)$, we have

Corollary 3.13.

$$[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\mathrm{virt}} = \sum_{\eta \in \bar{\Omega}} \mathbf{m}(\eta) [\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)]^{\mathrm{virt}}.$$

We now state how virtual moduli cycles $[\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)]^{\mathrm{virt}}$ is related to $[\mathfrak{M}(\mathfrak{Y}_i^{\mathrm{rel}}, \Gamma_i)]^{\mathrm{virt}}$. As mentioned in [Li], we have the natural evaluation morphism $\mathfrak{M}(\mathfrak{Y}_i^{\mathrm{rel}}, \Gamma_i) \rightarrow D^r$ and the induced Cartesian diagram

$$(3.14) \quad \begin{array}{ccc} \mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\mathrm{rel}}, \Gamma_2) & \longrightarrow & \mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}}, \Gamma_1) \times \mathfrak{M}(\mathfrak{Y}_2^{\mathrm{rel}}, \Gamma_2) \\ \downarrow & & \downarrow \\ D^r & \xrightarrow{\Delta} & D^r \times D^r \end{array}$$

Here the arrow Δ is the diagonal morphism. Let Φ_η be the finite étale morphism in (3.2), which has pure degree $|\mathrm{Eq}(\eta)|$ (see [Li, Section 4]).

Lemma 3.14. *We have the identity*

$$\frac{1}{|\text{Eq}(\eta)|} \Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}) = [\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)]^{\text{virt}}.$$

We will prove these Lemmas in the next section.

The main degeneration formula of the Gromov-Witten invariants of W follows immediately from these Lemmas and the first version of the degeneration formula proved in the previous subsection.

Theorem 3.15. *Let the notation be as before. Then as elements in $A_*\mathfrak{M}(\mathfrak{W}_0, \Gamma)$,*

$$[\mathfrak{M}(\mathfrak{W}_0, \Gamma)]^{\text{virt}} = \sum_{\eta \in \bar{\Omega}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}).$$

Finally, we state the numerical corollary of this theorem. Let $j_i: Y_i \rightarrow W$ be the inclusion and let

$$j_i^*: H_C^0(R^* \pi_* \mathbb{Q}_W)^{\times k} \rightarrow H^*(Y_i, \mathbb{Q})^{\times k}$$

be the induced pull back homomorphism. Now let $\eta = (\Gamma_1, \Gamma_2, I) \in \bar{\Omega}$ be any admissible triple. For $i = 1$ or 2 , we let $\mathfrak{M}_{\Gamma_i^\circ}$ be the moduli space of stable curves of topological type Γ_i° (See the definition before (2.9)). It is naturally a Deligne-Mumford stack. Further, we have a natural local immersion of stacks

$$\phi_\eta: \mathfrak{M}_{\Gamma_1^\circ} \times \mathfrak{M}_{\Gamma_2^\circ} \longrightarrow \mathfrak{M}_{g,k}$$

that associates to any pair of curves $(C_1, C_2) \in \mathfrak{M}_{\Gamma_1^\circ} \times \mathfrak{M}_{\Gamma_2^\circ}$ the gluing $C_1 \sqcup C_2$ by identifying the i -th distinguished marked point of C_1 with the i -th distinguished marked point of C_2 for all i . Now let $\beta \in H^*(\mathfrak{M}_{g,k})$. We assume β has the following Kunnet type decomposition

$$\phi_\eta^*(\beta) = \sum_{j \in K_\eta} \beta_{\eta,1,j} \boxtimes \beta_{\eta,2,j}, \quad \beta_{i,j} \in H^*(\mathfrak{M}_{\Gamma_i^\circ}).$$

Corollary 3.16. *Let W/C be the family and let $\Gamma = (g, b, k)$ be as before. Then for any closed point $\xi \neq 0 \in C$, $\alpha \in H_C^0(R^* \pi_* \mathbb{Q}_W)^{\times k}$ and $\beta \in H^*(\mathfrak{M}_{g,n})$ as before,*

$$\Phi_\Gamma^{W\xi}(\alpha(\xi), \beta) = \sum_{\eta \in \bar{\Omega}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \sum_{j \in K_\eta} [\Psi_{\Gamma_1^1}^{Y_1^{\text{rel}}} (j_1^* \alpha, \beta_{\eta,1,j}) \bullet \Psi_{\Gamma_2^2}^{Y_2^{\text{rel}}} (j_2^* \alpha, \beta_{\eta,2,j})]_0.$$

Here $\bullet$ is the intersection of the homology groups

$$H_*(D^r) \times H_*(D^r) \xrightarrow{\cap} H_*(D^r)$$

and $[\gamma]_0$ is the degree 0 part of the homology class $\gamma \in H_*(D^r)$.

4. PROOF OF THE MAIN THEOREM

The goal of this section is to prove Lemma 3.12-3.14. In essence, the proofs of these Lemmas (except Lemma 3.12) rely on the comparison of the virtual moduli cycles of stacks with the virtual moduli cycles of their substacks. This is precisely the situation studied in [LT2, Lemma 3.4], except that there we used the existence of certain locally free sheaves to construct the virtual moduli cycles. To prove Lemma 3.12, we need to study the situation more general than the one studied. In the first subsection we will revise [LT2, Lemma 3.4] to cover all the situations we need.

4.1. Comparison of the virtual moduli cycles. Let $\mathbf{M} \rightarrow \mathbf{N}$ be a representable morphism of stacks. In this subsection, we assume $\mathbf{M}$ is a DM-stack having a perfect-obstruction theory with the associated obstruction sheaves $\mathcal{O}b_{\mathbf{M}}$. For $\mathbf{N}$ we need to consider two possibilities: One is when $\mathbf{N}$ is a DM-stack having a perfect-obstruction theory with the obstruction sheaf $\mathcal{O}b_{\mathbf{N}}$. The other is when $\mathbf{N}$ is a closed substack of a smooth Artin stack $\mathbf{Q}$ defined by the vanishing of a section s of a vector bundle F on $\mathbf{Q}$.

Let $\mathcal{S} = \{S_\alpha\}_\Lambda$ be an atlas of $\mathbf{N}$ and let $\{(\mathcal{F}_\alpha^\bullet, \mathbf{ob}_{S_\alpha})\}_\Lambda$ be the data associated to the perfect-obstruction theory of $\mathbf{N}$. In case $\mathbf{N}$ is a DM-stack this is specified in the Definition 2.1. In case $\mathbf{N}$ is an Artin stack, we assume that there is an atlas $\tilde{\mathcal{S}} = \{\tilde{S}_\alpha\}_\Lambda$ of $\mathbf{Q}$ so that $S_\alpha = \tilde{S}_\alpha \times_{\mathbf{Q}} \mathbf{N}$. Then we simply take $\mathcal{F}_\alpha^\bullet = [\mathcal{F}_\alpha^1 \rightarrow \mathcal{F}_\alpha^2]$ to be $\mathcal{F}_\alpha^1 = \mathcal{O}_{S_\alpha}(T\tilde{S}_\alpha)$, $\mathcal{F}_\alpha^2 = \mathcal{O}_{S_\alpha}(F|_{S_\alpha})$ and the arrow $\mathcal{F}_\alpha^1 \rightarrow \mathcal{F}_\alpha^2$ to be the one induced by the differential of the section $s \in H^0(F)$. The obstruction assignment $\mathbf{ob}_{S_\alpha}$ taking values in $\mathcal{O}b_{S_\alpha} = h^2(\mathcal{F}_\alpha^\bullet)$ is the obvious one induced by the defining equation s . Since $\mathbf{M} \rightarrow \mathbf{N}$ is representable, $S_\alpha \times_{\mathbf{N}} \mathbf{M}$ is a scheme. For each $\alpha \in \Lambda$ we pick an affine étale universal open $R_\alpha \rightarrow S_\alpha \times_{\mathbf{N}} \mathbf{M}$. Without loss of generality, we can assume $\{R_\alpha\}_\Lambda$ is a covering of $\mathbf{M}$ in the sense that the image $\mathbf{R}_\alpha \triangleq \rho_\alpha(R_\alpha)$ of the tautological $p_\alpha : R_\alpha \rightarrow \mathbf{M}$ is an open substack and the collection $\{\mathbf{R}_\alpha\}_\Lambda$ forms an open covering of $\mathbf{M}$. Note that when $\mathbf{N}$ is a DM-stack, $R_\alpha \rightarrow \mathbf{M}$ are étale and then $\{R_\alpha\}_\Lambda$ forms an étale cover of $\mathbf{M}$. In case $\mathbf{N}$ is an Artin stack, then $R_\alpha \rightarrow \mathbf{M}$ are smooth morphisms. In this case we shall view $\{R_\alpha\}_\Lambda$ as an atlas in the smooth site¹⁶ of $\mathbf{M}$.

We let $\{(\mathcal{E}_\alpha^\bullet, \mathbf{ob}_{R_\alpha})\}_\Lambda$ be the data given by the perfect obstruction theory of $\mathbf{M}$ associated to the covering $\{R_\alpha\}_\Lambda$. Namely, each $(\mathcal{E}_\alpha^\bullet, \mathbf{ob}_{R_\alpha})$ is a perfect obstruction theory of R_α , the sheaves $\text{Coker}\{\mathcal{E}_\alpha^1 \rightarrow \mathcal{E}_\alpha^2\}$ descends to the sheaf $\mathcal{O}b_{\mathbf{M}}$ of $\mathcal{O}_{\mathbf{M}}$ -modules and the obstruction assignments $\{\mathbf{ob}_{R_\alpha}\}$ are compatible over all $R_{\alpha\beta} = R_\alpha \times_{\mathbf{M}} R_\beta$.

We next assume $R_\alpha \rightarrow S_\alpha$ admits a perfect relative obstruction theory given by $(\mathcal{L}_\alpha^\bullet, \mathbf{ob}_{R_\alpha/S_\alpha})$ as defined in Definition 1.11. We say $\mathbf{M} \rightarrow \mathbf{N}$ admits a perfect relative obstruction theory if we can choose $\{(\mathcal{L}_\alpha^\bullet, \mathbf{ob}_{R_\alpha/S_\alpha})\}_\Lambda$ so that the relative obstruction sheaves $\mathcal{O}b_{R_\alpha/S_\alpha} \triangleq \text{Coker}\{\mathcal{L}_\alpha^1 \rightarrow \mathcal{L}_\alpha^2\}$ descends to a global sheaf on $\mathbf{M}$ and the obstruction assignments $\mathbf{ob}_{R_\alpha/S_\alpha}$ are compatible on the overlaps $R_{\alpha\beta}$.

Definition 4.1. *The perfect (relative) obstruction theories $\{\mathcal{E}_\alpha^\bullet, \mathbf{ob}_{R_\alpha}\}$, $\{\mathcal{F}_\alpha^\bullet, \mathbf{ob}_{S_\alpha}\}$ and $\{\mathcal{L}_\alpha^\bullet, \mathbf{ob}_{R_\alpha/S_\alpha}\}$ are said to be compatible if to each $\alpha \in \Lambda$ there is an exact triangle of complexes*

$$(4.1) \quad \implies \mathcal{L}_\alpha^\bullet \implies \mathcal{E}_\alpha^\bullet \implies \mathcal{F}_\alpha^\bullet \otimes_{\mathcal{O}_{S_\alpha}} \mathcal{O}_{R_\alpha} \implies \mathcal{L}_\alpha^{\bullet+1} \implies$$

which induces a long exact sequence of cohomologies

$$(4.2) \quad \rightarrow h^i(\mathcal{L}_\alpha^\bullet \otimes \mathcal{I}) \xrightarrow{\tau_{1,i}} h^i(\mathcal{E}_\alpha^\bullet \otimes \mathcal{I}) \xrightarrow{\tau_{2,i}} h^i(\mathcal{F}_\alpha^\bullet \otimes_{\mathcal{O}_{S_\alpha}} \mathcal{I}) \xrightarrow{\tau_{0,i+1}} h^{i+1}(\mathcal{L}_\alpha^\bullet \otimes \mathcal{I}) \rightarrow$$

for any sheaf of $\mathcal{O}_{R_\alpha}$ -modules $\mathcal{I}$ that satisfies the following properties:

(1). *The first part of the above exact sequence is identical to the exact sequence*

$$0 \longrightarrow \mathfrak{Def}_{R_\alpha/S_\alpha}^1(\mathcal{I}) \longrightarrow \mathfrak{Def}_{R_\alpha}^1(\mathcal{I}) \longrightarrow \mathfrak{Def}_{S_\alpha}^1(\mathcal{I}) \longrightarrow$$

under the canonical isomorphisms $\mathfrak{Def}^1 = h^1(\cdot \otimes \mathcal{I})$ given by the definition of the perfect obstruction theories.

(2). *Let ξ be any object in $\mathfrak{Tri}_{R_\alpha}$, which is also an object in $\mathfrak{Tri}_{S_\alpha}$. Then under*

¹⁶Namely, the open covering are universal open smooth morphisms.

the arrow $\tau_{2,2}$ in (4.2) we have $\mathbf{ob}_{S_\alpha}(\xi) = \tau_{2,2}(\mathbf{ob}_{R_\alpha}(\xi))$.

(3). Let $\xi = (B, I, \varphi_0)$ be an object in $\mathfrak{Tri}_{R_\alpha/S_\alpha}$, which is also an object in $\mathfrak{Tri}_{R_\alpha}$. Then $\mathbf{ob}_{R_\alpha}(\xi) = \tau_{2,1}(\mathbf{ob}_{R_\alpha/S_\alpha}(\xi))$. Further, suppose $\mathbf{ob}_{R_\alpha}(\xi) = 0$ and $\varphi_0: \text{Spec } B/I \rightarrow R_\alpha$ extends to $\varphi: \text{Spec } B \rightarrow R_\alpha$. Let $e \in h^1(\mathcal{F}_\alpha^\bullet \otimes_{B/I} I)$ be the difference of the tautological $\text{Spec } B \rightarrow \text{Spec } S_\alpha$ and the composite of $\text{Spec } B \xrightarrow{\varphi} \text{Spec } R_\alpha$ and $\text{Spec } R_\alpha \xrightarrow{pr} \text{Spec } S_\alpha$. Then $\mathbf{ob}_{R_\alpha/S_\alpha}(\xi) = \tau_{0,2}(e)$.

Finally, the collection of exact sequences $h^2(\mathcal{L}_\alpha^\bullet) \rightarrow h^2(\mathcal{E}_\alpha^\bullet) \rightarrow h^2(\mathcal{F}_\alpha^\bullet) \rightarrow 0$ descends to an exact sequence of sheaves

$$\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \longrightarrow \mathcal{O}b_{\mathbf{M}} \longrightarrow \mathcal{O}b_{\mathbf{N}} \otimes_{\mathcal{O}_{\mathbf{N}}} \mathcal{O}_{\mathbf{M}} \longrightarrow 0.$$

The goal of this subsection is to show that with the data given in Definition 4.1, we can construct a class $[\mathbf{M}, \mathbf{N}]^{\text{virt}} \in A_*\mathbf{M}$, called the relative virtual moduli cycle. We will then show that it is equal to $[\mathbf{M}]^{\text{virt}}$ in $A_*\mathbf{M}$. This allows us to give a different interpretation of $[\mathbf{M}]^{\text{virt}}$, useful in the proof of the key lemmas in the previous section.

We first study the local situation. We let $p \in \mathbf{M}$ be any closed point, $q \in \mathbf{N}$ be the image of p , $\bar{p} \in R_\alpha$ be a lift of $p \in \mathbf{M}$ and $\bar{q} \in S_\alpha$ be the image of $\bar{p}$. We let $T_1 = h^1(\mathcal{E}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$, $T_2 = h^1(\mathcal{F}_\alpha^\bullet \otimes \mathbb{k}_{\bar{q}})$ and let $T_{1/2} = h^1(\mathcal{L}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$. Similarly, we let $O_1 = h^2(\mathcal{E}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$, $O_2 = h^2(\mathcal{F}_\alpha^\bullet \otimes \mathbb{k}_{\bar{q}})$ and let $O_{1/2} = h^2(\mathcal{L}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$, all implicitly depending on the lift $\bar{p}$. Note that they fit into the exact sequence

$$(4.3) \quad 0 \longrightarrow T_{1/2} \longrightarrow T_1 \longrightarrow T_2 \xrightarrow{\delta} O_{1/2} \longrightarrow O_1 \longrightarrow O_2 \longrightarrow 0$$

induced by (4.1). We now let $T = T_1 \oplus T_2 / \ker(\delta)$ and pick a surjective homomorphism $T \rightarrow T_2$ that is an extension of $T_1 \rightarrow T_2$. Next we let $O = O_1 \oplus \text{Im}(\delta)$. We then pick an injective homomorphism $\eta: O_{1/2} \rightarrow O$ so that the composite $O_{1/2} \rightarrow O \rightarrow O_{1/2}$ is the identity map. This way we have two exact sequences

$$(4.4) \quad 0 \rightarrow T_{1/2} \rightarrow T \rightarrow T_2 \rightarrow 0 \quad \text{and} \quad 0 \rightarrow O_{1/2} \rightarrow O \rightarrow O_2 \rightarrow 0.$$

Further, with various homomorphisms chosen and the isomorphism $T_2 / \ker(\delta) \cong \text{Im}(\delta)$, we have the following induced exact sequence

$$0 \longrightarrow T_1 \longrightarrow T \longrightarrow O \longrightarrow O_1 \longrightarrow 0.$$

In the following for any vector space V we denote by $\mathbb{k}[[V]]$ the ring of formal power series $\lim_n \oplus_{i=0}^n S^i(V)$. We now let

$$(4.5) \quad f \in \mathbb{k}[[T^\vee]] \otimes O, \quad g \in \mathbb{k}[[T_2^\vee]] \otimes O_2 \quad \text{and} \quad h \in \mathbb{k}[[T^\vee]] / (g) \otimes O_{1/2}$$

be the Kuranishi maps of the (relative) obstruction theories of R_α , of S_α and of R_α/S_α at $\bar{p}$ (or $\bar{q}$) respectively (see [LT2, Lemma 3.10]). By abuse of notation, we denote by (g) the idea generated by the components of g in $\mathbb{k}[[T_2^\vee]]$ and in $\mathbb{k}[[T^\vee]]$ via the inclusion $\mathbb{k}[[T_2^\vee]] \rightarrow \mathbb{k}[[T^\vee]]$.

Lemma 4.2. *We can choose the Kuranishi maps f , g and h so that*

- (1) $\varphi_1(f_p) = \tau(g_p)$ under the naturally induced maps $\varphi_1: \mathbb{k}[[T^\vee]] \otimes O \rightarrow \mathbb{k}[[T^\vee]] \otimes O_2$ and $\tau: \mathbb{k}[[T_2^\vee]] \otimes O_2 \rightarrow \mathbb{k}[[T^\vee]] \otimes O_2$;
- (2) The differential $dh_p(0): T \rightarrow O_{1/2}$ is identical to the composite $T \rightarrow T_2 / \ker(\delta) \rightarrow O_{1/2}$ induced by δ in (4.3).
- (3) Let $\tilde{h}$ and $\tilde{f}$ in $\mathbb{k}[[T_{1/2}^\vee]] / (g) \otimes O$ be the images of h and f under the obvious maps induced by the arrows mentioned before, then $\tilde{f} = \tilde{h}$.

Proof. The proof follows from the construction of Kuranishi maps, as was demonstrated in [LT2, Lemma 3.10]. $\square$

We now let $X = \operatorname{Spec} \mathbb{k}[[T^\vee]]$, $V_{1/2} = O_{1/2} \times X$, $V = O \times X$ and let $V_2 = O_2 \times X$, all viewed as vector bundles (or their total spaces) over X . We let $\rho_1 : V_{1/2} \rightarrow V$ and $\rho_2 : V \rightarrow V_2$ be the vector bundle homomorphisms induced by arrows in (4.4). Then V_2 is the quotient vector bundle of V by $V_{1/2}$. We consider the subscheme $\Gamma \subset V_1$ that is the graph of f , consider $Y = V_{1/2} \times_X \Gamma \subset V_{1/2} \times_X V$ and consider

$$\Theta_1 = \tilde{V}_{1/2} \times_{V_{1/2} \times_X V} Y \subset Y, \quad \Theta_2 = \tilde{V}_1 \times_{V_{1/2} \times_X V} Y \subset Y.$$

Here $\tilde{V}_{1/2}$ is the image scheme of the immersion $1_{V_{1/2}} \times \rho_1 : V_{1/2} \rightarrow V_{1/2} \times_X V$ and $\tilde{V} \triangleq 0_{V_{1/2}} \times V \subset V_{1/2} \times_X V$, where $0_{V_{1/2}}$ is the zero section of $V_{1/2}$. Note that $\tilde{V}_{1/2}$ and $\tilde{V}$ are isomorphic to $V_{1/2}$ and V , respectively. Following [LT2, Page 145], we denote the normal cone to $C_{\Theta_2/Y} \times_Y \Theta_1$ in $C_{\Theta_2/Y}$ ¹⁷ by $\mathcal{B}(p)_1$ and denote the normal cone to $C_{\Theta_1/Y} \times_Y \Theta_2$ in $C_{\Theta_1/Y}$ by $\mathcal{B}(p)_2$. Both $\mathcal{B}(p)_1$ and $\mathcal{B}(p)_2$ are subcones in $\hat{V}_{1/2} \times_{\hat{X}} \hat{V}$, where $\hat{X} = \operatorname{Spec} \mathbb{k}[[T_1^\vee]]/(f)$ and $\hat{V}_i = V_i \times_X \hat{X}$. As argued in [LT2, 146], based on the work of [Vis] (see also the recent [Kr1]) there is a canonical rational equivalence¹⁸ $\mathcal{Q}(p) \in W_*(\hat{V}_{1/2} \times_{\hat{X}} \hat{V})$ so that

$$(4.6) \quad \partial_0 \mathcal{Q}(p) = \mathcal{B}(p)_1 \quad \text{and} \quad \partial_\infty \mathcal{Q}(p) = \mathcal{B}(p)_2.$$

The cones $\mathcal{B}(p)_1$ and $\mathcal{B}(p)_2$ have the following interpretations as shown in [LT2, page 145]. Let $\mathcal{D}(p)_1 \subset \hat{V}$ be the normal cone to $\hat{X}$ in X , then

$$(4.7) \quad \mathcal{B}(p)_1 = \phi_1^* \mathcal{D}(p)_1$$

where $\phi_1 : \hat{V}_{1/2} \times_{\hat{X}} \hat{V} \rightarrow \hat{V}$ is the projection. Next, we let $W = \operatorname{Spec} \mathbb{k}[[T_2^\vee]]$ and let $\hat{W} = \operatorname{Spec} \mathbb{k}[[T_2^\vee]]/(g)$. We then form the normal cone to $C_{\hat{W}/W} \times_W \hat{X}$ in $C_{\hat{W}/W} \times_W X$, denoted by $\mathcal{D}(p)_2$. It is naturally a subcone in $\hat{V}_{1/2} \times_{\hat{X}} \hat{V}_2$. Then

$$(4.8) \quad \mathcal{B}(p)_2 = \phi_2^* \mathcal{D}(p)_2,$$

where $\phi_2 = (1, \rho_2) : \hat{V}_{1/2} \times_{\hat{X}} \hat{V} \rightarrow \hat{V}_{1/2} \times_{\hat{X}} \hat{V}_2$.

We caution that all the objects so far constructed depends on the lift $\bar{p}$, on the choices of arrows before (4.4) and on the Kuranishi maps. Later we will show that they are canonical in certain degree, up to the symmetry $\operatorname{Aut}(p)$.

In the following, we first construct the relative moduli cycle $[\mathbf{M}, \mathbf{N}]^{\text{virt}}$ and several related cycles based on the collection of the infinitesimal models $\mathcal{D}(p)_2$, etc. First of all, Lemma 2.7 holds for the collection of cones $\{\mathcal{D}(p)_2\}_{p \in \mathbf{M}}$. Namely at each $p \in \mathbf{M}$ the cone

$$\mathcal{D}(p)_2 \times_{\hat{X}} 0 \subset (\hat{V}_{1/2} \times_{\hat{X}} \hat{V}_2) \times_X 0$$

is independent of the choices of the Kuranishi maps f , g and h and is invariant under the natural $\operatorname{Aut}(p)$ action. (Here $\hat{X}$ is the subscheme defined before that depends on p implicitly.) Similar statements hold for the cycles $\mathcal{D}(p)_1 \in Z_* \hat{V}$, the cycles $\mathcal{B}(p)_i \in Z_*(\hat{V}_{1/2} \times_{\hat{X}} \hat{V})$ and the rational equivalence $\mathcal{Q}(p) \in W_*(\hat{V}_{1/2} \times_{\hat{X}} \hat{V})$. These were proved in [LT2, Section 3]. Secondly, for each α we pick a pair of

¹⁷For closed subscheme $A \subset B$, $C_{A/B}$ is the normal cone to A in B .

¹⁸In this paper we use the convention that a rational equivalence $Q \in W_* Z$ is a cycle in $Z_*(Z \times \mathbf{P}^1)$ so that all its irreducible components are flat over $\mathbf{P}^1$. We then define $\partial_0 Q$ and $\partial_\infty Q$ to be $Q \cap Z \times \{0\}$ and $Q \cap Z \times \{\infty\}$ respectively. In case $\varphi : Z \rightarrow W$ is a flat morphism, we denote by $\varphi^* Q$ the flat pull back of the rational equivalence.

locally free subsheaves¹⁹ $\mathcal{L}_\alpha \subset \mathcal{E}_\alpha$ over R_α , viewed as a complex $[\mathcal{L}_\alpha \rightarrow \mathcal{E}_\alpha]$, and a surjective homomorphism of complexes

$$(4.9) \quad [\mathcal{L}_\alpha \rightarrow \mathcal{E}_\alpha] \implies [\mathcal{O}b_{R_\alpha/S_\alpha} \rightarrow \mathcal{O}b_{R_\alpha}].$$

For simplicity, we denote by W_α the vector bundle $\text{Vect}(\mathcal{L}_\alpha \oplus \mathcal{E}_\alpha)$. As shown in [LT2, Section 3] there is a unique cycle $B_{i,\alpha} \in Z_* W_\alpha$ so that the collection $\mathcal{B}_i = \{(R_\alpha, B_{i,\alpha}, \mathcal{L}_\alpha \oplus \mathcal{E}_\alpha)\}_\Lambda$ satisfies the cycle consistency criteria for the infinitesimal models $\mathcal{B}(p)_i$ over the pair $(\mathbf{M}, \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \oplus \mathcal{O}b_{\mathbf{M}})$. We let $[\mathcal{B}_1]$ and $[\mathcal{B}_2]$ be the associated class in $A_* \mathbf{M}$ following the basic construction. Next we let $\mathcal{F}_\alpha \triangleq \mathcal{E}_\alpha / \mathcal{L}_\alpha$ be the quotient sheaf, which is locally free by our choice. Then $\mathcal{O}b_{S_\alpha} \otimes_{\mathcal{O}S_\alpha} \mathcal{O}R_\alpha$ is canonically a quotient sheaf of $\mathcal{F}_\alpha$. Again following [LT2, Section 3] we can find a unique cone cycle $\mathcal{D}_{2,\alpha} \subset \text{Vect}(\mathcal{L}_\alpha \oplus \mathcal{F}_\alpha)$ so that the collection $\mathcal{D}_2 = \{(R_\alpha, \mathcal{D}_{2,\alpha}, \mathcal{L}_\alpha \oplus \mathcal{F}_\alpha)\}$ satisfies the cycle consistency criteria with the infinitesimal models $\mathcal{D}(p)_2 \subset \hat{V}_{1/2} \times_{\hat{X}} \hat{V}_2$ over the pair $(\mathbf{M}, \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \oplus \rho^* \mathcal{O}b_{\mathbf{N}})$, where $\rho : \mathbf{M} \rightarrow \mathbf{N}$ is the projection. Thus by applying the basic construction to this collection we obtain the class $[\mathcal{D}_2] \in A_* \mathbf{M}$. We will call the class $[\mathcal{D}_2]$ the relative cycle and denoted it by $[\mathbf{M}, \mathbf{N}]^{\text{virt}}$.

Lemma 4.3. *We have $[\mathbf{M}]^{\text{virt}} = [\mathbf{M}, \mathbf{N}]^{\text{virt}}$ in $A_* \mathbf{M}$.*

To prove the Lemma we need to construct a rational equivalence $[\mathcal{Q}] \in W_* \mathbf{M}$ (or equivalently a class $[\mathcal{Q}] \in A_* \mathbf{M} \times \mathbf{P}^1$) so that $\partial[\mathcal{Q}]$ provides the identity in the Lemma. Here when $[\mathcal{Q}]$ is a class, we define $\partial[\mathcal{Q}] = \partial_0[\mathcal{Q}] - \partial_\infty[\mathcal{Q}]$ with $\partial_0[\mathcal{Q}]$ the image of the Gysin map $0^![\mathcal{Q}]$ associated to $0 \rightarrow \mathbf{P}^1$. The ∂_∞ is defined similarly. The proof is parallel to that of [LT2, Lemma 3.4] and will occupy the rest of this subsection. We need to provide a revised proof since the current construction does not rely on the existence of a global vector bundle as was assumed in [LT2].

We first set up the notation relevant to the construction of the cycles $[\mathcal{B}_i]$, etc, following the basic construction of the virtual cycles. First, in constructing the class $[\mathcal{B}_i]$ we first built the index set $\Xi(\mathcal{B}_i)$. Each $a \in \Xi(\mathcal{B}_i)$ associates to an integral closed substack $\mathbf{Y}_a \subset \mathbf{M}$, a multiplicity m_a , a maximal open substack $j_a : \mathbf{Y}_a^0 \rightarrow \mathbf{Y}_a$ so that $j_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \oplus j_a^* \mathcal{O}b_{\mathbf{M}}$ is locally free. Over $\mathbf{Y}_a^0$ we have a canonical cone representative $\mathbf{N}_a^0 \subset \mathbf{F}_a^0$, where $\mathbf{F}_a^0$ is the vector bundle (stack) $\text{Vect}(j_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \oplus j_a^* \mathcal{O}b_{\mathbf{M}})$ over $\mathbf{Y}_a^0$. We then pick a projective variety $\varphi_a : Y_a \rightarrow \mathbf{Y}_a$ generically finite over $\mathbf{Y}_a$, a locally free sheaf $\mathcal{F}_a$ of $\mathcal{O}_{Y_a}$ -modules and a quotient sheaf homomorphism $\mathcal{F}_a \rightarrow \varphi_a^*(\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \oplus \mathcal{O}b_{\mathbf{M}})$. We form the vector bundle $F_a = \text{Vect}(\mathcal{F}_a)$ with the induced flat morphism $F_a|_{\varphi_a^{-1}(\mathbf{Y}_a^0)} \rightarrow \mathbf{F}_a^0$. We let $N_a \subset F_a$ be the closure of the flat pull back of $\mathbf{N}_a^0$. Then $[\mathcal{B}_i]$ is the sum of $m_a \xi(a)$ over all $a \in \Xi(\mathcal{B}_i)$, where $\xi(a)$ is $\deg(\varphi_a)^{-1} \varphi_{a*} 0_{F_a}^! [N_a]$.

We set up the notation for the cycle $[\mathcal{D}_i]$ according to the same rule. We first prove $[\mathcal{B}_1] = [\mathbf{M}]^{\text{virt}}$. This identity follows from the relation (4.7). The actual proof goes as follows: First the class $[\mathcal{D}_1] = [\mathbf{M}]^{\text{virt}}$ (see [LT2, Section 3]). From (4.7) the index sets $\Xi(\mathcal{B}_1) \equiv \Xi(\mathcal{D}_1)$ naturally. For $a \in \Xi(\mathcal{B}_1)$ with $\bar{a} \in \Xi(\mathcal{D}_1)$ the corresponding element, their base substacks $\mathbf{Y}_a = \mathbf{Y}_{\bar{a}}$. Further the intrinsic representative $\mathbf{N}_a^0$ is the flat pull back of $\mathbf{N}_{\bar{a}}^0$ under an obvious homomorphism of vector bundles. Based on these, we can choose $Y_a = Y_{\bar{a}}$, choose locally free sheaves $\mathcal{F}_a = \mathcal{F}_{\bar{a}}$ and then the cone representatives $N_a \equiv N_{\bar{a}}$. This proves $\xi(a) = \xi(\bar{a})$ and

¹⁹In this paper by a pair of locally free sheaves we mean a locally free subsheaf of a locally free sheaf with locally free quotient sheaf, all of finite ranks.

hence $[\mathcal{B}_1] = [\mathcal{D}_1] = [\mathbf{M}]^{\text{virt}}$. Since the proof is straightforward, we will omit the details here. For the same reason, we prove $[\mathcal{B}_2] = [\mathbf{M}, \mathbf{N}]^{\text{virt}}$ based on the identity (4.8). Again we will omit the details here.

We now construct the required cycle $[\mathcal{Q}]$ so that

$$(4.10) \quad \partial_0[\mathcal{Q}] = [\mathcal{B}_1] \quad \text{and} \quad \partial_\infty[\mathcal{Q}] = [\mathcal{B}_2].$$

For each $\alpha \in \Lambda$ we fix the pair $\mathcal{L}_\alpha \subset \mathcal{E}_\alpha$ over R_α as in (4.9). As before we let $W_\alpha = \text{Vect}(\mathcal{L}_\alpha \oplus \mathcal{E}_\alpha)$. We claim that we can find a collection of rational equivalence $Q_\alpha \subset W_* W_\alpha$ indexed by $\alpha \in \Lambda$ that satisfy the following existence Lemma.

Lemma 4.4. *To each $\alpha \in \Lambda$ there is a unique rational equivalence $Q_\alpha \in W_* W_\alpha$ of which the following holds. Let $p \in \mathbf{M}$ be any point and let $\bar{p}$ be a lift of p in some chart R_β . We let $\hat{V}_{1/2}$, $\hat{V}$ and $\hat{V}_2$ be the vector bundles over $\hat{X}$ (associated to $\bar{p}$) defined before and after Lemma 4.2. Let R_α be the chart, let $R_{\alpha,p} = R_\alpha \times_{\mathbf{M}} p$ and let $\hat{R}_\alpha$ be the formal completion of R_α along $R_{\alpha,p}$. We also pick a morphism $\hat{R}_\alpha \rightarrow \hat{X}$ that commutes with $\hat{R}_\alpha \rightarrow \mathbf{M}$ and $\hat{X} \rightarrow \mathbf{M}$. Now let $O_{1/2} \hookrightarrow O$ be the pair in (4.4) associated to $\bar{p}$. Then up to $\text{Aut}(p)$ there are canonical induced homomorphisms of vector bundles over $R_{\alpha,p}$*

$$O_{1/2} \times R_{\alpha,p} \rightarrow h^2(\mathcal{L}_\alpha)|_{R_{\alpha,p}}, \quad O \times R_{\alpha,p} \rightarrow h^2(\mathcal{E}_\alpha)|_{R_{\alpha,p}} \quad \text{and} \quad \mathcal{L}_\alpha|_{R_{\alpha,p}} \rightarrow O_{1/2} \times R_{\alpha,p}.$$

We then pick a surjective homomorphism $\varphi_2 : \mathcal{E}_\alpha|_{R_{\alpha,p}} \rightarrow O \times R_{\alpha,p}$ so that the following diagram of the complexes of vector bundles over $R_{\alpha,p}$

$$(4.11) \quad \begin{array}{ccc} [\mathcal{L}_\alpha|_{R_{\alpha,p}} \rightarrow \mathcal{E}_\alpha|_{R_{\alpha,p}}] & \xRightarrow{\quad} & [O_{1/2} \times R_{\alpha,p} \rightarrow O \times R_{\alpha,p}] \\ & \searrow & \downarrow \\ & & [h^2(\mathcal{L}_\alpha^\bullet)|_{R_{\alpha,p}} \rightarrow h^2(\mathcal{E}_\alpha^\bullet)|_{R_{\alpha,p}}] \end{array}$$

is commutative. Then there is a vector bundle homomorphism

$$\Phi_1 : W_\alpha \times_{R_\alpha} \hat{R}_\alpha \longrightarrow (\hat{V}_{1/2} \times_{\hat{X}} \hat{V}) \times_{\hat{X}} \hat{R}_\alpha$$

extending the obvious homomorphism

$$W_\alpha \times_{R_\alpha} R_{\alpha,p} \longrightarrow (V_{1/2} \times V) \times R_{\alpha,p}$$

induced by the φ_2 before (4.11) so that

$$\Phi_1^*(\psi^* \mathcal{Q}(p)) = \Phi_2^*(Q_\alpha),$$

where $\psi : (\hat{V}_{1/2} \times_{\hat{X}} \hat{V}) \times_{\hat{X}} \hat{R}_\alpha \rightarrow \hat{V}_{1/2} \times_{\hat{X}} \hat{V}$ and $\Phi_2 : W_\alpha \times_{R_\alpha} \hat{R}_\alpha \rightarrow W_\alpha$ are the obvious (flat) projections.

We will call this Lemma the *rational equivalence consistency criteria*.

This Lemma is proved in [LT2, Section 3] for the case where $\mathbf{M}$ is a quotient stack. The general case is exactly the same. We will not repeat the argument here.

We now fix the collection $\mathcal{Q} = \{(R_\alpha, Q_\alpha, \mathcal{L}_\alpha \oplus \mathcal{E}_\alpha)\}_\Lambda$. To each $\alpha \in \Lambda$ we let $\Xi(Q_\alpha)$ be the set of irreducible components of Q_α . For each $a \in \Xi(Q_\alpha)$, we let $T_a \subset Q_\alpha$ be the corresponding irreducible component, let m_a be the multiplicity of Q_α along T_a and let $\mathbf{Y}_a \subset \mathbf{M}$ be the base substack of a , which is the closure of the image of $T_a \rightarrow \mathbf{M}$. We then let $\iota_a : \mathbf{Y}_a^0 \rightarrow \mathbf{Y}_a$ be an open substack so that $\iota_a^* \mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}}$, $\iota_a^* \mathcal{O}_{b_{\mathbf{M}}}$ and $\mathcal{K}_a^0 = \ker\{\iota_a^* \mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}} \rightarrow \iota_a^* \mathcal{O}_{b_{\mathbf{M}}}\}$ are locally free. We then pick a homomorphism

$$(4.12) \quad \eta : \iota_a^* \mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}} \rightarrow \mathcal{K}_a^0$$

so that the composite $\mathcal{K}_a^0 \rightarrow \iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{K}_a^0$ is the identity. (Such lift exists possibly after shrinking $\mathbf{Y}_a^0 \subset \mathbf{Y}$ if necessary.) We let $\mathbf{V}_{1/2,a}^0 = \text{Vect}(\iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}})$, let $\mathbf{V}_a^0 = \text{Vect}(\mathcal{K}_a^0 \oplus \iota_a^* \mathcal{O}b_{\mathbf{M}})$ and let $\bar{\eta}: \mathbf{V}_{1/2,a}^0 \rightarrow \mathbf{V}_a^0$ be the immersion induced by η and the tautological $\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{O}b_{\mathbf{M}}$.

We now construct the intrinsic representative of a as a cycle in $\mathbf{F}_a^0 \times \mathbf{P}^1$, where $\mathbf{F}_a^0 = \mathbf{V}_{1/2,a}^0 \times_{\mathbf{Y}_a^0} \mathbf{V}_a^0$ is a vector bundle stack over $\mathbf{Y}_a^0$. Let $\text{pr}: W_\alpha \rightarrow R_\alpha$ be the projection. It is easy to see that for some dense open subset U of $\text{pr}(T_a) \subset R_a$ we can lift the tautological $U \rightarrow \mathbf{Y}_a$ to $g: U \rightarrow \mathbf{Y}_a^0$ and lift the left vertical arrow (below) to surjective horizontal arrow (the top one) as shown in the commutative diagram

$$(4.13) \quad \begin{array}{ccc} [\mathcal{L}_\alpha \rightarrow \mathcal{E}_\alpha] \otimes_{\mathcal{O}_{R_\alpha}} \mathcal{O}_U & \xrightarrow{\text{lift}} & g^*[\iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{K}_a^0 \oplus \iota_a^* \mathcal{O}b_{\mathbf{M}}] \\ \downarrow & & \downarrow \\ g^*[\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{O}b_{\mathbf{M}/\mathbf{N}}] & \longrightarrow & g^*[\iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}}]. \end{array}$$

Note that the other two arrows are tautological ones. We let

$$(4.14) \quad (W_\alpha \times \mathbf{P}^1)|_{U \times \mathbf{P}^1} \longrightarrow \mathbf{F}_a^0 \times \mathbf{P}^1$$

be the induced projection. Then an easy argument shows that the statement in Lemma 4.4 implies that there is a reduced and irreducible cycle $\mathbf{Q}_{a,\eta}^0 \subset \mathbf{F}_a^0 \times \mathbf{P}^1$ so that the restriction of T_a to fibers over $U \times \mathbf{P}^1$ is a dense open subset of the flat pull back of $\mathbf{Q}_{a,\eta}^0$ via the arrow in (4.14). We call $\mathbf{Q}_{a,\eta}^0$ an intrinsic representative of a .

Once we have constructed the intrinsic representative of $a \in \Xi(Q_\alpha)$, we can define an equivalence relation on $\cup \Xi(Q_\alpha)$ as we did before. Let $a \in \Xi(Q_\alpha)$ and $b \in \Xi(Q_\beta)$ be two elements. In case $\mathbf{Y}_a \neq \mathbf{Y}_b$, then $a \not\sim b$. In case $\mathbf{Y}_a = \mathbf{Y}_b$, we let $\mathbf{Q}_{a,\eta}^0$ and $\mathbf{Q}_{b,\eta}^0$ be the respective representatives of a and b , based on the same η in (4.12). Then we say $a \sim b$ if $\mathbf{Q}_{a,\eta}^0 \cap \mathbf{Q}_{b,\eta}^0$ is dense in both $\mathbf{Q}_{a,\eta}^0$ and $\mathbf{Q}_{b,\eta}^0$. This defines an equivalence relation. Further, whenever $a \sim b$ then $m_a = m_b$. Now let $\Xi(\mathcal{Q})$ be the set of equivalence classes. For any $a \in \Xi(\mathcal{Q})$, we pick a projective variety Y_a and a generically finite morphism $\varphi_a: Y_a \rightarrow \mathbf{Y}_a$. We then pick a pair of locally free subsheaves $\mathcal{L}_a \hookrightarrow \mathcal{E}_a$ over Y_a , a surjective homomorphism of complexes

$$[\mathcal{L}_a \rightarrow \mathcal{E}_a] \implies \varphi_a^*[\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{O}b_{\mathbf{M}}]$$

and a lift (of the above arrow)

$$[\mathcal{L}_a \rightarrow \mathcal{E}_a] \implies \varphi_a^*[\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{K}_a^0 \oplus \mathcal{O}b_{\mathbf{M}}]$$

over a dense open subset $Y_a^0 \subset Y_a$. After that, we let $W_a = \text{Vect}(\mathcal{L}_a \oplus \mathcal{E}_a)$ and let $j_a: W_a|_{Y_a^0} \rightarrow \mathbf{F}_a^0$ be the induced morphism. By shrinking Y_a^0 if necessary, we can assume that $W_a|_{Y_a^0} \rightarrow \mathbf{F}_a^0$ is flat. With all these chosen, we let Q_a be the closure in $W_a \times \mathbf{P}^1$ of the flat pull back of $\mathbf{Q}_{a,\eta}^0$ and define $\xi(a) = \deg(\varphi_a)^{-1} \varphi_{a*} 0_{W_a}^! [Q_a]$. We define

$$[\mathcal{Q}] = \sum_{a \in \Xi(\mathcal{Q})} m_a \xi(a) \in A_* \mathbf{M} \times \mathbf{P}^1.$$

It remains to prove the identity

$$(4.15) \quad \partial_\infty[\mathcal{Q}] = \sum_{a \in \Xi(\mathcal{Q})} m_a \partial_\infty \xi(a) = [\mathcal{B}_2] \in A_* \mathbf{M}$$

and the similar identity with ∂_∞ (resp. $[\mathcal{B}_2]$) replaced by ∂_0 (resp. $[\mathcal{B}_1]$). We will prove the identity (4.15). The proof of the other identity is similar. To achieve this, we will define a function $\mu : \Xi(\mathcal{Q}) \times \Xi(\mathcal{B}_2) \rightarrow \mathbb{Q}$ that satisfies

$$(4.16) \quad \sum_{a \in \Xi(\mathcal{Q})} m_a \mu(a, b) = m_b, \quad \forall b \in \Xi(\mathcal{B}_2)$$

and

$$(4.17) \quad \partial_\infty \xi(a) = \sum_{b \in \Xi(\mathcal{B}_2)} \mu(a, b) \xi(b), \quad \forall a \in \Xi(\mathcal{Q}).$$

Once these two are established, then

$$\partial_\infty[\mathcal{Q}] = \sum m_a \partial_\infty \xi(a) = \sum m_a \sum \mu(a, b) \xi(b) = \sum m_b \xi(b) = [\mathcal{B}_2],$$

which is (4.15).

We first construct the function μ . We begin with any (smooth) chart R_α . We let $\mathcal{L}_\alpha \oplus \mathcal{E}_\alpha$ be the locally free sheaves on R_α chosen before. For simplicity we denote by $\mathcal{W}_\alpha$ the sheaf $\mathcal{L}_\alpha \oplus \mathcal{E}_\alpha$ and continue to denote by W_α the vector bundle $\text{Vect}(\mathcal{L}_\alpha \oplus \mathcal{E}_\alpha)$. We let $B_{2,\alpha} \subset W_\alpha$ and $Q_\alpha \subset W_\alpha \times \mathbf{P}^1$ be the cycles constructed before. For each $a \in \Xi(\mathcal{Q})$ we let $Q_{\alpha,a}$ be the closure in $W_\alpha \times \mathbf{P}^1$ of the pull back of $\mathbf{Q}_{a,\eta}^0$ via the arrow in (4.14). The cycle $Q_{\alpha,a}$ is also the union of irreducible components of Q_α in the equivalence class a . With this choice of $Q_{\alpha,a}$, we have the decomposition

$$(4.18) \quad Q_\alpha = \sum_{a \in \Xi(\mathcal{Q})} m_a Q_{\alpha,a}.$$

Similarly, we have a canonical decomposition

$$B_{2,\alpha} = \sum_{b \in \Xi(\mathcal{B}_2)} m_b B_{2,\alpha,b}.$$

As before, we let $\Xi(B_{2,\alpha})$ be the index set of irreducible components of $B_{2,\alpha}$. For $c \in \Xi(B_{2,\alpha})$, we denote by T_c the corresponding component. Then because $\partial_\infty Q_\alpha = B_{2,\alpha}$, there is a unique function $\mu_\alpha(a, \cdot) : \Xi(B_{2,\alpha}) \rightarrow \mathbb{Z}$ so that $\partial_\infty Q_{\alpha,a} = \sum \mu_\alpha(a, c) T_c$. Again, by cycle (rational equivalence) consistency criteria and the invariance of the cycle under $\text{Aut}(p)$, we conclude that whenever $c_1 \sim c_2$ then $\mu_\alpha(a, c_1) = \mu_\alpha(a, c_2)$. Thus $\mu_\alpha(a, \cdot)$ descends to a function $\mu_\alpha(a, \cdot)$ from $\Xi(\mathcal{B}_2)$ to $\mathbb{Z}$. Finally, for $b \in \Xi(\mathcal{B}_2)$ we define $B_{2,\alpha,b}$ to be the union of T_c for those $c \in \Xi(B_{2,\alpha})$ such that $c \sim b$. Then we have the identity

$$\partial_\infty Q_{\alpha,a} = \sum_{b \in \Xi(\mathcal{B}_2)} \mu_\alpha(a, b) B_{2,\alpha,b}.$$

For the same reason whenever $R_\alpha \times_{\mathbf{M}} R_\beta \times_{\mathbf{M}} \mathbf{Y}_b \neq \emptyset$, then $\mu_\alpha(a, b) = \mu_\beta(a, b)$. Hence, we can define $\mu(a, b)$ to be $\mu_\alpha(a, b)$ for those α so that $R_\alpha \times_{\mathbf{M}} \mathbf{Y}_b \neq \emptyset$. Then the identity (4.16) follows from collecting term $B_{2,\alpha,b}$ in

$$\sum_{b \in \Xi(\mathcal{B}_2)} m_b B_{2,\alpha,b} = B_{2,\alpha} = \partial_\infty Q_\alpha = \sum_{a \in \Xi(\mathcal{Q})} m_a \sum_{b \in \Xi(\mathcal{B}_2)} \mu(a, b) B_{2,\alpha,b}$$

for some α so that $R_\alpha \times_{\mathbf{M}} \mathbf{Y}_b \neq \emptyset$.

We now investigate (4.17). Let $a \in \Xi(\mathcal{Q})$ be any element, let $\varphi_a : Y_a \rightarrow \mathbf{Y}_a$ be the generically finite morphism and let $\mathcal{L}_a \subset \mathcal{E}_a$ be the pair of sheaves chosen before. We still denote by W_a the vector bundle $\text{Vect}(\mathcal{L}_a \oplus \mathcal{E}_a)$. We let $Q_a \in W_* W_a$ be the

representative of a and let $\partial_\infty Q_a = \sum m_i D_i$ be the decomposition into irreducible components. Since $Q_a \rightarrow \mathbf{M}$ factor through $\mathbf{Y}_a \subset \mathbf{M}$, all D_i lie over $\mathbf{Y}_a$. We now let $\mathbf{A} \subset \mathbf{Y}_a$ be a closed integral substack. We let $\Upsilon_{\mathbf{A}}$ be (the index set of) those D_i so that the image $D_i \rightarrow \mathbf{M}$ is exactly $\mathbf{A}$. Similarly we let $\Xi(\mathcal{B}_2)_{\mathbf{A}}$ be those $b \in \Xi(\mathcal{B}_2)$ so that $\mathbf{Y}_b = \mathbf{A}$. Clearly if we can show that

$$(4.19) \quad \deg(\varphi_a)^{-1} \sum_{i \in \Upsilon_{\mathbf{A}}} m_i \varphi_{a*} 0_{W_a}^! [D_i] = \sum_{b \in \Xi(\mathcal{B}_2)_{\mathbf{A}}} \mu(a, b) \xi(b),$$

since $\mathbf{A} \subset \mathbf{Y}_a$ is arbitrary, (4.17) will follow immediately.

In the remainder of this section we will prove the identity (4.19) for any pair $(a, \mathbf{A})$. We pick an étale $\varphi_U : U \rightarrow \mathbf{Y}_a$ so that $U \times_{\mathbf{Y}_a} \mathbf{A} \rightarrow \mathbf{A}$ is dominant. We then pick a pair of locally free sheaves $\mathcal{L}_U \subset \mathcal{E}_U$ and surjective homomorphisms ϕ_1 and ϕ_2 of complexes as shown that make the diagram

$$\begin{array}{ccc} [\mathcal{L}_U \rightarrow \mathcal{E}_U] & \xrightarrow{\phi_1} & \varphi_U^* [\iota_a^* \mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}} \rightarrow \mathcal{K}_a^0 \oplus \iota_a^* \mathcal{O}_{b_{\mathbf{M}}}] \\ \downarrow \phi_2 & & \downarrow \\ \varphi_U^* [\mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}} \rightarrow \mathcal{O}_{b_{\mathbf{M}}}] & \longrightarrow & \varphi_U^* [\iota_a^* \mathcal{O}_{b_{\mathbf{M}/\mathbf{N}}} \rightarrow \iota_a^* \mathcal{O}_{b_{\mathbf{M}}}] \end{array}$$

commutative. Here the top-right corner is the complex in (4.13) and the right vertical arrow is the standard projection. As before, we let W_U be the vector bundle $\text{Vect}(\mathcal{L}_U \oplus \mathcal{E}_U)$. Then the homomorphism ϕ_1 defines a flat morphism $W_U|_{\varphi_U^{-1}(\mathbf{Y}_a^0)} \rightarrow \mathbf{F}_a^0$. We let $Q_U \subset W_* W_U$ be the closure of the flat pull back of $\mathbf{Q}_{a,\eta}^0 \in W_* \mathbf{F}_a^0$.

We next consider the projections

$$q_1 : U \times_{\mathbf{Y}_a} Y_a \rightarrow U \quad \text{and} \quad \tilde{q}_1 : W_U \times_{\mathbf{Y}_a} Y_a \rightarrow W_U.$$

We let $\tilde{W} \subset W_U \times_{\mathbf{Y}_a} Y_a$ be a dense open subset so that the tautological $\tilde{W} \rightarrow \mathbf{Y}_a$ factor through $\mathbf{Y}_a^0$ and $\tilde{W} \rightarrow \mathbf{F}_a^0$ is quasi-finite. We then let $\tilde{Q}_U$ be the closure in $(W_U \times_{\mathbf{Y}_a} Y_a) \times \mathbf{P}^1$ of the flat pull back of $\mathbf{Q}_{a,\eta}^0$ via the obvious $\tilde{W} \rightarrow \mathbf{F}_a^0$. Then since $\mathbf{Q}_{a,\eta}^0$ is dominant over $\mathbf{Y}_a^0$ and since $\tilde{q}_1$ is proper, we have $\tilde{q}_{1*}(\tilde{Q}_U) = \deg(\varphi_a) Q_U$ and then

$$(4.20) \quad \tilde{q}_{1*}(\partial_\infty \tilde{Q}_U) = \partial_\infty \tilde{q}_{1*}(\tilde{Q}_U) = \deg(\varphi_a) \partial_\infty Q_U.$$

For the convenience of the readers, we list the related rational equivalence relations constructed:

$$\mathbf{Q}_{a,\eta}^0 \in W_* \mathbf{F}_a^0, \quad Q_U \in W_* W_U, \quad \tilde{Q}_U \in W_*(W_U \times_{\mathbf{Y}_a} Y_a) \quad \text{and} \quad Q' \in W_* W.$$

(Q' will be constructed in (4.23).) We now let Υ (resp. $\tilde{\Upsilon}$) be the (index) set of those irreducible components of $\partial_\infty Q_U$ (resp. $\partial_\infty \tilde{Q}_U$) that lie over and dominate $\mathbf{A}$. For $i \in \Upsilon$ or $\tilde{\Upsilon}$ we denote by T_i the corresponding component and by m_i the multiplicity of Q_U or $\tilde{Q}_U$ along T_i . We need to divide $\tilde{\Upsilon}$ into two parts: One is $\tilde{\Upsilon}_1$ which consists of those T_i so that $\tilde{q}_{1*}(T_i) \neq 0$. We let $\tilde{\Upsilon}_0 = \tilde{\Upsilon} - \tilde{\Upsilon}_1$. Because of (4.20), there is a map $\lambda : \tilde{\Upsilon}_1 \rightarrow \Upsilon$ so that under $\tilde{q}_{1*}$ the component T_i is mapped onto $T_{\lambda(i)}$, say via $\psi_{i,\lambda(i)} : T_i \rightarrow T_{\lambda(i)}$. Then (4.20) implies

$$(4.21) \quad \sum_{i \in \lambda^{-1}(j)} \deg(\psi_{i,j}) = \deg(\varphi_a), \quad j \in \Upsilon.$$

We now compare the collection $\tilde{\Upsilon}$ with the collection $\Upsilon_{\mathbf{A}}$. We let p_1 and p_2 be the projections of $U \times_{\mathbf{Y}_a} Y_a$ to U and Y_a , respectively. We pick a pair of locally

free sheaves $\mathcal{L} \hookrightarrow \mathcal{E}$ on $U \times_{\mathbf{Y}_a} Y_a$ so that there are two surjective homomorphisms ϕ_1 and ϕ_2 of complexes as shown in the commutative diagram

$$(4.22) \quad \begin{array}{ccc} [\mathcal{L} \rightarrow \mathcal{E}] & \xrightarrow{\phi_1} & p_2^*[\mathcal{L}_a \rightarrow \mathcal{E}_a] \\ \downarrow \phi_2 & & \downarrow \\ p_1^*[\mathcal{L}_U \rightarrow \mathcal{E}_U] & \longrightarrow & [\iota_a^* \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{K}_a^0 \oplus \iota_a^* \mathcal{O}b_{\mathbf{M}}]. \end{array}$$

Here the two remaining arrows in the diagram are the ones chosen before. We let W be the vector bundle $\text{Vect}(\mathcal{L} \oplus \mathcal{E})$ over $U \times_{\mathbf{Y}_a} Y_a$ and let $\zeta_1: W \rightarrow W_U \times_{\mathbf{Y}_a} Y_a$ and $\zeta_2: W \rightarrow W_a$ be the morphisms to vector bundles over $U \times_{\mathbf{Y}_a} Y_a$ and Y_a induced by ϕ_1 and ϕ_2 , respectively. The map ζ_1 is obviously smooth whose fibers are vector spaces. Since $U \times_{\mathbf{Y}_a} Y_a \rightarrow Y_a$ is étale, ζ_2 is also smooth with affine fibers. Further, because of the commutative diagram (4.22) and the rational equivalence consistency criteria, the flat pull back of Q_U via ζ_1 and the flat pull back of Q_a via ζ_2 are identical over a dense open subset $\mathcal{U} \subset U \times_{\mathbf{Y}_a} Y_a$ that is flat over both U and Y_a . Hence since both Q_U and Q_a are closed and since all their irreducible components dominate $\mathbf{Y}_a$,

$$(4.23) \quad Q' \triangleq \zeta_1^*(Q_U) = \zeta_2^*(Q_a).$$

We now let Υ' be the set of those irreducible components of $\partial_\infty Q'$ that lie over and dominate $\mathbf{A}$. Since the fibers of ζ_1 are vector spaces, Υ' is naturally identical to $\tilde{\Upsilon}$, say via $\lambda': \Upsilon' \rightarrow \tilde{\Upsilon}$. Further, for each $i \in \Upsilon'$ the component T_i is a vector bundle over $T_{\lambda'(i)}$. On the other hand, since $p_2: U \times_{\mathbf{Y}_a} Y_a \rightarrow Y_a$ is smooth, each irreducible component of $\partial_\infty Q'$ is a component of the flat pull back of a T_i for some $i \in \Upsilon_{\mathbf{A}}$. Thus there is a map $\lambda_a: \Upsilon' \rightarrow \Upsilon_{\mathbf{A}}$ so that for each $i \in \Upsilon_{\mathbf{A}}$ the union $\sum_{j \in \lambda_a^{-1}(i)} T_j$ is exactly the pull back of T_i . For $i \in \Upsilon'$ and $j \in \lambda_a(i)$, we let $\varphi_{i,j}: T_i \rightarrow T_j$ be the tautological map.

We are now ready to prove the identity (4.19). For each $i \in \Upsilon'$, we let $b(T_i)$ be the image of $T_i \rightarrow U \times_{\mathbf{Y}_a} Y_a$. We pick a projective variety Z_i so that a dense open subset $Z_i^0 \subset Z_i$ is a finite branched cover of $b(T_i)$ and the induced $Z_i^0 \rightarrow \mathbf{A}$ extends to $\varphi_i: Z_i \rightarrow \mathbf{A}$. We then pick a pair of locally free sheaves $\mathcal{L}_i \subset \mathcal{E}_i$ over Z_i and a surjective homomorphism of complexes ϕ'_1 that lifts to a surjective ϕ'_2 as shown in the commutative diagram

$$\begin{array}{ccc} [\mathcal{L}_i \rightarrow \mathcal{E}_i] & \xrightarrow{\phi'_2} & [\mathcal{L}|_{Z_i^0} \rightarrow \mathcal{E}|_{Z_i^0}] \\ \downarrow \phi'_1 & & \downarrow \\ \varphi_i^*[\mathcal{O}b_{\mathbf{M}/\mathbf{N}} \rightarrow \mathcal{O}b_{\mathbf{M}}] & \longrightarrow & \varphi_i^*[\mathcal{O}b_{\mathbf{M}/\mathbf{N}}|_{Z_i^0} \rightarrow \mathcal{O}b_{\mathbf{M}}|_{Z_i^0}]. \end{array}$$

Then following the definition of ξ we let $W_i = \text{Vect}(\mathcal{L}_i \oplus \mathcal{E}_i)$ and let $C_i \subset W_i \times \mathbf{P}^1$ be the closure of the flat pull back of $T_i \subset W|_{Z_i^0} \times \mathbf{P}^1$ via the induced $W_i|_{Z_i^0} \rightarrow W|_{Z_i^0}$. We then define (the non-normalized) $\tilde{\xi}(i) = \varphi_{i*} 0_{W_i}^! [C_i] \in A_* \mathbf{M} \times \mathbf{P}^1$.

For $i \in \Upsilon$, $\tilde{\Upsilon}$ or $\Upsilon_{\mathbf{A}}$ we define the variety Z_i^0 to be the image of T_i in U , in $U \times_{\mathbf{Y}_a} Y_a$ or Y_a respectively, and then define $\varphi_i: Z_i \rightarrow \mathbf{A}$ and the class $\tilde{\xi}(i)$ along the same line. We let $\psi_{U,\mathbf{A}}$ be the induced morphism $U \times_{\mathbf{Y}_a} \mathbf{A} \rightarrow \mathbf{A}$. We have the following Lemma concerning the relations $\lambda: \tilde{\Upsilon}_1 \rightarrow \Upsilon$, $\lambda': \Upsilon' \rightarrow \tilde{\Upsilon}$ and $\lambda_a: \Upsilon' \rightarrow \Upsilon_{\mathbf{A}}$ defined before.

Lemma 4.5. *The following relations hold: 1. For each $i \in \Upsilon'$ we have $\bar{\xi}(i) = \bar{\xi}(\lambda'(i))$; 2. For each $i \in \tilde{\Upsilon}$ we have $\deg(\varphi_{i,\lambda_a(i)}) \bar{\xi}(\lambda_a(i)) = \bar{\xi}(i)$ and for any $j \in \Upsilon_{\mathbf{A}}$ we have $\sum_{i \in \lambda_a^{-1}(j)} \deg(\varphi_{i,j}) = \deg(\psi_{U,\mathbf{A}})$; 3. For any $i \in \tilde{\Upsilon}_1$ we have $\deg(\psi_{i,\lambda(i)}) \bar{\xi}(\lambda(i)) = \bar{\xi}(i)$; 4. For $i \in \tilde{\Upsilon}_0$ we have $\bar{\xi}(i) = 0$.*

Proof. The proof of (1) is parallel to the proof of Lemma 2.6, using the fact that the fibers of ζ_1 are vector spaces. We will omit the proof here. The proof of the identity in (2) concerning $\bar{\xi}(\cdot)$ is similar and will be omitted too. The identity concerning the degrees in (2) follows from the fact that $U \times_{\mathbf{Y}_a} Y_a \rightarrow Y_a$ is étale. We now prove (3). Let $i \in \tilde{\Upsilon}$ and $j = \lambda(i) \in \Upsilon$. We let $Z_i^0 \subset Z_i$ and $Z_j^0 \subset Z_j$ be the pair of varieties constructed in defining the classes $\bar{\xi}(i)$ and $\bar{\xi}(j)$. By definition, there is a canonical dominant morphism $Z_i^0 \rightarrow Z_j^0$. Hence without loss of generality we can assume that it extends to $\rho: Z_i \rightarrow Z_j$. With this choice of Z_i and Z_j , we can choose the pairs of sheaves $\mathcal{L}_i \subset \mathcal{E}_i$ and $\mathcal{L}_j \subset \mathcal{E}_j$ be so that the former is the pull back of the later via ρ . Then because of the relation (4.20), the cycle representatives $C_i \in Z_* \text{Vect}(\mathcal{L}_i \oplus \mathcal{E}_i)$ and $C_j \in Z_* \text{Vect}(\mathcal{L}_j \oplus \mathcal{E}_j)$ satisfies $\tilde{\rho}_* C_i = \deg(\psi_{i,j}) C_j$. Here we used the fact that C_i is supported on a single variety. This relation implies (3) immediately. The proof of (4) is parallel and will be omitted. This proves the Lemma. $\square$

It follows from the Lemma that

$$\deg(\psi_{U,\mathbf{A}}) \sum_{i \in \Upsilon_{\mathbf{A}}} m_i 0_{W_a}^! [D_i] = \deg(\psi_{U,\mathbf{A}}) \sum_{i \in \Upsilon_{\mathbf{A}}} m_i \bar{\xi}(i) = \sum_{j \in \Upsilon'} m_j \bar{\xi}(j).$$

Here we used (2) to derive the last identity. Because $\bar{\xi}(i) = 0$ for $i \in \tilde{\Upsilon}_0$, by (1) of the Lemma the right hand side above is

$$\begin{aligned} \sum_{i \in \tilde{\Upsilon}_1} m_i \bar{\xi}(i) &= \sum_{i \in \tilde{\Upsilon}_1} \deg(\psi_{i,\lambda(i)}) m_i \bar{\xi}(\lambda(i)) = \deg(\varphi_a) \sum_{i \in \Upsilon} m_i \bar{\xi}(i) \\ &= \deg(\varphi_a) \deg(\psi_{U,\mathbf{A}}) \sum_{b \in \Xi(\mathcal{B}_2)_{\mathbf{A}}} \mu(a,b) \xi(b). \end{aligned}$$

Here the first equality follows from (3) in the Lemma, the second equality follows from (4.21) and the last identity follows from the definition of $\mu(a,b)$, using the fact that U is étale over $\mathbf{Y}_a$. This proves the identity (4.19) and hence completes the proof of Lemma 4.3.

4.2. Three special cases. In this subsection, we will study three cases of relative obstruction theories and derive some identities.

We first study the case where $\mathbf{N}$ is a DM-stack and $\mathbf{M} \rightarrow \mathbf{N}$ is a substack defined by the vanishing of a C-divisor $(\mathbf{L}, \mathbf{s})$. Namely if $S_\alpha \rightarrow \mathbf{N}$ is a chart (in an atlas Λ) and (L_α, s_α) is the associated C-divisor on S_α , then $\mathbf{M} \times_{\mathbf{N}} S_\alpha$ is defined by the vanishing of s_α . We denote $\mathbf{M} \times_{\mathbf{N}} S_\alpha$ by R_α . We claim that there is a canonical relative obstruction theory of R_α/S_α taking values in the sheaf $\mathcal{O}_{R_\alpha}(L_\alpha)$ ($= \mathcal{O}_{R_\alpha}(L_\alpha|_{R_\alpha})$). Since R_α is a subscheme of S_α , the relative first order deformations $\mathfrak{Def}_{R_\alpha/S_\alpha}^1 = 0$. Now let $\xi = (B, I, \varphi)$ be any object in $\mathfrak{Tri}_{R_\alpha/S_\alpha}$. Since B is a $\Gamma(\mathcal{O}_{S_\alpha})$ -algebra, its associated morphism $\phi: \text{Spec } B \rightarrow S_\alpha$ is an extension of $\text{Spec } B/I \rightarrow S_\alpha$. We consider the section $s_\alpha \circ \phi$ of $\phi^* L_\alpha$. Since $s_\alpha \circ \phi|_{\text{Spec } B/I} \equiv 0 \in \Gamma(\varphi^* L_\alpha)$, $s_\alpha \circ \phi$ is an element in $\Gamma(R_\alpha, L_\alpha) \otimes I$. We define this element to be the relative obstruction class $\mathbf{ob}_{R_\alpha/S_\alpha}(\xi)$. Of course, $\mathbf{ob}_{R_\alpha/S_\alpha}(\xi) = 0$ if and only

if $\phi: \text{Spec } B \rightarrow S_\alpha$ factor through $R_\alpha \subset S_\alpha$. Since $R_\alpha \rightarrow S_\alpha$ is an immersion, φ extends to $\text{Spec } B \rightarrow R_\alpha$ as S_α -morphism if and only if ϕ already factor through $R_\alpha \subset S_\alpha$. This proves that $\mathbf{ob}_{R_\alpha/S_\alpha}$ is an obstruction assignment. Since $(\mathbf{L}, \mathbf{s})$ is a $\mathbb{C}$ -divisor on $\mathbf{N}$, this defines a relative obstruction theory of $\mathbf{M}/\mathbf{N}$.

Now assume both $\mathbf{M}$ and $\mathbf{N}$ have perfect obstruction theories provided by the data $\{\mathcal{E}_\alpha^\bullet, \mathbf{ob}_{R_\alpha}\}_\Lambda$ and $\{\mathcal{F}_\alpha^\bullet, \mathbf{ob}_{R_\alpha/S_\alpha}\}_\Lambda$. We assume further that the obstruction theories of $\mathbf{M}$ and $\mathbf{N}$ are compatible to the relative obstruction theory of $\mathbf{M}/\mathbf{N}$ in the sense of Definition 4.1 with $\mathcal{L}_\alpha^\bullet = [0 \rightarrow \mathcal{O}_{R_\alpha}(L_\alpha)]$.

Lemma 4.6. *Let the notation be as before. Then $[\mathbf{M}]^{\text{virt}} = c_1(\mathbf{L}, \mathbf{s})[\mathbf{N}]^{\text{virt}}$.*

Proof. We will follow the notation developed before and after the Lemma 4.2. Let $p \in \mathbf{N}$ be any point, S_α be an étale chart of $\mathbf{N}$ with a lift $\bar{p} \in S_\alpha$ of p . We let $T_2 = h^1(\mathcal{F}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$ and let $O_2 = h^2(\mathcal{F}_\alpha^\bullet \otimes \mathbb{k}_{\bar{p}})$. We let $\hat{W}_p$ be the formal completion of S_α along $\bar{p}$. Here we use subscript p instead $\bar{p}$ since $\hat{W}_p$ depends on p up to $\text{Aut}(p)$. We know $\hat{W}_p$ is $\text{Spec } \mathbb{k}[[T_2^\vee]]/(g)$, where g is a Kuranishi map of the obstruction theory of S_α at $\bar{p}$. We denote by C_p the normal cone to $\hat{W}_p$ in $W_p \triangleq \text{Spec } \mathbb{k}[[T_2^\vee]]$. The cone C_p is naturally embedded in $O_2 \times \hat{W}_p$. Again, the pair $C_p \subset O_2 \times \hat{W}_p$ only depends on the point $p \in \mathbf{N}$, up to the symmetry $\text{Aut}(p)$. By the construction of the virtual moduli cycles in [LT2] and in section 2.2, this collection of cone cycles $\{C_p\}_{p \in \mathbf{N}}$ can be algebraized and hence gives rise to a cycle, the virtual moduli cycle $[\mathbf{N}]^{\text{virt}}$.

Now assume $p \in \mathbf{M}$. Then $\bar{p} \in R_\alpha$ where $R_\alpha = S_\alpha \times_{\mathbf{N}} \mathbf{M}$. As in (4.3), we let $T_1 = h^1(\mathcal{E}_\alpha \times \mathbb{k}_{\bar{p}})$ and $O_1 = h^2(\mathcal{E}_\alpha \otimes \mathbb{k}_{\bar{p}})$, etc. There are two cases to consider. One is when $\delta: T_2 \rightarrow O_{1/2}$ in (4.3) is 0. In this case $T = T_1 = T_2$, $O = O_1$ and $O_2 = O_1/O_{1/2}$. Further, we can choose the relative Kuranishi map $h \in \mathbb{k}[[T^\vee]]/(g) \otimes O_{1/2}$ to be $\hat{s}_\alpha$, the pull back of s_α via $\text{Spec } \mathbb{k}[[T^\vee]]/(g) \cong \hat{W}_p \rightarrow S_\alpha$. Here we have used the canonical isomorphism $O_{1/2} = \mathcal{L}_\alpha|_{\bar{p}}$. Now let X_p and $\hat{X}_p$ be the schemes $\text{Spec } \mathbb{k}[[T^\vee]]$ and $\text{Spec } \mathbb{k}[[T^\vee]]/(f)$, where f is the Kuranishi map in (4.5). Then $X_p = W_p$ and the cone $\mathcal{D}(p)_2$ is the normal cone to $C_p \times_{\hat{W}_p} \{\hat{s}_\alpha = 0\}$ in C_p . The cone $\mathcal{D}(p)_2$ is naturally embedded in $\hat{V}_{1/2} \times_{\hat{X}_p} \hat{V}_2$. For the same reason, when δ is non-zero, thus surjective since $\dim O_{1/2} = 1$, the cone $\mathcal{D}(p)_2$ is also the normal cone to $C_p \times_{\hat{W}_p} \{\hat{s}_\alpha = 0\}$ in C_p .

We are now ready to prove the Lemma. First, from the discussion before Lemma 4.3 we know the collection $\{\mathcal{D}(p)_2\}_{p \in \mathbf{M}}$ can be algebraized. By definition, the cycle constructed based on $\{\mathcal{D}(p)_2\}$ following the basic construction in Section 2.2 is the relative virtual cycle $[\mathbf{M}, \mathbf{N}]^{\text{virt}}$. However, since $\hat{s}_\alpha$ are pull back of the section s_α , the cone $\mathcal{D}(p)_2$ is the normal cone to $C_p \times_{S_\alpha} \{s_\alpha = 0\}$ in C_p . Further, since the collection $\{(L_\alpha, s_\alpha)\}$ is the restriction of $(\mathbf{L}, \mathbf{s})$ to charts S_α , a repetition of the proof of Lemma 4.3 shows that $[\mathbf{M}, \mathbf{N}]^{\text{virt}} = c_1(\mathbf{L}, \mathbf{s})[\mathbf{N}]^{\text{virt}}$. Then combined with the identity $[\mathbf{M}]^{\text{virt}} = [\mathbf{M}, \mathbf{N}]^{\text{virt}}$ in Lemma 4.3, we have $[\mathbf{M}]^{\text{virt}} = c_1(\mathbf{L}, \mathbf{s})[\mathbf{N}]^{\text{virt}}$. This completes the proof of the Lemma. $\square$

We now investigate the second case. We let $\mathbf{N}$ be a DM-stack with a morphism $\mathbf{N} \rightarrow X$ to a scheme X . For simplicity we assume X is smooth. Let $\xi: X_0 \rightarrow X$ be a smooth subvariety and let $\mathbf{M}$ be defined by the Cartesian product

$$\mathbf{M} = \mathbf{N} \times_X X_0.$$

Then $\mathbf{M}$ is a substack of $\mathbf{N}$. Let $\mathbf{L}$ over $\mathbf{M}$ be the pull back of the normal bundle to X_0 in X . Similar to the case just studied, there is a canonical relative obstruction theory of $\mathbf{M}/\mathbf{N}$ taking values in the cohomology of the complex $\mathcal{L}^\bullet = [0 \rightarrow \mathcal{O}_{\mathbf{M}}(\mathbf{L})]$. We now assume $\mathbf{M}$ and $\mathbf{N}$ both have perfect obstruction theories and are compatible to the relative obstruction theory of $\mathbf{M}/\mathbf{N}$. By the intersection theory of DM-stacks [Vis], the Gysin map $\eta^![\mathbf{N}]^{\text{virt}} \in A_*\mathbf{M}$.

Lemma 4.7. *Let the notation be as above. Then $[\mathbf{M}, \mathbf{N}]^{\text{virt}} = \eta^![\mathbf{N}]^{\text{virt}}$.*

Proof. The proof is similar to that of the previous Lemma with slight modification. We shall omit the proof here. $\square$

We now investigate the third case. We let $\mathbf{Q}$ be a smooth Artin stack with k C-divisor $(\mathbf{L}_i, \mathbf{s}_i)$ and k positive integers n_i , $i = 1, \dots, k$. We define $\mathbf{N} \subset \mathbf{Q}$ be the substack defined by the vanishing of the sections $\mathbf{s}_1^{n_1}, \dots, \mathbf{s}_k^{n_k}$. Then if we let $\bar{\rho}_\alpha : \bar{S}_\alpha \rightarrow \mathbf{Q}$ be a smooth chart and let $S_\alpha = \bar{S}_\alpha \times_{\mathbf{Q}} \mathbf{N}$, then S_α has a natural perfect obstruction theory taking values in the complex

$$(4.24) \quad \mathcal{F}_\alpha^\bullet = [\mathcal{O}_{S_\alpha}(T\bar{S}_\alpha|_{S_\alpha}) \longrightarrow \oplus \mathcal{O}_{S_\alpha}(\rho_\alpha^* \mathbf{L}_i^{\otimes n_i})],$$

where $\rho_\alpha : S_\alpha \rightarrow \mathbf{N}$ is the tautological morphism. We now let $\mathbf{M}$ be a DM-stack with a morphism $\mathbf{M} \rightarrow \mathbf{N}$ and let $R_\alpha = S_\alpha \times_{\mathbf{N}} \mathbf{M}$. We assume $\mathbf{M}$ has a perfect obstruction theory given by $\{\mathcal{E}_\alpha^\bullet, \text{ob}_{R_\alpha}\}_\Lambda$, where Λ is the atlas $\{R_\alpha\}_\Lambda$ of $\mathbf{M}$. We also assume $\mathbf{M}/\mathbf{N}$ has a perfect relative obstruction theory given by $\{\mathcal{L}_\alpha^\bullet, \text{ob}_{R_\alpha/S_\alpha}\}$. Finally, we assume all these obstruction theories are compatible. Hence by Lemma 4.3 we have $[\mathbf{M}]^{\text{virt}} = [\mathbf{M}, \mathbf{N}]^{\text{virt}}$.

What we are interested is to compare this cycle with the virtual moduli cycle of the substack $\mathbf{M}_0 \subset \mathbf{M}$ defined by $\mathbf{M}_0 = \mathbf{M} \times_{\mathbf{N}} \mathbf{N}_0$, where $\mathbf{N}_0 \subset \mathbf{Q}$ is defined by the vanishing of the sections $\mathbf{s}_1, \dots, \mathbf{s}_k$. Note that $\mathbf{M}$ is homeomorphic to $\mathbf{M}_0$. Again we assume $\mathbf{M}_0$ has perfect obstruction theory so that it is compatible to the perfect obstruction theory of $\mathbf{N}_0$ and the perfect relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$. For simplicity, we only consider the case where $\mathbf{N}_0$ is smooth and $\text{Codim}(\mathbf{N}_0, \mathbf{Q}) = k$.

Lemma 4.8. *Suppose the relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$ is induced from that of $\mathbf{M}/\mathbf{N}$. Then $[\mathbf{M}, \mathbf{N}]^{\text{virt}} = (\prod_{i=1}^k n_i)[\mathbf{M}_0, \mathbf{N}_0]^{\text{virt}}$.*

Here by the relative obstruction theory of $\mathbf{M}/\mathbf{N}$ inducing a relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$ we mean that the relative obstruction sheaf $\mathcal{O}b_{\mathbf{M}_0/\mathbf{N}_0} = \mathcal{O}b_{\mathbf{M}/\mathbf{N}} \otimes_{\mathcal{O}_{\mathbf{M}}} \mathcal{O}_{\mathbf{M}_0}$ and the relative obstruction class assignment $\text{ob}_{\mathbf{M}_0/\mathbf{N}_0}(\xi) = \text{ob}_{\mathbf{M}/\mathbf{N}}(\xi)$ for any triple $\xi \in \text{Ob}(\mathfrak{T}\mathbf{ri}_{\mathbf{M}_0/\mathbf{N}_0})$.

Proof. Without loss of generality, we can assume all $n_i \geq 2$ since otherwise we can replace $\mathbf{Q}$ by $\mathbf{Q} \cap \{s_i = 0 | n_i = 1\}$. Let $\bar{S}_\alpha \rightarrow \mathbf{Q}$, $S_\alpha = \bar{S}_\alpha \times_{\mathbf{Q}} \mathbf{N}$ and let $R_\alpha = S_\alpha \times_{\mathbf{N}} \mathbf{M}$ be as before. We let $(L_{i,\alpha}, s_{i,\alpha})$ be the restriction of $(\mathbf{L}_i, \mathbf{s}_i)$ to $\bar{S}_\alpha$. Then $S_\alpha = \bar{S}_\alpha \cap \{s_{i,\alpha}^{n_1} = \dots = s_{i,\alpha}^{n_k} = 0\}$. We let $\mathcal{L}_\alpha^\bullet$ and $\mathcal{E}_\alpha^\bullet$ be the complexes of sheaves over R_α that are part of the (relative) obstruction theories of $\mathbf{M}/\mathbf{N}$ and $\mathbf{M}$ as stated in Definition 4.1. As before, we let $\mathcal{F}_\alpha^\bullet$ be the complex (4.24). Again, we assume that the exact sequences (4.1) hold. We now let $\bar{p} \in R_\alpha$ be any closed point and let $\bar{q} \in S_\alpha$ be the image of $\bar{p}$. We let T_i and O_i ($i = 1/2, 1, 2$ or $\emptyset$) be the vector spaces defined before (4.3) associated to $\bar{p}$ and the pair R_α/S_α . Since all $n_i \geq 2$, the vector space O_2 is $\oplus_{i=1}^k L_{i,\alpha}|_{\bar{p}}$. Also we can choose the Kuranishi map $g_p \in \mathbb{k}[[T_2^\vee]] \otimes O_2$ to be the germ of $s_\alpha^{[n]} = (s_{1,\alpha}^{n_1}, \dots, s_{k,\alpha}^{n_k})$. We let $W = \text{Spec } \mathbb{k}[[T_2]]$ and let $\hat{W}$, X and $\hat{X}$ be the schemes defined after Lemma 4.2. Then since $\text{Codim}(S_\alpha, \bar{S}_\alpha) = k$ and since

$\bar{S}_\alpha$ is smooth, the cone $C_{\hat{W}/W}$ (which is defined after (4.7)) is the vector bundle $O_2 \times \hat{W}$ over $\hat{W}$. In this case the corresponding germ $\mathcal{D}(p)_2$, which is the normal cone to $C_{\hat{W}/W} \times_W \hat{X}$ in $C_{\hat{W}/W} \times_W X$, is $C_{\hat{W}/W} \times_{\hat{W}} C_{\hat{X} \times_W \hat{W}/X \times_W \hat{W}}$.

Now we consider the parallel situation for the pair $\mathbf{M}_0 \rightarrow \mathbf{N}_0$. We let $S_{0,\alpha} = \bar{S}_\alpha \times_{\mathbf{Q}} \mathbf{N}_0$ and let $R_{0,\alpha} = R_\alpha \times_{\mathbf{Q}} \mathbf{N}_0$. We will denote by $\bar{p} \in R_{0,\alpha}$ and $\bar{q} \in S_{0,\alpha}$ the same points $\bar{p} \in R_\alpha$ and $\bar{q} \in S_\alpha$, via the inclusion $R_{0,\alpha} \subset R_\alpha$ and $S_{0,\alpha} \subset S_\alpha$. We then let $T_{0,i}$ and $O_{0,i}$ be the vector spaces defined before (4.3) associated to the pair $R_{0,\alpha}/S_{0,\alpha}$ over the point $\bar{p}$. By assumption, $T_{0,2} \subset T_2$, $T_{0,1/2} \equiv T_{1/2}$, $T_0 \subset T$ is a codimension k linear subspace, $O_{0,2} = 0$ and $O_{0,1/2} = O_{1/2}$. Thus we have the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & O_{1/2} & \longrightarrow & O & \longrightarrow & O_2 \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & O_{0,1/2} & \longrightarrow & O_0 & \longrightarrow & 0 \end{array}$$

Further, we can choose the residue of $h \in \mathbb{k}[[T^\vee]] \otimes O_{1/2}$ in $\mathbb{k}[[T_0^\vee]] \otimes O_{1/2} \equiv \mathbb{k}[[T_{0,1/2}^\vee]] \otimes O_{0,1/2}$, denoted by h_0 , be the relative Kuranishi map of $R_{0,\alpha}/S_{0,\alpha}$ at $\bar{p}$. We let $\hat{W}_0$, $\hat{W}_0$, X_0 and $\hat{X}_0$ be the similarly defined formal schemes associated to $\bar{q} \in S_{0,\alpha}$ and $\bar{p} \in R_{0,\alpha}$. Note that with this choice, the Kuranishi map $g_0 = 0$ and $\hat{W}_0 = W_0$. Hence the associated germ $\mathcal{D}(p)_{0,2}$ is the normal cone $C_{\hat{X}_0/X_0}$, which is a cycle in $O_{0,1/2} \times \hat{X}_0$.

Since $\hat{W}_0$ is $\hat{W}$ with the reduced scheme structure and since the relative Kuranishi map h_0 is the restriction of the Kuranishi map h to $\hat{W}_0$, $Z_0 = Z \times_W W_0$ and $\hat{Z}_0 = \hat{Z} \times_{\hat{W}} W_0$. From this we see that the cycle $\mathcal{D}(p)_2$ is a multiple of the pull-back of $\mathcal{D}(p)_{0,2}$ under the projection

$$(O_{1/2} \times O_2) \times \hat{Z} \xrightarrow{\text{proj}} O_{0,1/2} \times \hat{Z}$$

with the multiplier given by the multiplicity of $\hat{W}$ along $\hat{W}_0$, which is $\prod_{i=1}^k n_i$.

Based on this, we see that $\Xi(\mathbf{M}/\mathbf{N})$ is canonically isomorphic to $\Xi(\mathbf{M}_0/\mathbf{N}_0)$. Further, for any $a \in \Xi(\mathbf{M}/\mathbf{N})$ with the corresponding $\bar{a} \in \Xi(\mathbf{M}_0/\mathbf{N}_0)$, a representative (A_a, F_a, φ_a) of a is also a representative of $\bar{a}$. Of course their multiplicities obey $m_a = m_{\bar{a}} \prod n_i$. Therefore

$$[\mathbf{M}/\mathbf{N}]^{\text{virt}} = \sum_{a \in \Xi(\mathbf{M}/\mathbf{N})} m_a \xi(a) = \sum_{a \in \Xi(\mathbf{M}_0/\mathbf{N}_0)} m_a \prod_{i=1}^k n_i \cdot \xi(a) = \left(\prod_{i=1}^k n_i \right) [\mathbf{M}_0/\mathbf{N}_0]^{\text{virt}}.$$

This proves the Lemma. $\square$

4.3. Proof of Lemma 3.10 and 3.11. The proof of Lemma 3.10 is similar to that of Lemma 3.11 while technically less involved. Hence we will prove Lemma 3.11 and omit the other. The strategy to prove Lemma 3.11 is to apply Lemma 4.6 to the case where $\mathbf{M} = \mathfrak{M}(\mathfrak{W}_0, \eta)$ and $\mathbf{N} = \mathfrak{M}(\mathfrak{W}, \Gamma)$. To this end, we need to work out the relative obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)/\mathfrak{M}(\mathfrak{W}, \Gamma)$.

Following the argument in Section 2.1, we only need to look at the relative obstruction theory of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}/\mathfrak{M}(W[n], \Gamma)^{\text{st}}$. We first cover $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ by affine étale charts $S_\alpha \rightarrow \mathfrak{M}(W[n], \Gamma)^{\text{st}}$. We let $(L_{\eta, \alpha}, s_{\eta, \alpha})$ be the restriction of $(\mathbf{L}_\eta, \mathbf{s}_\eta)$ to S_α . By definition, $R_\alpha \triangleq S_\alpha \times_{\mathfrak{M}(W[n], \Gamma)^{\text{st}}} \mathfrak{M}(W_0[n], \eta)^{\text{st}}$ is the subscheme of S_α defined by the vanishing of s_α . Hence R_α/S_α admits an obvious relative

obstruction theory induced by the pair $(L_{\eta,\alpha}, s_{\eta,\alpha})$, as defined in the first case in subsection 4.2. Namely, for any $\xi = (B, I, \varphi) \in \mathfrak{Tri}_{R_\alpha/S_\alpha}$ with $\varphi: \text{Spec } B/I \rightarrow R_\alpha$, the relative obstruction class is

$$\mathbf{ob}_{R_\alpha/S_\alpha}(\xi) = \mathbf{d}(s_\alpha \circ \varphi) \in \Gamma(R_\alpha, L_{\eta,\alpha}) \otimes_{\Gamma(R_\alpha)} I.$$

Hence the relative obstruction theory of R_α/S_α takes values in the cohomology of the complex $[0 \rightarrow \mathcal{O}_{R_\alpha}(L_{\eta,\alpha})]$. Because $L_{\eta,\alpha}$ are the restriction of a global line bundle $\mathbf{L}_\eta$ on $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ and the sections s_α over S_α are the restrictions of the global section $\mathbf{s}_\eta$, the collection of the relative obstruction assignments $\{\mathbf{ob}_{R_\alpha/S_\alpha}\}$ are compatible over $R_{\alpha\beta}$ and thus defines a global relative obstruction theory of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}/\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ taking values in the cohomology of

$$[0 \rightarrow \mathcal{O}_{\mathfrak{M}(W_0[n], \eta)^{\text{st}}}(\mathbf{L}_\eta)].$$

We now let $A_\alpha = \Gamma(\mathcal{O}_{S_\alpha})$ and $A_{\eta,\alpha} = \Gamma(\mathcal{O}_{R_\alpha})$. Without lose of generality, we can assume all S_α are η -admissible (cf. Definition 3.3). We then form the complex of A_α -modules $\mathbf{E}_\alpha^\bullet$ as in (1.20) and the complex of $A_{\eta,\alpha}$ -modules $\mathbf{E}_{\eta,\alpha}^\bullet$ as in (3.10), with the subscript α added to emphasize their dependence on α . Recall that they are the respective complexes that are part of the perfect obstruction theories of S_α and R_α . Further, we have the exact sequence of complexes (3.11). Thus to apply Lemma 4.6 we only need to show that the relative obstruction theory of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}/\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ is compatible to the obstruction theories of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}$ and of $\mathfrak{M}(W[n], \Gamma)^{\text{st}}$, in the sense of Definition 4.1. Because of the exact sequence (3.11), we only need to show that for any triple $\xi = (B, I, \varphi)$ in $\mathfrak{Tri}_{R_\alpha/S_\alpha}$ we have

$$\zeta(\mathbf{ob}_{R_\alpha/S_\alpha}(\xi)) = \mathbf{ob}_{R_\alpha}(\xi).$$

Here ζ is the homomorphism $h^1(\mathcal{C}^{\bullet-1}(\mathcal{O}_{R_\alpha}(L_{\eta,\alpha}))) \rightarrow h^2(\mathbf{E}_{\eta,\alpha}^\bullet(\mathcal{O}_{R_\alpha}))$. But this follows directly from the construction of the respective (relative) obstruction theories. As argued before, the relative obstruction theory of $\mathfrak{M}(W_0[n], \eta)^{\text{st}}/\mathfrak{M}(W[n], \Gamma)^{\text{st}}$ descends to a relative obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)/\mathfrak{M}(\mathfrak{W}, \Gamma)$ taking values in the cohomology of the complex $[0 \rightarrow \mathcal{O}_{\mathfrak{M}(\mathfrak{W}_0, \eta)}(\mathbf{L}_\eta)]$, and this relative obstruction theory is compatible to the obstruction theories of $\mathfrak{M}(\mathfrak{W}_0, \eta)$ and $\mathfrak{M}(\mathfrak{W}, \Gamma)$. Thus by applying the result proved in subsection 4.2, we conclude $[\mathfrak{M}(\mathfrak{W}_0, \eta)]^{\text{virt}} = c_1(\mathbf{L}_\eta, \mathbf{s}_\eta)[\mathfrak{M}(\mathfrak{W}, \Gamma)]^{\text{virt}}$. This proves Lemma 3.11.

4.4. Proof of Lemma 3.12. We now prove Lemma 3.12. Here is our strategy. Let $\mathbf{M}_{g,n}$ be the (Artin) stack of k -pointed genus g nodal curves and let

$$\mu : \mathfrak{M}(\mathfrak{W}, \Gamma) \longrightarrow \mathbf{M}_{g,n}$$

be the forgetful morphism. Let $\eta = (\Gamma_1, \Gamma_2, I) \in \Omega$ be as in Lemma 3.12 that has r ordered roots of weights $\mu_1, \dots, \mu_r$. We will show that in the formal neighborhood of $\mathfrak{M}(\mathfrak{W}_0, \eta)$ in $\mathfrak{M}(\mathfrak{W}, \Gamma)$ (possibly after an étale base change) there are divisors $(\mathbf{L}_i, \mathbf{s}_i)$ for $1 \leq i \leq r$ so that $\mathfrak{M}(\mathfrak{W}_0, \eta)$ is defined by the vanishing of $\mathbf{s}_1^{\mu_1}, \dots, \mathbf{s}_r^{\mu_r}$ while $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ is defined by the vanishing of $\mathbf{s}_1, \dots, \mathbf{s}_r$. This way we can reduce the proof of Lemma 3.12 to the situation studied in Lemma 4.8.

We now provide the detail of the proof. We first construct the desired base change of the formal neighborhood of $\mathfrak{M}(\mathfrak{W}_0, \eta)$ in $\mathfrak{M}(\mathfrak{W}, \Gamma)$. We let $\mathbf{M}_{\Gamma_i^\circ}$ be the moduli stack of pointed nodal curves (not necessary connected) of topological type Γ_i° (see the definition before (2.9)). Since we do not impose stability condition on such curves, $\mathbf{M}_{\Gamma_i^\circ}$ is an Artin stack. Then for any pair $(C_1, C_2) \in \mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}$ we can form a new curve $C_1 \sqcup C_2$ by gluing the pairs of the i -th distinguished marked

point of C_1 and of C_2 for all $i = 1, \dots, r$. This construction extends to families, thus defines a (local embedding) morphism $\mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ} \rightarrow \mathbf{M}_{g,k}$.

Definition 4.9. Let $\mathbf{A}$ be an Artin stack. We say $\mathbf{B}$ is a formal extension of $\mathbf{A}$ if $\mathbf{A}$ is a closed substack of $\mathbf{B}$ and the inclusion $\mathbf{A} \rightarrow \mathbf{B}$ is a homeomorphism.

Lemma 4.10. We can find a formal extension $\mathbf{Q}$ of $\mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}$ so that the morphism $\mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ} \rightarrow \mathbf{M}_{g,k}$ extends to an étale morphism $\mathbf{Q} \rightarrow \mathbf{M}_{g,k}$.

Proof. For schemes, this is the topological equivalence of étale morphisms [Mil]. Note that once such extensions exist, then they are canonical. The proof of the general case can be proved by applying this topological equivalence theorem to charts of the stacks. We will leave the details to the readers. $\square$

We next consider the gluing morphism

$$\Phi_\eta : \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \longrightarrow \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) \subset \mathfrak{M}(\mathfrak{W}, \Gamma).$$

Lemma 4.11. There is a formal extension $\mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge$ of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)$ (as DM-stack) so that the morphism Φ_η extends to an étale morphism

$$\hat{\Phi}_\eta : \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \longrightarrow \mathfrak{M}(\mathfrak{W}, \Gamma).$$

Proof. The proof exactly the same to Lemma 4.10, and will be omitted. $\square$

Because $\mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge$ is homeomorphic to $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)$, the forgetful morphism

$$(4.25) \quad \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2) \longrightarrow \mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}$$

extends to $\mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \rightarrow \mathbf{Q}$.

Our next task is to define the PD-divisors $(\mathbf{L}_i, \mathbf{s}_i)$ on $\mathbf{Q}$ for $i = 1, \dots, r$ as mentioned. We let ξ and ζ be the universal families over $\mathbf{M}_{g,k}$ and $\mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}$. We let ξ_{node} be the natural substack of all nodal points of the fibers of ξ . Then $\xi_{\text{node}} \subset \xi$ is a smooth divisor. On the other hand, each fiber ζ_p (over $p \in \mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}$) contains r ordered distinguished nodes. They define r ordered distinguished sections

$$(4.26) \quad \mathbf{n}_1, \dots, \mathbf{n}_r : \mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ} \longrightarrow \zeta.$$

The nodal locus $\zeta_{\text{node}} \subset \zeta$ is the union of a smooth divisor with the images of these r sections. We now consider the universal family $\bar{\zeta}$ over $\mathbf{Q}$. Since $\mathbf{Q} \rightarrow \mathbf{M}_{g,k}$ is étale, the nodal locus $\bar{\zeta}_{\text{node}}$ is a smooth divisor in $\bar{\zeta}$. On the other hand, since $\bar{\zeta}|_{\mathbf{M}_{\Gamma_1^\circ} \times \mathbf{M}_{\Gamma_2^\circ}} \cong \zeta$, $\zeta_{\text{node}} \subset \bar{\zeta}_{\text{node}}$ and is a homeomorphism. Hence each $\text{Im}(\mathbf{n}_i)$ is an open subset in $\bar{\zeta}_{\text{node}}$. We let $\bar{\mathbf{B}}_i \subset \bar{\zeta}_{\text{node}}$ be the open subscheme that contains and is homeomorphic to $\text{Im}(\mathbf{n}_i)$. We let $\mathbf{B}_i \subset \mathbf{Q}$ be the image stack of $\bar{\mathbf{B}}_i$ under the projection $\bar{\zeta} \rightarrow \mathbf{Q}$. Clearly, $\mathbf{B}_i$ are smooth divisors of $\mathbf{Q}$. We define $(\mathbf{L}_i, \mathbf{s}_i)$ be the C-divisor on $\mathbf{Q}$ so that $\mathbf{B}_i = \mathbf{s}_i^{-1}(0)$.

For later application, we now give trivializations of $(\mathbf{L}_i, \mathbf{s}_i)$ on charts of $\mathbf{Q}$. let $T_\alpha \rightarrow \mathbf{Q}$ be any chart. Then $B_{i,\alpha} = T_\alpha \times_{\mathbf{Q}} \mathbf{B}_i$ is a smooth divisor in T_α . Without loss of generality, we can assume that $B_{i,\alpha}$ is defined by the vanishing of a $u_{i,\alpha} \in \Gamma(\mathcal{O}_{T_\alpha})$. We then choose $L_{i,\alpha}$ be the line bundle over T_α so that $\mathcal{O}_{T_\alpha}(L_{i,\alpha}) = u_{i,\alpha}^{-1} \mathcal{O}_{T_\alpha}$ and let $s_{i,\alpha} \in \Gamma(L_{i,\alpha})$ be the constant 1. Put it differently, $e_{i,\alpha} = u_{i,\alpha}^{-1} 1$ is a global holomorphic basis of $L_{i,\alpha}$ while the section $s_{i,\alpha} = 1 = u_{i,\alpha} e_{i,\alpha}$ vanishes on $B_{i,\alpha}$. In case $\tilde{u}_{i,\alpha}$ is another defining equation of $B_{i,\alpha}$, we define $(\tilde{L}_{i,\alpha}, \tilde{s}_{i,\alpha})$

similarly via a basis $\tilde{e}_{i,\alpha} = \tilde{u}_{i,\alpha}^{-1}1$ and $\tilde{s}_{i,\alpha} = \tilde{u}_{i,\alpha}\tilde{e}_{i,\alpha}$. The transition function is via $\tilde{e}_{i,\alpha} = u_{i,\alpha}/\tilde{u}_{i,\alpha} e_{i,\alpha}$.

Now let μ_i be the weight of the i -th root of η . We define

$$(4.27) \quad \mathbf{N}_0 = \{\mathbf{s}_1 = \cdots = \mathbf{s}_r = 0\} \quad \text{and} \quad \mathbf{N} = \{\mathbf{s}_1^{\mu_1} = \cdots = \mathbf{s}_r^{\mu_r} = 0\} \subset \mathbf{Q},$$

both are substacks of $\mathbf{Q}$. Clearly, $\mathbf{N}_0 = \mathbf{M}_{\Gamma_1^0} \times \mathbf{M}_{\Gamma_2^0}$. Hence

$$\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta) = \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \times_{\mathbf{Q}} \mathbf{N}_0.$$

We define

$$\mathfrak{M}(\mathfrak{W}_0, \eta)^{et} = \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \times_{\mathbf{Q}} \mathbf{N}.$$

Lemma 4.12. *Let $\pi_Q : \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \rightarrow \mathbf{Q}$ and $\pi_M : \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ be the tautological projections. Then we have isomorphisms of C -divisors*

$$\pi_Q^*(\mathbf{L}_i, \mathbf{s}_i)^{\otimes \mu_i} \cong \pi_M^*(\mathbf{L}_\eta, \mathbf{s}_\eta).$$

Further, $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et} \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)$ factor through an étale $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et} \rightarrow \mathfrak{M}(\mathfrak{W}_0, \eta)$.

Proof. We cover $\mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge$ by an atlas of étale charts $S_\alpha \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge$ indexed by Λ . To each $\alpha \in \Lambda$, we let $f_\alpha : \mathcal{X}_\alpha \rightarrow W[n_\alpha]$ be the pull back of the universal family over $\mathfrak{M}(\mathfrak{W}, \Gamma)$. We then cover $\mathbf{Q}$ by charts T_α indexed by the same set Λ . Without loss of generality, we can assume that the composite of $S_\alpha \rightarrow \mathfrak{M}(\mathfrak{W}, \Gamma)^\wedge \rightarrow \mathbf{Q}$ factor through $h_\alpha : S_\alpha \rightarrow T_\alpha \subset \mathbf{Q}$ and the family $\mathcal{X}_\alpha$ is the pull back of the universal family $\bar{\zeta}_\alpha$ on T_α . As before, we let $B_{i,\alpha} = T_\alpha \times_{\mathbf{Q}} \mathbf{B}_i \subset T_\alpha$ and let $u_{i,\alpha}$ be a defining equation of $B_{i,\alpha}$; We let $(L_{i,\alpha}, s_{i,\alpha})$ be the restriction of $(\mathbf{L}_i, \mathbf{s}_i)$ to the chart T_α . As mentioned, we can choose $e_{i,\alpha} = u_{i,\alpha}^{-1}1$ to be a holomorphic basis of $L_{i,\alpha}$, and hence $s_{i,\alpha} = u_{i,\alpha} e_{i,\alpha}$. Now we let $n_{i,\alpha} : B_{i,\alpha} \rightarrow \bar{\zeta}_\alpha|_{B_{i,\alpha}}$ be the lift of the section $\mathbf{n}_i$ in (4.26) to $B_{i,\alpha}$. Since $\mathbf{Q}$ is homeomorphic to $\mathbf{M}_{\Gamma_1^0} \times \mathbf{M}_{\Gamma_2^0}$, such a lift exists and is unique. We then let $N_{i,\alpha} = n_{i,\alpha}(B_{i,\alpha})$ and let $\hat{N}_{i,\alpha}$ be the formal completion of $\bar{\zeta}_\alpha$ along $N_{i,\alpha}$. Without loss of generality, we can assume

$$\hat{N}_{i,\alpha} \cong \text{Spec } \mathbb{k}[[z_1, z_2]] \times_{\text{Spec } \mathbb{k}[t]} T_\alpha.$$

Here $\text{Spec } \mathbb{k}[[z_1, z_2]] \rightarrow \text{Spec } \mathbb{k}[t]$ is defined by $t \mapsto z_1 z_2$ and $T_\alpha \rightarrow \text{Spec } \mathbb{k}[t]$ is defined by $t \mapsto u_{i,\alpha}$.

We now back to the family $f_\alpha : \mathcal{X}_\alpha \rightarrow W[n_\alpha]$. Recall $\mathcal{X}_\alpha = \bar{\zeta}_\alpha \times_{\mathbf{Q}} S_\alpha$. We let $\hat{\mathcal{X}}_{i,\alpha}$ be the formal completion of $\mathcal{X}_\alpha$ along $N_{i,\alpha} \times_{\mathbf{Q}} S_\alpha$, which is isomorphic to $\mathcal{X}_\alpha \times_{\bar{\zeta}_\alpha} \hat{N}_{i,\alpha}$. By shrinking S_α if necessary, we can assume that there is a parameterization of a neighborhood of nodes $\mathcal{W}_\alpha \subset W[n_\alpha]$ given by

$$\psi_\alpha : \mathcal{W}_\alpha \longrightarrow \text{Spec } \mathbb{k}[w_1, w_2] \otimes_{\mathbb{k}[t_{i,\alpha}]} \Gamma(\mathbf{A}^{n_\alpha+1})$$

for some $l_\alpha \in [n_\alpha + 1]$, as in (1.1), so that the induced morphism $\hat{f}_\alpha : \hat{\mathcal{X}}_{i,\alpha} \rightarrow W[n_\alpha]$ factor through

$$(4.28) \quad \hat{f}_\alpha : \hat{\mathcal{X}}_{i,\alpha} \rightarrow \mathcal{W}_\alpha \quad \text{via} \quad \hat{f}_\alpha^*(w_j) = \beta_{j,\alpha} \cdot z_j^{\mu_j}, \quad j = 1, 2.$$

Now let $(L_{\eta,\alpha}, s_{\eta,\alpha})$ be the restriction to S_α of $(\mathbf{L}_\eta, \mathbf{s}_\eta)$. By definition, a trivialization of $(L_{\eta,\alpha}, s_{\eta,\alpha})$ is given by $\mathcal{O}(L_{\eta,\alpha}) = t_{l_\alpha}^{-1} \mathcal{O}_{T_\alpha}$ with the basis $\epsilon_{\eta,\alpha} = t_{l_\alpha}^{-1}1$ and $s_{\eta,\alpha} \equiv t_{l_\alpha} \epsilon_{\eta,\alpha}$. As mentioned, a trivialization of $(L_{i,\alpha}, s_{i,\alpha})$ is given by $\mathcal{O}(L_{i,\alpha}) = u_{i,\alpha}^{-1} \mathcal{O}_{S_\alpha}$ with $e_{i,\alpha} = u_{i,\alpha}^{-1}1$ and $s_{i,\alpha} = u_{i,\alpha} e_{i,\alpha}$. We then define an isomorphism $L_{\eta,\alpha} \cong L_{i,\alpha}^{\otimes \mu_i}$ via

$$\epsilon_{\eta,\alpha} = (\beta_{\alpha,1} \beta_{\alpha,2}) e_{i,\alpha}^{\otimes \mu_i}.$$

Note that since $\beta_{\alpha,1}\beta_{\alpha,2} \in \Gamma(\mathcal{O}_{S_\alpha}^\times)$, the above identity defines an isomorphism $L_{\eta,\alpha} \cong L_{i,\alpha}^{\otimes \mu_i}$. Further, because of the relations (4.28) and $w_1 w_2 = t_{l_\alpha}$,

$$(4.29) \quad s_{\eta,\alpha} = s_{i,\alpha}^{\otimes \mu_i}$$

under this isomorphism. Hence it defines an isomorphism of the corresponding C-divisors. It is routine to check that the so defined isomorphisms extend to an isomorphism of C-divisors

$$\pi_Q^*(\mathbf{L}_\eta, \mathbf{s}_\eta) \cong \pi_M^*(\mathbf{L}_i, \mathbf{s}_i)^{\otimes \mu_i}.$$

This proves the isomorphisms of the C-divisors.

The last statement follows directly from the construction of $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et}$. $\square$

We now consider the pair $\rho : \mathfrak{M}(\mathfrak{W}_0, \eta)^{et} \rightarrow \mathbf{N}$. Since $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et}$ is étale over $\mathfrak{M}(\mathfrak{W}_0, \eta)$, we can take the obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et}$ to be the pull back of that of $\mathfrak{M}(\mathfrak{W}_0, \eta)$. We now workout the relative obstruction theory of $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et}/\mathbf{N}$ and to show that it satisfies the set up in Lemma 4.8. We will follow closely the notation developed in the beginning of Section 4.1. For convenience, we denote $\mathfrak{M}(\mathfrak{W}_0, \eta)^{et}$ by $\mathbf{M}$.

We begin with a smooth chart $\bar{S}$ of $\mathbf{Q}$ and the associated chart $S = \bar{S} \times_{\mathbf{Q}} \mathbf{N}$. We let (L_i, s_i) be the restriction of $(\mathbf{L}_i, \mathbf{s}_i)$ to $\bar{S}$. Since $\mathbf{Q}$ and hence $\bar{S}$ are smooth, the pairs (L_i, s_i) for $i = 1, \dots, r$ define a natural obstruction theory of S taking values in the cohomology of the complex

$$(4.30) \quad \mathcal{F}^\bullet = [\mathcal{O}_S(T\bar{S}) \xrightarrow{ds^{[\mu]}} \oplus_{i=1}^r \mathcal{O}_S(L_i^{\otimes \mu_i})],$$

as defined in the beginning of the section 4.1. Here $ds^{[\mu]}$ is the abbreviation of $(ds_1^{\mu_1}, \dots, ds_r^{\mu_r})$. Now let $\mathcal{X}$ be the universal family over S and let $\mathcal{D} \subset \mathcal{X}$ be the divisor of the marked sections of $\mathcal{X}$. Then by the deformation theory of nodal curves, there is a canonical homomorphism of sheaves (the Kodaira map)

$$\mathcal{O}_S(T\bar{S}) \longrightarrow \mathcal{E}xt_{\mathcal{X}/S}^1(\Omega_{\mathcal{X}/S}(\mathcal{D}), \mathcal{O}_{\mathcal{X}}).$$

Next we pick an affine étale universal open $R \rightarrow S \times_{\mathbf{N}} \mathbf{M}$ with $\rho : R \rightarrow S$ the projection. Without loss of generality we can assume that the universal family of R is of the form $f : \rho^* \mathcal{X} \rightarrow W[n]$ for some integer n . We now construct the standard obstruction theory of R . Since R is a smooth chart (not necessary étale) of $\mathbf{M}$, its obstruction theory is slightly different from that defined in Section 3.1, which is for étale charts of $\mathbf{M}$.

We begin with the construction of the complex $\mathbf{E}_\eta^\bullet$ that will be part of the obstruction theory of R . We let $B = \Gamma(\mathcal{O}_R)$ with $A = \Gamma(\mathcal{O}_S)$ as before. First of all, we choose the complex $F^\bullet$ in (1.15) to be

$$F^0 = 0 \quad \text{and} \quad F^1 = \Gamma(R, \rho^* T\bar{S}) = \Gamma(\mathcal{O}_S(T\bar{S})) \otimes_A B.$$

We pick a collection of charts $(\mathcal{U}_\alpha/\mathcal{V}_\alpha, f_\alpha)$ of f indexed by Λ , as mentioned in the paragraph after (1.17). By shrinking R and/or make an étale base change of $\bar{S}$, we can assume that there is a collection of charts $\{U_\alpha/V_\alpha\}$ of $\mathcal{X}/S$ indexed by the same Λ so that $\mathcal{V}_\alpha = R \times_S V_\alpha$ and $\mathcal{U}_\alpha = R \times_S U_\alpha$. We form the modules $\text{Hom}_{\mathcal{U}}(f^* \Omega_{W[n]}, B)^\dagger$, after picking the necessary data as shown in the paragraphs after equation (1.17). Since we have chosen $F^0 = 0$, the homomorphism $\zeta_\alpha(\cdot)$ in (1.19) is zero. The homomorphism $\zeta_{\alpha\beta}(\cdot)$ is exactly the one defined in (1.18). We then form the complex $\mathbf{D}^\bullet$, the homomorphism δ and the complex $\mathbf{E}^\bullet$, following the definitions after Lemma 1.14, line by line except that we replace F^0 by 0. To form

the complex $\mathbf{E}_\eta^\bullet$, we shall follow the discussion after (3.8). We let $\mathbf{C}_\eta^\bullet$ be the complex defined in (3.7) and let $\delta : \mathbf{E}^\bullet \Rightarrow \mathbf{C}_\eta^{\bullet-1}$ be the homomorphism defined exactly as in (3.8). We then let $\mathbf{E}_\eta^\bullet$ be the complex defined in (3.10). It follows from the discussion before that the obstruction theory of $\mathbf{M}$ induces a natural obstruction theory of R taking values in the cohomology of the complex $\mathbf{E}_\eta^\bullet$.

We now construct a complex $\mathbf{F}^\bullet$ of A -modules that is quasi-isomorphic to $\mathcal{F}^\bullet$ (in (4.30)) and a homomorphism of complexes $\mathbf{E}_\eta^\bullet \Rightarrow \mathbf{F}^\bullet$. Let i be any integer between 1 and r . We let $\Lambda_i \subset \Lambda$ be the subset of the indices α so that $U_\alpha \rightarrow \mathcal{X}$ contains the i -th distinguished nodes of some fibers of $\mathcal{X}/S$. Then the collection $\{U_\alpha\}_{\alpha \in \Lambda_i}$ forms a covering of a neighborhood of the i -th distinguished nodes of the fibers of $\mathcal{X}/S$ and the collection $\{V_\alpha\}_{\Lambda_i}$ forms an étale covering of S . We let $\mathbf{C}_i^k = \mathcal{C}^k(\Lambda_i, L_i^{\otimes \mu_i})$ be the Čech complex of k -cochains of the line bundle $L_i^{\otimes \mu_i}$ associated to the covering $\{V_\alpha\}_{\Lambda_i}$. The complex $\mathbf{C}_i^k$ comes with the standard coboundary operator $\mathbf{C}_i^k \rightarrow \mathbf{C}_i^{k+1}$. We define $\mathbf{F}^1 = \Gamma(\mathcal{O}_S(T\bar{S}))$ and define $\mathbf{F}^k = \bigoplus_{i=1}^r \mathbf{C}_i^{k-2}$ for $k \geq 2$.

We next define the differential $\partial^k : \mathbf{F}^k \rightarrow \mathbf{F}^{k+1}$. Let $\mathcal{I}$ be the ideal sheaf of the zero section of the total space $T\bar{S}$ and let $\bar{S}^{(2)}$ be the subscheme of $T\bar{S}$ defined by the ideal sheaf $\mathcal{I}^2$. Then there is a tautological morphism $\kappa : \bar{S}^{(2)} \rightarrow \bar{S}$ characterized by the following property: Let $v \in T_p\bar{S}$ with $\text{Spec } \mathbb{k}[t]/(t^2) \rightarrow T_p\bar{S}$ its associated morphism that lifts to the morphism $[v] : \text{Spec } \mathbb{k}[t]/(t^2) \rightarrow \bar{S}^{(2)}$. Then the composite $\kappa \circ [v] : \text{Spec } \mathbb{k}[t]/(t^2) \rightarrow \bar{S}$ is the tangent vector $v \in T_p\bar{S}$. Now let $\xi \in \Gamma(\mathcal{O}_S(T\bar{S}))$ be any element. We let $\tilde{S} = \text{Spec } \Gamma(\mathcal{O}_S) * A$ be the trivial extension of S by A and let $\theta_\xi : \tilde{S} \rightarrow \bar{S}$ be the composite $\tilde{S} \rightarrow \bar{S}^{(2)} \xrightarrow{\kappa} \bar{S}$, where $\tilde{S} \rightarrow \bar{S}^{(2)}$ is defined by ξ . Since $\theta_\xi^*(s_i^{\mu_i}) \in \Gamma(\theta_\xi^* L_i^{\otimes \mu_i})$ and $\theta_\xi^*(s_i^{\mu_i})|_S \equiv 0$,

$$\mathbf{d}[\theta_\xi^*(s_i^{\mu_i}) - 0] \in \Gamma(S, \mathcal{O}_S(L_i^{\otimes \mu_i}) \otimes \mathcal{I}_{S \subset \tilde{S}}) = \Gamma(S, \mathcal{O}_S(L_i^{\otimes \mu_i})).$$

Here $\mathcal{I}_{S \subset \tilde{S}}$ is the ideal of $S \subset \tilde{S}$, which is isomorphic to A . Now let $\rho_\alpha^* \theta_\xi^*(s_i^{\mu_i}) \in \Gamma(\mathcal{V}_\alpha, \mathcal{O}_S(L_i^{\otimes \mu_i}))$ be the pull back under $\rho_\alpha : \mathcal{V}_\alpha \rightarrow S$. We then define

$$\partial_i^1(\xi)_\alpha = \mathbf{d}[\rho_\alpha^* \theta_\xi^*(s_i^{\mu_i}) - 0] \in \Gamma(\mathcal{V}_\alpha, \mathcal{O}_S(L_i^{\otimes \mu_i})), \quad \alpha \in \Lambda_i,$$

and define $\partial^1 = \bigoplus_i \partial_i^1$. For $k \geq 1$ we let $\partial^k : \mathbf{F}^k \rightarrow \mathbf{F}^{k+1}$ to be the direct sum of the coboundary operators of $\mathbf{C}_i^\bullet$. Clearly, the so defined operator $\partial^\bullet$ satisfies $\partial^k \circ \partial^{k+1} = 0$, and hence defines a complex $\mathbf{F}^\bullet = (\mathbf{F}^k, \partial^k)$. Further, it follows from our construction that the complex $\mathbf{F}^\bullet$ is quasi-isomorphic to the complex $\mathcal{F}^\bullet$ in (4.30).

We now define the promised homomorphism $\varphi^\bullet : \mathbf{E}_\eta^\bullet \Rightarrow \mathbf{F}^\bullet \otimes_A B$. Recall that $\mathbf{E}_\eta^1 = \Gamma(\mathcal{O}_S(T\bar{S})) \otimes_A B \oplus \mathbf{D}^0$ and $\mathbf{F}^1 = \Gamma(\mathcal{O}_S(T\bar{S}))$. The homomorphism $\varphi^1 : \mathbf{E}_\eta^1 \rightarrow \mathbf{F}^1 \otimes_A B$ is the one induced by the identity of $\Gamma(\mathcal{O}_S(T\bar{S})) \otimes_A B$. For $k \geq 2$, we notice that $\mathbf{E}_\eta^k = \mathbf{E}^k \oplus \mathbf{C}_\eta^{k-2}$ and $\mathbf{F}^k = \bigoplus_{i=1}^r \mathbf{C}_i^{k-2}$. The homomorphism φ^k will be induced by $\varphi_i^k : \mathbf{C}_\eta^k \rightarrow \mathbf{C}_i^k \otimes_A B$, which we define now. Let $\xi \in \mathbf{C}_\eta^k$ be any element and let $(\alpha_0 \cdots \alpha_k)$ be a $(k+1)$ -tuple in Λ_i . Then $\xi_{\alpha_0 \cdots \alpha_k}$ is an element in $\Gamma(\mathcal{V}_{\alpha_0 \cdots \alpha_k}, L_\eta)$. Here L_η is the restriction of L_η to R . Using the canonical isomorphism $L_\eta \cong \rho^* L_i^{\otimes \mu_i}$, where $\rho : R \rightarrow S$ is the tautological projection,

$$\xi_{\alpha_0 \cdots \alpha_k} \in \Gamma(\mathcal{V}_{\alpha_0 \cdots \alpha_k}, \rho^* L_i^{\otimes \mu_i}) = \Gamma(\mathcal{V}_{\alpha_0 \cdots \alpha_k}, L_i^{\otimes \mu_i}) \otimes_A B.$$

We denote this element by $\tilde{\xi}_{\alpha_0 \cdots \alpha_k}$. We define

$$\varphi_i^k(\xi)_{\alpha_0 \cdots \alpha_k} = \tilde{\xi}_{\alpha_0 \cdots \alpha_k} \in \Gamma(\mathcal{V}_{\alpha_0 \cdots \alpha_k}, L_i^{\otimes \mu_i}) \otimes_A B.$$

This defines a homomorphism $\mathbf{C}_\eta^k \rightarrow \bigoplus_{i=1}^r \mathbf{C}_i^k \otimes_A B$.

We claim that the so defined homomorphisms form a homomorphism of complexes $\mathbf{E}_\eta^\bullet \Rightarrow \mathbf{F}^\bullet \otimes_A B$. For this, we need to check the commutativity of the following diagram

$$(4.31) \quad \begin{array}{ccc} \mathbf{E}_\eta^k & \xrightarrow{d^k} & \mathbf{E}_\eta^{k+1} \\ \varphi^k \downarrow & & \varphi^{k+1} \downarrow \\ \mathbf{F}^k \otimes_A B & \xrightarrow{\partial^k} & \mathbf{F}^{k+1} \otimes_A B. \end{array}$$

We will prove the commutativity of (4.31) for the case $k = 1$. The other cases are similar and will be omitted. First, we show that (4.31) commutes on $\mathbf{D}^0 \subset \mathbf{E}_\eta^1$. Let $\xi \in \mathbf{D}^0 \subset \mathbf{E}_\eta^1$ be any element given by a collection $\{\xi_\alpha\}$ of $\xi_\alpha \in \text{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, \mathcal{O}_S)^\dagger$, as defined in (1.11). Because of the relation (1.12), the composite $\mathbf{D}^0 \subset \mathbf{E}_\eta^1 \rightarrow \mathbf{E}_\eta^2 \xrightarrow{\varphi^2} \mathbf{F}^2 \otimes_A B$ is zero. On the other hand, by definition we have $\varphi^1(\xi) = 0$. Hence $\varphi^2(\partial^1(\xi)) = d^1(\varphi^1(\xi))$ for all $\xi \in \mathbf{D}^0$.

We next check that (4.31) commutes on $\Gamma(\mathcal{O}_S(T\bar{S})) \subset \mathbf{E}_\eta^1$. We first recall the definition of $\varphi^2 \circ d^1$. Let $\xi \in \Gamma(\mathcal{O}_S(T\bar{S}))$ be any element. It determines a morphism $\tilde{\kappa}_\xi: \tilde{R} = \text{Spec } \Gamma(\mathcal{O}_R) * B \rightarrow \bar{S}$ that is the pull back of $\kappa_\xi: \tilde{S} \rightarrow \bar{S}$ via the tautological extension $\tilde{\rho}: \tilde{R} \rightarrow \tilde{S}$ of $\rho: R \rightarrow S$. We let $\alpha \in \Lambda_i$ be any index with U_α/V_α the associated chart of the universal family $\mathcal{X}/S$ and with $\tilde{U}_\alpha/\tilde{V}_\alpha$ its minimal extension to the family $\tilde{\mathcal{X}}/\tilde{S}$. We let $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ and $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ be the corresponding pull back charts of $\rho^*\mathcal{X}/R$ and $\tilde{\rho}^*\tilde{\mathcal{X}}/\tilde{R}$. We then pick a local parameterization of the nodes of U_α/V_α and its extension to $\tilde{U}_\alpha/\tilde{V}_\alpha$. We let $(z_{\alpha,i}, s_\alpha)$ and $(\tilde{z}_{\alpha,i}, \tilde{s}_\alpha)$ be the relevant functions associated to these parameterizations²⁰. We let the parameterizations of the nodes of $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ and $\tilde{\mathcal{U}}_\alpha/\tilde{\mathcal{V}}_\alpha$ be the pull back of those from U_α/V_α and $\tilde{U}_\alpha/\tilde{V}_\alpha$. We let $f_\alpha: \mathcal{U}_\alpha \rightarrow W[n]$ be the restriction of f to $\mathcal{U}_\alpha$ and let $\tilde{f}_\alpha: \tilde{\mathcal{U}}_\alpha \rightarrow W[n]$ be a pre-deformable extension of f_α . We let $\iota_\alpha: \mathcal{U}_\alpha \rightarrow U_\alpha$, $\tilde{\iota}: \tilde{\mathcal{U}}_\alpha \rightarrow \tilde{U}_\alpha$, $j_\alpha: \mathcal{V}_\alpha \rightarrow V_\alpha$, $\tilde{j}_\alpha: \tilde{\mathcal{V}}_\alpha \rightarrow \tilde{V}_\alpha$, $\rho_\alpha: U_\alpha \rightarrow V_\alpha$ and $\tilde{\rho}_\alpha: \tilde{U}_\alpha \rightarrow \tilde{V}_\alpha$ be the tautological projections. Then after picking a local parameter of $f_\alpha(\mathcal{U}_\alpha) \subset W[n]$, say (w_1, w_2) with $w_1 w_2 = t_{l_\alpha}$, we have

$$f_\alpha^*(w_{\alpha,i}) = \iota_\alpha^*(z_{\alpha,i}^{\mu_i}) \cdot h_{\alpha,i} \quad \text{and} \quad \tilde{f}_\alpha^*(w_{\alpha,i}) = \tilde{\iota}_\alpha^*(\tilde{z}_{\alpha,i}^{\mu_i}) \cdot \tilde{h}_{\alpha,i}$$

for some $h_{\alpha,i} \in \Gamma(\mathcal{O}_{\mathcal{U}_\alpha}^\times)$ and their extensions $\tilde{h}_{\alpha,i} \in \Gamma(\mathcal{O}_{\tilde{\mathcal{U}}_\alpha}^\times)$ that satisfy $h_{\alpha,1} h_{\alpha,2} \in \Gamma(\mathcal{O}_{\mathcal{V}_\alpha})$ and $\tilde{h}_{\alpha,1} \tilde{h}_{\alpha,2} \in \Gamma(\mathcal{O}_{\tilde{\mathcal{V}}_\alpha})$. Since $\tilde{z}_{\alpha,1} \tilde{z}_{\alpha,2} = \tilde{s}_\alpha$, we have

$$\tilde{f}_\alpha^*(t_{l_\alpha}) = (\tilde{h}_{\alpha,1} \tilde{h}_{\alpha,2}) \tilde{\iota}_\alpha^*(\tilde{s}_\alpha^{\mu_i}).$$

Then

$$\varphi^2(d^1(\xi))_\alpha = \mathbf{d}[\tilde{f}_\alpha^*(t_{l_\alpha}) - 0] = (h_{\alpha,1} h_{\alpha,2}) \mathbf{d}[\tilde{j}_\alpha^*(\tilde{s}_\alpha^{\mu_i}) - 0] \in \Gamma(\mathcal{V}_\alpha, \rho^* L_{\eta,\alpha} \otimes I_{R \subset \tilde{R}}).$$

Here the last relation follows because $\tilde{f}_\alpha$ is pre-deformable.

As to $d^1 \circ \varphi^1$, by definition

$$d^1(\varphi^1(\xi))_\alpha = \mathbf{d}[\tilde{s}_\alpha^{\mu_i} - 0] \in \Gamma(V_\alpha, \rho^* L_i^{\otimes \mu_i} \otimes I_{S \subset \tilde{S}}).$$

Therefore we have $\varphi^2(\partial^1(\xi))_\alpha = d^1(\varphi^1(\xi))_\alpha$ because of the isomorphism $L_\eta \cong L_i^{\otimes \mu_i}$, (4.31), the relation $t_{l_\alpha} = (h_{\alpha,1} h_{\alpha,2}) s_\alpha^{\mu_i}$ and $I_{R \subset \tilde{R}} \cong B$ and $T_{S \subset \tilde{S}} \cong A$. This proves the commutativity of (4.31) for $k = 1$. The case $k \geq 2$ is similar and will be omitted.

²⁰By abuse of notation we will view s_α as functions on $\mathcal{V}_\alpha$ and on $\mathcal{U}_\alpha$ via the pull back $\mathcal{O}_{\mathcal{V}_\alpha} \rightarrow \mathcal{O}_{\mathcal{U}_\alpha}$. The same convention applies to $\tilde{s}_\alpha$ as well.

We define $\mathbf{G}^k = \ker\{\mathbf{E}_\eta^k \rightarrow \mathbf{F}^k \otimes_A B\}$. Since $\mathbf{E}_\eta^k \rightarrow \mathbf{F}^k \otimes_A B$ is surjective, $\mathbf{G}^k$ is a flat $\mathcal{O}_S$ -module. The differentials of $\mathbf{E}_\eta^\bullet$ induces differentials of $\mathbf{G}^\bullet$ and the resulting complex fits into the following exact sequence

$$0 \Rightarrow \mathbf{G}^\bullet \Rightarrow \mathbf{E}_\eta^\bullet \Rightarrow \mathbf{F}^\bullet \otimes_A B \Rightarrow 0.$$

It is routine to check that for each $\xi = (B', I, \varphi_0) \in \mathfrak{Tri}_{S/R}$ there is a canonical obstruction class $\mathbf{ob}_{R/S}(\xi) \in h^2(\mathbf{G}^\bullet \otimes I)$ to extending $\varphi_0 : \mathrm{Spec} B'/I \rightarrow R$ to an S -morphism $\mathrm{Spec} B' \rightarrow R$, and further such assignment satisfies the requirement in Definition 4.1. Finally, we remark that though the complexes $\mathbf{G}^\bullet$, $\mathbf{E}_\eta^\bullet$ and $\mathbf{F}^\bullet$ depend on the choice of the covering $\{\mathcal{U}_\alpha/\mathcal{V}_\alpha\}$ of f , they as elements in the derived category are unique. In particular the modules (sheaves) $\mathcal{O}b_{R/S} \triangleq h^2(\mathbf{G}^\bullet)$, $\mathcal{O}b_R \triangleq h^2(\mathbf{E}_\eta^\bullet)$ and $\mathcal{O}b_S \triangleq h^2(\mathbf{F}^\bullet)$ are independent of the choice of the coverings.

We now cover $\mathbf{N}$ by an atlas $\{S_\alpha\}_\Xi$ and for each $\alpha \in \Xi$ we pick an open étale $R_\alpha \rightarrow \mathbf{M} \times_{\mathbf{N}} S_\alpha$ so that $\{R_\alpha\}_\Xi$ forms an atlas of $\mathbf{M}$. For each α we pick a sufficiently fine covering of its universal family and then form the associated complexes $\mathbf{G}_\alpha^\bullet$, $\mathbf{E}_{\eta,\alpha}^\bullet$ and $\mathbf{F}_\alpha^\bullet$. Here we added the subscript α to indicate the dependence on the chart R_α/S_α . To be consistent with the notation in Definition 4.1, we let $\mathcal{L}_\alpha^\bullet = \mathbf{G}_\alpha^\bullet$, $\mathcal{E}_\alpha^\bullet = \mathbf{E}_{\eta,\alpha}^\bullet$ and $\mathcal{F}_\alpha^\bullet = \mathbf{F}_\alpha^\bullet$, viewed as complexes of sheaves of $\mathcal{O}_{R_\alpha}$ or $\mathcal{O}_{S_\alpha}$ -modules accordingly.

Lemma 4.13. *There are standard relative obstruction theories of R_α/S_α for $\alpha \in \Xi$ taking values in the complexes $\mathcal{L}_\alpha^\bullet$ such that the (relative) obstruction theories $\{\mathcal{E}_\alpha^\bullet, \mathbf{ob}_{R_\alpha}\}$, $\{\mathcal{F}_\alpha^\bullet, \mathbf{ob}_{S_\alpha}\}$ and $\{\mathcal{L}_\alpha^\bullet, \mathbf{ob}_{R_\alpha/S_\alpha}\}$ are compatible in the sense of definition 4.1. Further, the so defined obstruction theories of R_α/S_α define a relative obstruction theory of $\mathbf{M}/\mathbf{N}$ that is compatible to the obstruction theories of $\mathbf{M}$ and $\mathbf{N}$.*

Proof. The proof is routine and will be omitted. $\square$

Let $\mathbf{N}_0 \subset \mathbf{N}$ be as in (4.27) and let $\mathbf{M}_0 = \mathbf{M} \times_{\mathbf{N}} \mathbf{N}_0$. By definition $\mathbf{M}_0$ is an étale cover of $\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)$. Note that $\mathbf{N}_0$ is smooth. As to $\mathbf{M}_0$, we endow it with the induced obstruction theory of $\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)$, which is perfect since that of $\mathfrak{M}(\mathfrak{Y}_1^{\mathrm{rel}} \sqcup \mathfrak{Y}_2^{\mathrm{rel}}, \eta)$ is. We let $[\mathbf{M}_0]^{\mathrm{virt}}$ be the virtual moduli cycle of $\mathbf{M}_0$. Then Lemma 3.12 is equivalent to

$$(4.32) \quad \mathbf{m}(\eta)[\mathbf{M}_0]^{\mathrm{virt}} = [\mathbf{M}]^{\mathrm{virt}} \in A_*\mathbf{M}.$$

By Lemma 4.8, to prove this identity it suffices to show that 1) *there is a relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$ that is compatible to the obstruction theory of $\mathbf{M}_0$ and $\mathbf{N}_0$* and 2) *the relative obstruction theory $\mathbf{M}_0/\mathbf{N}_0$ is compatible to the relative obstruction theory of $\mathbf{M}/\mathbf{N}$.*

The proof of 1) is parallel to the construction of the relative obstruction theory of $\mathbf{M}/\mathbf{N}$. The proof of 2) is immediate once the relative obstruction theory was constructed. Since the proof is routine, we will leave it to the readers. We state it as a Lemma.

Lemma 4.14. *The standard relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$ is compatible to the obstruction theories of $\mathbf{M}_0$ and $\mathbf{N}_0$. Further, the relative obstruction theory of $\mathbf{M}_0/\mathbf{N}_0$ is induced from the obstruction theory of $\mathbf{M}/\mathbf{N}$.*

In the end, we apply Lemma 4.8 to the pairs $\mathbf{M}_0/\mathbf{N}_0 \subset \mathbf{M}/\mathbf{N}$ to conclude (4.32). This completes the proof of Lemma 3.12.

4.5. **Proof of Lemma 3.14.** It remains to prove Lemma 3.14. Let

$$(4.33) \quad \Phi_\eta : \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2) \longrightarrow \mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$$

be the étale morphism in (3.2). Using the Cartesian product (3.14), we can give $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$ a canonical obstruction theory. We call such obstruction theory the obstruction theory induced by the Cartesian product. On the other hand, since Φ_η is étale, the obstruction theory of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$ induces an obstruction theory of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$ as well.

Lemma 4.15. *The two obstruction theories of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$, one defined by the Cartesian product and the other induced by that the $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)$, are identical.*

Proof. The proof is similar to that in [LT2], and will be omitted. $\square$

We now consider the virtual moduli cycle of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$. Since the obstruction theory of $\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)$ is induced by the Cartesian product,

$$[\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}} = \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}).$$

On the other hand, because of Lemma 4.15 and because Φ_η is étale of pure degree $|\text{Eq}(\eta)|$,

$$\frac{1}{|\text{Eq}(\eta)|} \Phi_{\eta*} \Delta^!([\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}}, \Gamma_1)]^{\text{virt}} \times [\mathfrak{M}(\mathfrak{Y}_2^{\text{rel}}, \Gamma_2)]^{\text{virt}}) = [\mathfrak{M}(\mathfrak{Y}_1^{\text{rel}} \sqcup \mathfrak{Y}_2^{\text{rel}}, \eta)]^{\text{virt}}.$$

This is exactly Lemma 3.14.

This completes the proof of the degeneration formula of the Gromov-Witten invariants stated in the beginning of this paper.

5. APPENDIX

5.1. **The tangent and the obstruction spaces.** In this appendix, we will express the first order deformation and the obstruction spaces of an $\mathfrak{M}(\mathfrak{W}, \Gamma)$ and $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ in terms of some known cohomology groups. As a corollary, we will show that the obstruction theories we constructed in this paper are all perfect.

Let $S \rightarrow \mathfrak{M}(W[n], \Gamma)^{\text{st}}$ be an affine étale chart. As before, we denote by $f : \mathcal{X} \rightarrow W[n]$ be the universal family with $\mathcal{D} \subset \mathcal{X}$ the divisor of the ordinary marked sections. As before, we let $\pi : \mathcal{X} \rightarrow S$ be the projection and let $\rho : S \rightarrow \mathbf{A}^{n+1}$ be the morphism under f . By shrinking and making an étale base change, we can assume the following holds for S : For each l the projection induced by π

$$f^{-1}(\mathbf{D}_l)_{\text{red}} \longrightarrow (S \times_{\mathbf{A}^{n+1}} \mathbf{H}^l)_{\text{red}}$$

is a union of r_l disjoint sections $\sigma_{l,i} : (S \times_{\mathbf{A}^{n+1}} \mathbf{H}^l)_{\text{red}} \rightarrow f^{-1}(\mathbf{D}_l)_{\text{red}}$ for $i = 1, \dots, r_l$. (Note that r_l could be zero.) We pick an atlas $(\mathcal{U}_\alpha / \mathcal{V}_\alpha, f_\alpha)$ of f indexed by Λ so that for each l and $i \in [1, r_l]$ there is exactly one and only one α so that $\mathcal{U}_\alpha \cap \text{Im}(\sigma_{l,i}) \neq \emptyset$, and hence $\text{Im}(\sigma_{l,i})$ is covered by $\mathcal{U}_\alpha$. As before, we let Λ_l be the collection of those α so that $\mathcal{U}_\alpha / \mathcal{V}_\alpha$ covers $\text{Im}(\sigma_{l,i})$ for some i . When α is of the second kind, we let $(z_{\alpha,1}, z_{\alpha,2}, s_\alpha)$ be the parameterization of the distinguished nodes of $\mathcal{U}_\alpha / \mathcal{V}_\alpha$. We require $s_\alpha^{m_\alpha} = g^*(t_l)$ in case $\alpha \in \Lambda_l$. With such assumptions and choices made, the standard log structure of S is given by the pre-log structure $\mathcal{N}_S = \oplus N_l \rightarrow \mathcal{O}_S$ given in (1.7).

We now let $\mathbf{E}^\bullet$ and $\mathbf{D}^\bullet$ be the complexes associated to the perfect obstruction theory of S constructed in section 1. Let $A = \Gamma(\mathcal{O}_S)$. For simplicity, we give an ordering of Λ_l and thus the r_l charts in Λ_l are $\mathcal{U}_{l,1} / \mathcal{V}_{l,1}, \dots, \mathcal{U}_{l,r_l} / \mathcal{V}_{l,r_l}$. We let $\mathbf{R}_l^\bullet$ be the complex

$$(s_{l,1}^{m_{l,1}}, \dots, s_{l,r_l}^{m_{l,r_l}}) : \mathcal{O}_S^{\oplus r_l} \longrightarrow (\mathcal{O}_S^{\oplus r_l} / \mathcal{O}_S) \otimes_{\mathcal{O}_{\mathbf{A}^{n+1}}} \mathcal{O}_{\mathbf{A}^{n+1}}(\mathbf{H}^l),$$

where $\mathcal{O}_S^{\oplus r_l}/\mathcal{O}_S$ is the quotient of $\mathcal{O}_S^{\oplus r_l}$ by the diagonal $\mathcal{O}_S \hookrightarrow \mathcal{O}_S^{\oplus r_l}$. In case $r_l = 0$, we agree $\mathbf{R}_l^\bullet = [A \rightarrow 0]$. Here the complex $\mathbf{R}_l^\bullet$ is indexed at $[0, 1]$.

Proposition 5.1. *For any A -module I , we have the following two exact sequences*

$$\begin{aligned} 0 \longrightarrow \mathrm{Ext}_{\mathcal{X}}^0(\Omega_{\mathcal{X}/S}(\mathcal{D}), \mathcal{I}) &\longrightarrow H^0(\mathbf{D}^\bullet \otimes_A I) \longrightarrow H^1(\mathbf{E}^\bullet \otimes_A I) \longrightarrow \\ &\longrightarrow \mathrm{Ext}_{\mathcal{X}}^1(\Omega_{\mathcal{X}/S}(\mathcal{D}), \mathcal{I}) \longrightarrow H^1(\mathbf{D}^\bullet \otimes_A I) \longrightarrow H^2(\mathbf{E}^\bullet \otimes_A I) \longrightarrow 0 \\ 0 \longrightarrow H^0(\mathrm{Hom}(f^*\Omega_{W[n]^\dagger/\mathbf{A}^{n+1}^\dagger}, \mathcal{I})) &\longrightarrow H^0(\mathbf{D}^\bullet \otimes_A I) \xrightarrow{b_0} \bigoplus_{l=1}^{n+1} H^0(\mathbf{R}_l^\bullet \otimes_{\mathcal{O}_S} I) \longrightarrow \\ &\xrightarrow{\delta} H^1(\mathrm{Hom}(f^*\Omega_{W[n]_0^\dagger/\mathbf{A}^{n+1}^\dagger}, \mathcal{I})) \longrightarrow H^1(\mathbf{D}^\bullet \otimes_A I) \xrightarrow{b_1} \bigoplus_{l=1}^{n+1} H^0(\mathbf{R}_l^\bullet \otimes_{\mathcal{O}_S} I) \longrightarrow 0 \end{aligned}$$

and the vanishing $H^i(\mathbf{E}^\bullet) = 0$ for $i > 2$ and $H^i(\mathbf{D}^\bullet) = 0$ for $i \geq 2$. Here $\mathcal{I} = I \otimes_A \mathcal{O}_{\mathcal{X}}$.

Proof. The first exact sequence follows directly from the construction of the complexes $\mathbf{D}^\bullet$ and $\mathbf{E}^\bullet$. We now prove the second exact sequence. We first construct the arrow b_0 . Let $\xi \in H^0(\mathbf{D}^\bullet \otimes I)$ be any element. Then ξ is represented by $\{\xi_\alpha\}$ where $\xi_\alpha \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, I)^\dagger$. In case $\alpha \in \Lambda_l$ then

$$(5.1) \quad \xi_\alpha = (\varphi_\alpha, \eta_{\alpha,1}, \eta_{\alpha,2}) \in \mathrm{Hom}_{\mathcal{U}_\alpha}(f^*\Omega_{W[n]}, \mathcal{I}_\alpha) \oplus \mathcal{I}_\alpha^{\oplus 2}, \quad \mathcal{I}_\alpha = \mathcal{I} \otimes_{\mathcal{O}_{\mathcal{X}}} \mathcal{O}_{\mathcal{U}_\alpha},$$

following the convention in (1.10). Specifically, the η means $dw_i/w_i \mapsto \eta_{\alpha,i}$, under the appropriate parameterization (w_1, w_2) of $f_\alpha(\mathcal{U}_\alpha)$. Recall Λ_l was given an ordering, thus for $\alpha = i \in \Lambda_l$ we denote $\eta_{(l,i)} = \eta_{\alpha,1} + \eta_{\alpha,2}$. Then we define

$$b_0(\xi)_l = (\eta_{(l,1)}, \dots, \eta_{(l,r_l)}) \in A^{\oplus r_l}$$

and then the arrow b_0 is

$$b_0(\xi) = (b_0(\xi)_1, \dots, b_0(\xi)_{r_l}).$$

In case $r_l = 0$, then we pick an $\alpha \in \Lambda$ and define $\eta_l = \varphi_\alpha(dt_l)$, where φ_α is part of the ξ_α as in (5.1). It is direct to check that if ξ is a cohomology class then $b_0(\xi)_l \in H^0(\mathbf{R}_l^\bullet \otimes I)$. This defines the corresponding exact sequence in the second exact sequence.

We next construct the arrow $H^1(\mathbf{D}^\bullet) \rightarrow H^1(\mathbf{R}^\bullet)$. Let $\Sigma_l = f^{-1}(\mathbf{D}_l) \subset \mathcal{X}$ and let $I_{\Sigma_l \subset \mathcal{X}}$ be the relative locally constant ideal sheaf defined by $\Gamma(\mathcal{U}_\alpha, I_{\Sigma_l \subset \mathcal{X}}) = \Gamma(\mathcal{V}_\alpha, \mathcal{O}_{\mathcal{V}_\alpha})$ in case $\mathcal{U}_\alpha \cap f^{-1}(\mathbf{D}_l) = \emptyset$ and is $s_\alpha^{m_\alpha} \Gamma(\mathcal{V}_\alpha, \mathcal{O}_{\mathcal{V}_\alpha})$ otherwise. Similarly, we let $\pi^{-1}\mathcal{O}_S$ be the pull back sheaf, namely $\Gamma(\mathcal{U}_\alpha, \pi^{-1}\mathcal{O}_S) = \Gamma(\mathcal{V}_\alpha, \mathcal{O}_{\mathcal{V}_\alpha})$. Then we have the exact sequence in étale site

$$0 \longrightarrow I_{\Sigma_l \subset \mathcal{X}} \longrightarrow \pi^{-1}\mathcal{O}_S \longrightarrow \bigoplus_{\alpha \in \Lambda_l} (\mathcal{O}_S / s_\alpha^{m_\alpha} \mathcal{O}_S) \longrightarrow 0$$

and its induced exact sequence in cohomologies (of $\mathcal{O}_S$ -modules)

$$0 \longrightarrow g^*(t_l)\mathcal{O}_S \longrightarrow \mathcal{O}_S \longrightarrow \bigoplus_{\alpha \in \Lambda_l} (\mathcal{O}_S / s_\alpha^{m_\alpha} \mathcal{O}_S) \longrightarrow H^1(\mathbf{R}_l^\bullet) \longrightarrow 0.$$

Now we back to the complex $\mathbf{D}^\bullet$. It follows from the construction of $\mathbf{D}^\bullet$ that there is a homomorphism of complexes $\mathbf{D}^\bullet \rightarrow \mathbf{C}^\bullet(\Lambda, I_{\Sigma_l \subset \mathcal{X}})$. Thus we have a homomorphism of cohomologies $b_{1,l}: H^1(\mathbf{D}^\bullet \otimes_A I) \rightarrow H^1(\mathbf{R}_l^\bullet \otimes_A I)$ for each l . This forms the arrow in the second exact sequence of the Proposition. The arrow δ in the second sequence is the ordinary connecting homomorphism. We leave it to readers to check that with these arrows the second sequence is exact.

The vanishing result stated in Lemma 1.19 follows from these exact sequences. $\square$

The tangent and the obstruction to deformation of $\mathfrak{M}(\mathfrak{Z}^{\mathrm{rel}}, \Gamma)$ is similar. We let $S \rightarrow \mathfrak{M}(Z[n]^{\mathrm{rel}}, \Gamma)^{\mathrm{st}}$ be an affine étale chart, satisfying the similar property as in the case just studied. We let $f: \mathcal{X} \rightarrow Z[n]$ be the universal family with over S with $\mathcal{D} \subset \mathcal{X}$ be the divisor of the union of all ordinary and the distinguished marked sections. We let $(z_{\alpha,1}, z_{\alpha,2}, s_\alpha)$ be the parameterization of charts $\mathcal{U}_\alpha/\mathcal{V}_\alpha$ of f as in the previous case. We let Λ_l be those such that $\mathcal{U}_\alpha \cap f^{-1}(\mathbf{B}_l) \neq \emptyset$. Let $\mathbf{E}^\bullet$ and $\mathbf{D}^\bullet$ be the complexes constructed in the section 1 that is part of the perfect obstruction theory of S .

Proposition 5.2. *The two exact sequences in Proposition 5.1 still hold with the sheaf $\Omega_{W[n]^\dagger/\mathbf{A}^{n+1}^\dagger}$ replaced by $\Omega_{Z[n]^\dagger/\mathbf{A}^{n+1}^\dagger}$. The same vanishing results hold as well.*

We will close this section by working out the obstruction sheaf of an example suggested to us by E. Ionel.

Let $Z_{\text{rel}} = (Z, D)$ be a pair of smooth variety and a smooth divisor. We let Γ be the graph consisting of one vertex and one leg. We assign the weights of the vertex to be $g = 1$ and $d = 0$. Thus $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ is the moduli of relative stable morphisms to Y from 1-pointed genus 1 curves to Z of degree 0. Since $d = 0$, all $f : X \rightarrow Z$ in $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ are constant maps. Hence $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ is isomorphic to $\mathfrak{M}_{1,1} \times Z$. We now show that its obstruction sheaf is

$$(5.2) \quad \mathcal{O}b = \pi_2^* \Omega_Z(\log D)^\vee,$$

where $\pi_2 : \mathfrak{M}_{1,1} \times Z \rightarrow Z$ is the second projection.

Let $f_0 \in \mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ be a relative stable morphism. Since $d = 0$, we can always represent f_0 by a morphism $f_0 : X \rightarrow Z[1]^\circ$, where $Z[1]^\circ = Z[1] - D[1] \cup Z[1]_{0,\text{sing}}$ with $Z[1]_{0,\text{sing}}$ is the singular locus of $Z[1]_0$. Then $Z[1]^\circ/\mathbb{C}^* \equiv Z$. The obstruction to deforming f_0 as morphism to $Z[1]^\circ$ is $H^1(f_0^* T_{Z[1]^\circ/\mathbf{A}^1}) \equiv T_{Z[1]^\circ/\mathbf{A}^1}|_{f_0(X)}$, where $Z[1]^\circ \rightarrow \mathbf{A}^1$ is the tautological projection and $T_{Z[1]^\circ/\mathbf{A}^1}$ is the relative tangent bundle. We let $f : \mathcal{X} \rightarrow Z[1]^\circ$ be the family over $\mathfrak{M}_{1,1} \times Z[1]^\circ$ so that $\mathcal{X}$ is the pull back of the universal family over $\mathfrak{M}_{1,1}$ while the morphism f is the composite of the projection $\mathcal{X} \rightarrow \mathfrak{M}_{1,1} \times Z[1]^\circ$ with the second projection $\mathfrak{M}_{1,1} \times Z[1]^\circ \rightarrow Z[1]^\circ$. Clearly, f is the universal family of $\mathfrak{M}(Z[1]^\circ, \Gamma)^{\text{st}}$. The obstruction bundle to the moduli space $\mathfrak{M}(Z[1]^\circ, \Gamma)^{\text{st}}$ over $\mathfrak{M}_{1,1} \times Z[1]^\circ$ is $p_2^* T_{Z[1]^\circ/\mathbf{A}^1}$, where p_2 is the second projection of $\mathfrak{M}_{1,1} \times Z[1]^\circ$. The $\mathbb{C}^*$ -action lifts canonically to $p_2^* T_{Z[1]^\circ/\mathbf{A}^1}$ and the obstruction sheaf of $\mathfrak{M}(\mathfrak{Z}^{\text{rel}}, \Gamma)$ is the descent of $p_2^* T_{Z[1]^\circ/\mathbf{A}^1}$. It is direct to check that under the quotient map $\mathfrak{M}_{1,1} \times Z[1]^\circ/\mathbb{C}^* \cong \mathfrak{M}_{1,1} \times Z$, equivariant part $(p_2^* T_{Z[1]^\circ/\mathbf{A}^1})^{\mathbb{C}^*}$ is canonically isomorphic to $\pi_2^* \Omega_Z(\log D)^\vee$. This proves the identity (5.2).

5.2. Local and formal pre-deformable morphisms. In this part we prove that formal pre-deformable morphisms are automatically local pre-deformable.

Lemma 5.3. *Let $\hat{L} = (\mathbb{k}[[z_1, z_2]] \otimes_{\mathbb{k}[[s]]} A)^\wedge$, be as in Definition 1.2. Suppose there are units $f_1, f_2, g_1, g_2 \in \hat{L}$ and an integer $n \geq 1$ so that $f_1 f_2$ and $g_1 g_2 \in \hat{A}$ and that $z_i^m f_i = z_i^m g_i$ for $i = 1, 2$. Then $f_1 = g_1$ and $f_2 = g_2$.*

Proof. We let $r_i = f_i/g_i$. Then $z_i^m(r_i - 1) = 0$ for $i = 1$ and 2 . Recall that elements in $\hat{L}$ have unique normal form $a_0 + \sum_{i \geq 1} (a_i z_1^i + b_i z_2^i)$ for $a_i, b_i \in \hat{A}$ (cf. [Li]). By the uniqueness of the normal form, the normal form of r_1 (resp. r_2) must be of the form $r_1 = 1 + \sum_{j \geq 1} a_j z_2^j$ (resp. $r_2 = 1 + \sum_{j \geq 1} b_j z_1^j$). Then $f_1 f_2 / g_1 g_2 = r_1 r_2 \in \hat{A}$ implies that $1 + \sum a_j z_2^j = \epsilon(1 + \sum b_j z_1^j)^{-1}$ for some unit $\epsilon \in \hat{A}$, which is impossible unless all a_j and all b_j are zero. This proves the uniqueness Lemma. $\square$

Lemma 5.4. *The notion of pure contact is independent of the choice of the charts of the nodes of $\mathcal{U}/\mathcal{V}$.*

Proof. Let ϕ in (1.4) and $\tilde{\phi} : \mathbb{k}[[\tilde{z}_1, \tilde{z}_2]] \otimes_{\mathbb{k}[[s]]} A \rightarrow R$ be two charts of the nodes of $\mathcal{U}/\mathcal{V}$. Without loss of generality we can assume that the vanishing locus of $\phi(z_1)$ and $\tilde{\phi}(\tilde{z}_1)$ are identical in $\text{Spec } R$. Then by the proof of [Li, Lemma 2.9], $c_i = z_i/\tilde{z}_i \in R_S$ and hence $c_1 c_2 \in A_T$. Now let $\varphi : \mathbb{k}[[w_1, w_2]] \rightarrow R$ be of pure contact with respect to the chart ϕ . If $\varphi(w_i) = z_i^m h_i$ in R_S for $h_1, h_2 \in R_S$ satisfying $h_1 h_2 \in A_T$, then $\varphi(w_i) = \tilde{z}_i^m (c_i^m h_i)$ in R_S and $c_1^m h_1 c_2^m h_2 \in A_T$ as well. Thus φ is of pure contact with respect to $\tilde{\phi}$ as well. This proves the Lemma. $\square$

We now state and prove the following equivalence result.

Lemma 5.5. *Let the notation be as in Definition 1.4. Then φ is of pure contact if and only if it is formally of pure contact.*

Proof. Clearly, if φ is of pure m -contact, then it is so formally. We now prove the other direction. Assume φ is formally of pure m -contact. Then there are β_1 and $\beta_2 \in \hat{R}$ such that $\hat{\varphi}(w_i) = z_i^m \beta_i$, where as usual $\hat{\varphi}: \mathbb{k}[w_1, w_2] \rightarrow \hat{R}$ is the homomorphism induced by φ . We first show that there are f_1 and $f_2 \in R_S$ so that $\varphi(w_i) = z_i^m f_i$ in R_S for $i = 1$ and 2 . Let $x = \varphi(w_1) \in R$, then $\hat{x} = \hat{\varphi}(w_1) \in z_1^m \hat{R}$. Thus by [Ma, Thm 8.1], the residue class of $\bar{x} \in R_S/(z_1^m)$ is $(\bar{x})^\wedge = 0 \in (R/(z_1^m))^\wedge$. Hence $\bar{x} \in \cap_{m \geq 1} I^m R/(z_1^m)$, and by [Ma, Thm 8.9] there is an $a \in R/(z_1^m)$ satisfying $a \equiv 1 \pmod{I}$ such that $a\bar{x} = 0$. Then $a \in \mathcal{S}$ and by our assumption a is a unit in $R_S/(z_1^m)$. Hence $\bar{x} = 0$ in $R_S/(z_1^m)$. This proves that $\varphi(w_1) \in z_1^m R_S$ and hence there is an $f_1 \in R_S$ such that $\varphi(w_1) = z_1^m f_1$ in R_S . For the same reason, $\varphi(w_2) = z_2^m f_2$ in R_S for some $f_2 \in R_S$.

We next prove the following induction hypothesis: *For any non-negative integer k , there are g_1 and $g_2 \in R_S$ such that*

$$(5.3) \quad z_1^m f_1 - z_1^m g_1, \quad z_2^m f_2 - z_2^m g_2 \in s^k R_S \quad \text{and} \quad g_1 g_2 \in A_T + s^k R_S.$$

Clearly, this statement is true for $k = 0$. We now show that this statement is true for $k + 1$ if it is true for k . Let g_1 and g_2 be elements in R_S satisfying (5.3) for an integer k . To carry out the induction we need to find r_1 and $r_2 \in R_S$ so that

$$(5.4) \quad z_i^m f_i - z_i^m (g_i + r_i) \in s^{k+1} R_S \quad \text{for } i = 1, 2$$

and

$$(5.5) \quad (g_1 + r_1)(g_2 + r_2) \in A_T + s^{k+1} R_S.$$

Since φ formally is of pure m -contact, there are units η_1 and $\eta_2 \in \hat{R}_S$ such that $z_i^m \hat{f}_i = z_i^m \eta_i$ for $i = 1$ and 2 and $\eta_1 \eta_2 \in \hat{A}$. Because $\hat{g}_i$ and η_i satisfy the relation

$$z_i^m \hat{g}_i = z_i^m \eta_i \pmod{s^k} \quad \text{and} \quad \hat{g}_1 \hat{g}_2, \eta_1 \eta_2 \in \hat{A} + s^k \hat{R},$$

by the uniqueness Lemma 5.3, $\hat{g}_i \equiv \eta_i \pmod{s^k}$. Hence

$$(z_i^m f_i - z_i^m g_i)^\wedge = z_i^m \hat{f}_i - z_i^m \hat{g}_i = z_i^m (\eta_i - \hat{g}_i) \in z_i^m s^k \hat{R}.$$

As we argued in the existence of f_i , this implies that $z_i^m (f_i - g_i) \in z_i^m s^k R_S$.

We now let $l_i \in R_S$ be such that $z_i^m f_i - z_i^m g_i = z_i^m s^k l_i$. We let $r_1 = s^k (l_1 + z_2 h_1)$ and $r_2 = s^k (l_2 + z_1 h_2)$ with h_1 and $h_2 \in R_S$ to be determined. Clearly, (5.4) will hold with this choice of r_1 and r_2 . As for (5.5), we need

$$(g_1 + r_1)(g_2 + r_2) = g_1 g_2 + s^k (g_1 l_2 + g_1 z_1 h_1 + g_2 l_1 + g_2 z_2 h_2) \in A_T + s^{k+1} R_S.$$

Since $g_1 g_2 \in A_T + s^k R_S$, we can find $c \in R_S$ so that $g_1 g_2 - cs^k \in A_T + s^{k+1} R_S$. Hence we need to find h_1 and h_2 in R_S so that

$$(c + g_1 l_2 + g_2 l_1) + g_1 h_1 z_1 + g_2 h_2 z_2 \in A_T + s R_S.$$

Let $d = c + g_1 l_2 + g_2 l_1$. Clearly, there are γ_1 and $\gamma_2 \in \hat{R}$ and $\alpha \in A_T$ so that $\hat{d} - (\alpha + \gamma_1 z_1 + \gamma_2 z_2) \in s R_S$. Hence following the argument for the existence of f_i , there are c_1 and $c_2 \in R_S$ so that $d - (c_1 z_1 + c_2 z_2) \in A_T + s R_S$. Hence the choice $h_1 = -c_1 g_1^{-1}$ and $h_2 = -c_2 g_2^{-1}$ will do the job for (5.5). Here g_1 and g_2 are units since β_1 and β_2 are units. This proves that for each k we can find g_1 and g_2 that satisfy (5.3).

Now we show that there are h_1 and $h_2 \in R_S$ as required by the Lemma. We first let $M \subset R_S$ be the set of those elements that are annihilated by some power of s . It is an ideal, and since R_S is Noetherian, there is an N so that $s^N M = 0$. We let g_1 and g_2 be the pair satisfying (5.3) for $k = N + 1$. In case $R/M = 0$ then the Lemma is already proved. Now assume $R/M \neq 0$. We consider the ring R_S/M_S . Since $\hat{\varphi}(t) = \hat{\varphi}(w_1)\hat{\varphi}(w_2) = \epsilon s^m$ in $\hat{R}$ for some unit $\epsilon \in \hat{R}$, R_S/M_S is flat over $\mathbb{k}[t]$. Now consider the homomorphism $\bar{\varphi}: \mathbb{k}[w_1, w_2] \rightarrow R_S/M_S$ induced by φ . By [Li, Prop. 2.2], it is formally of pure contact. Hence there are $\bar{h}_1$ and $\bar{h}_2 \in R_S/M_S$ so that $\bar{\varphi}(w_i) = z_i^m \bar{h}_i$ in R_S/M_S for $i = 1$ and 2 . Because R_S/M_S is flat over $\mathbb{k}[t]$, $\bar{\varphi}_1(w_1)\bar{\varphi}_2(w_2) \in R_S/M_S$ implies that $\bar{h}_1 \bar{h}_2 \in R_S/M_S$. We then apply the uniqueness Lemma 5.3 to conclude that the residue classes of $\bar{h}_i$ and

g_i in $R_S/(M_S, s^k)$ are identical. Hence we can find $h_i \in R_S$ so that its residue class in R_S/M_S and $R_S/(s^k)$ are $\bar{h}_i$ and g_i respectively. Therefore we have $\varphi(w_i) = z_i^m h_i$ in R_S and $h_1 h_2 \in A_T$. This completes the proof of the Lemma. $\square$

REFERENCES

- [Art] M. Artin, *Versal deformations and algebraic stacks*, Invent. Math. 27 (1974), 165–189.
- [Beh] K. Behrend, *Gromov-Witten invariants in algebraic geometry*, Invent. Math. **127**(1997), no. 3, 601–617.
- [BF] K. Behrend and B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128**(1997), no. 1, 45–88.
- [DM] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, Publ. I.H.E.S. **45**(1969), 101–145.
- [EGH] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to symplectic field theory*, preprint.
- [FO] K. Fukaya and K. Ono, *Arnold conjecture and Gromov-Witten invariant for general symplectic manifolds*, The Arnoldfest (Toronto, ON, 1997), 173–190, Fields Inst. Commun., 24, Amer. Math. Soc., Providence, RI, 1999.
- [Ful] W. Fulton, *Intersection theory*, Ergebnisse der Math. und ihrer Grenzgebiete 3. Folge Band 2, 1984.
- [Gat] A. Gathmann, *Absolute and relative Gromov-Witten invariants of very ample hypersurfaces*, preprint, math.AG/9908054.
- [Gie] D. Gieseker, *A degeneration of the moduli space of stable bundles*, J. Differential Geom. **19**(1984), no. 1, 173–206.
- [GL] D. Gieseker, and J. Li, *Irreducibility of moduli of rank two vector bundles*, J. Diff. Geom. **40**(1994), 23–104.
- [GM] D. Gieseker and I. Morrison, *Hilbert stability of rank-two bundles on curves*, J. Differential Geom. **19**(1984), no. 1, 1–29.
- [Har] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics 52. Berlin, Heidelberg, New York, Springer, 1977.
- [HM] J. Harris and D. Mumford, *The Kodaira dimension of the moduli space of curves of genus ≥ 23* , Invent. Math. **90**(1987), no. 2, 359–387.
- [IP1] E-N. Ionel and T. Parker, *Gromov-Witten Invariants of Symplectic Sums*, Math. Res. Lett. **5**(1998), no. 5, 563–576.
- [IP2] E-N. Ionel and T. Parker, *Relative Gromov-Witten Invariants*, preprint, math.SG/9907155.
- [IP3] E-N. Ionel and T. Parker, *The Symplectic Sum Formula for Gromov-Witten Invariants*, preprint, math.SG/0010217.
- [Ka1] F. Kato, *Log smooth deformation theory*, Tohoku Math. J. (2) **48** (1996), no. 3, 317–354.
- [Ka2] K. Kato, *Logarithmic structures of Fontaine-Illusie*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), 191–224, Johns Hopkins Univ. Press, Baltimore, MD, 1989.
- [Kol] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics 32. Springer-Verlag, Berlin, 1996.
- [Kr1] A. Kresch, *Cycle groups for Artin stacks*, Invent. Math. **138** (1999), no. 3, 495–536.
- [Kr2] A. Kresch, *Canonical rational equivalence of intersections of divisors*, Invent. Math. **136** (1999), no. 3, 483–496.
- [KKP] B. Kim, A. Kresch, and T. Pantev, *Functoriality in intersection theory and a conjecture of Cox, Katz and Lee*, preprint.
- [Li] J. Li, *Stable morphisms to singular schemes and relative stable morphisms*, to appear in J. Diff. Geom., AG/0009097.
- [LR] A-M. Li and Y. Ruan, *Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds I*, Invent. Math. **145** (2001), no. 1, 151–218.
- [LT1] J. Li and G. Tian, *Algebraic and symplectic geometry of virtual moduli cycles*, Algebraic geometry—Santa Cruz 1995, 143–170, Proc. Sympos. Pure Math., 62, Part 2, Amer. Math. Soc., Providence, RI, 1997.
- [LT2] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*, J. Amer. Math. **11**(1998), no. 1, 119–174.

- [LT3] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds*, Topics in symplectic 4-manifolds (Irvine, CA, 1996), 47–83, First Int. Press Lect. Ser., I, Internat. Press, Cambridge, MA, 199.
- [LT4] J. Li and G. Tian, *Comparison of algebraic and symplectic Gromov-Witten invariants*, Asian J. Math. **3**(1999), no. 3, 689–728.
- [Ma] Matsumura, *Commutative ring theory*, Cambridge University press, 1986.
- [Mil] J. Milner, *Étale cohomology*, Princeton University press, Princeton, New Jersey, 1980.
- [RT] Y.B. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J. Diff. Geom. **42**(1995), no. 2, 259–367.
- [Ru1] Y. Ruan, *Topological Sigma model and Donaldson type invariants in Gromov theory*, Duke Math. J. **83**(1996), no. 2, 461–500.
- [Ru2] Y. Ruan, *Virtual neighborhoods and pseudo-holomorphic curves*, Proceedings of 6th Gökova Geometry-Topology Conference, Turkish J. Math. **23**(1999), no. 1, 161–231.
- [Si1] B. Siebert, *Symplectic Gromov-Witten invariants*, New trends in algebraic geometry (Warwick, 1996), 375–424, London Math. Soc. Lecture Note Ser., 264, Cambridge Univ. Press, Cambridge, 1999.
- [Si2] B. Siebert, *Algebraic and symplectic Gromov-Witten invariants coincide*, Ann. Inst. Fourier (Grenoble) **49**(1999), no. 6, 1743–1795.
- [Shl] M. Shlesinger, *Functors of Artin rings*, Trans. Amer. Math. Soc., **130**(1968), 208–222.
- [Tia] G. Tian, *Quantum cohomology and its associativity*, Current developments in mathematics, 1995 (Cambridge, MA), 361–401, Internat. Press, Cambridge, MA, 1994.
- [Vis] A. Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*, Invent. Math. **97**(1989), no. 3, 613–670.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305
E-mail address: `jli@math.stanford.edu`